

ETG Turbulence Isotropization

Stefan Tirkas¹
Hoatian Chen¹ Gabriele Merlo² Scott Parker¹

¹CIPS, University of Colorado, Boulder

²University of Texas, Austin

October 19, 2020

Outline

- 1 Drift Wave Instabilities
- 2 Hasegawa-Mima Fluid Model
- 3 Zonal Flow Excitation

Drift Wave Instabilities

- Drift waves are most simply characterized as density, temperature and electrostatic potential fluctuations in low- β plasmas.
- Modes relevant to tokamak physics include ion-temperature-gradient modes (ITG), electron-temperature-gradient modes (ETG), and trapped electron modes (TEM).
- Low-frequency drift wave turbulence is largely responsible for the anomalous transport of plasma particles across magnetic field lines.



Ion-Temperature-Gradient Mode Growth







Figure: Simple picture of ITG instability.

ETG Simulation in GENE



- Successful reproduction of Goerler benchmark with kinetic ions and electrons, showing ITG mode and trapped electron mode growth.
- Conversion of ITG to ETG flux-tube case prior to running non-linear simulations.

ETG "Streamers" in GENE



- ETG turbulence in toroidal gyrokinetic simulations is associated with elongated "streamers".
- Multiscale turbulence simulations have shown that streamers dominate electron heat flux and lead best reproduced experimental heat fluxes within experimental uncertainties.

Hasegawa-Mima Fluid ETG Model

- Partial differential equation derived from fluid continuity and momentum equations.
- Approximations made that are useful to describing turbulence in tokamak plasmas.
 - Cyclotron motion periods much smaller than time scales that quantities of interest change on (B, Φ, n) .
 - Long length scales along \hat{b} -direction - k_{\parallel} ignorable.
 - Quasi-neutrality of particle densities is enforced.
 - Isothermal equation of state, with adiabatic ions that have negligible temperatures.
- Shown to cause isotropic behavior for long wavelength modes as well as an inverse energy-cascade.

Hasegawa-Mima Equations

We start with the fluid continuity and momentum equations and $\tau = T_e/T_i$, where we have already taken the ion approximations discussed on the previous slide:

$$\frac{\partial n_e}{\partial t} + \nabla \cdot (n_e \vec{v}_e) = 0, \quad (1)$$

$$m_e \frac{d\vec{v}_e}{dt} = (1 + \tau) e \nabla \delta \Phi - \frac{e}{c} \vec{v}_e \times \vec{B} - \frac{\nabla P_e}{n_e}. \quad (2)$$

We break equation (2) up into parallel and perpendicular components, and break up \vec{v}_e in terms of higher and lower order terms to find,

$$\begin{aligned} \vec{v}_{e,0} &= \vec{v}_{\parallel} + \vec{v}_{\perp,0} = \vec{v}_{\parallel} + (1 + \tau) \vec{v}_E + \vec{v}_D \\ \vec{v}_{e,1} &= \vec{v}_{\perp,1} \end{aligned} \quad (3)$$

Hasegawa-Mima Equations

Taking the standard electron dynamic normalization,

$$\begin{aligned} \Phi &= \frac{e\delta\Phi}{T_i}, \quad -\frac{1}{r_n} = \frac{\partial_x n_e}{n_e}, \quad -\frac{1}{r_t} = \frac{\partial_x T_e}{T_e}, \quad \eta_e = \frac{r_n}{r_t}, \\ \rho_e &= \sqrt{\frac{\tau m_e}{m_i}} \rho_i, \quad \vec{x} = \frac{\vec{X}}{\rho_e}, \quad t = \frac{\rho_e}{r_n} \omega_{ce} t \end{aligned} \quad (4)$$

and plugging into equation () gives the form of the H-M ETG model,

$$\begin{aligned} & - \left(1 - \frac{1+\tau}{2\tau} \nabla_{\perp}^2\right) \partial_t \Phi + \frac{1+\tau}{2\tau} \frac{r_n^2}{\rho_e^2} \partial_t^{-1} \nabla_{\parallel}^2 \Phi + \frac{(1+\tau)(1+\eta_e)}{4\tau} \partial_y \nabla_{\perp}^2 \Phi \\ & + \frac{1+\eta_e}{2\tau} \partial_y \Phi + \frac{(1+\tau)^2}{\tau^2} \frac{r_n}{4\rho_e} (\hat{b} \times \nabla_{\perp} \Phi \cdot \nabla_{\perp}) \nabla_{\perp}^2 \Phi = 0. \end{aligned} \quad (5)$$

Hasegawa-Mima Equations

Finally we drop the parallel gradient term since $k_{\parallel}^2/k_{\perp}^2 \sim \epsilon^2$, and simplify the bracketed expression for a 2-D slab geometry to find the final form of our model,





$$\begin{aligned} \partial_t \left[\Phi - \frac{1+\tau}{2\tau} \zeta \right] = & \frac{(1+\tau)(1+\eta_e)}{4\tau} \zeta_y + \frac{1+\eta_e}{2\tau} \phi_y \\ & + \frac{(1+\tau)^2}{\tau} \frac{r_n}{4\rho_e} [\Phi_x \zeta_y - \zeta_x \Phi_y], \end{aligned} \quad (6)$$

where $\zeta = \nabla^2 \Phi$ and x, y subscripts denote partial derivatives.

Pseudo-Spectral Solver


- Equation (6) is solved numerically using the pseudo-spectral method.
 - Fourier transform the equation and get $\zeta = (k_x^2 + k_y^2)\Phi$.
 - Inverse Fourier transform $\zeta_{x,y}$ and $\Phi_{x,y}$ back into real space.
 - Calculate the non-linear products between ζ and Φ in real space and then Fourier transform the products so they can be added to the other terms in Fourier space.
 - Time advance Φ discretely.
- Time advancement is done using the 4th order Runge-Kutta method.
- The nonlinear term, due to the polarization drift, is responsible for isotropizing the turbulence.

ETG H-M Results



- Results of H-M code. Initial condition (above) isotropizes at later time (below).
- Add parameters?

GENE ETG Streamer Test



- Test of GENE saturated ETG results into H-M code. Initial conditions (above) isotropize at later times (below).
- Inverse energy cascade observed.

Zonal Flow Excitation




Figure: Figure shows a non-linear GENE sim initialized with two modes - a default ZF mode ($k_y = 0$) and the orange ETG ($k_y \rho_i = 6.36$) mode. This excites the blue ZF mode ($k_y = 0, k_x \rho_i = 4.955$).

- Zonal flow can be spontaneously excited by intermediate-scale ETG turbulence in tokamak geometries - $k_{\perp} \rho_e \ll 1 \ll k_{\perp} \rho_i$.
- Plan to carry out large-scale ETG simulations with GENE to compare to theory of zonal flow excitation by ETG modes.

References

