

Scalar and Representative Observables for Polarimetric SAR Data

Thanh-Hai Le, Ian McLoughlin, and Chan-Hua Vun

Abstract

This paper proposes a scalar and representative observable for the multi-dimensional POLSAR data, from which statistically consistent discrimination measures can be derived. Specifically, the statistical behaviour of the POLSAR covariance matrix determinant is used to derive a scalar and generic statistical model for the multi-dimensional POLSAR data, which is specifically applicable to the two and three dimensional versions of partial and full monostatic polarimetric SAR data. As the POLSAR covariance matrix determinant generalizes the SAR intensity towards the multidimensional extension, the proposed model is also shown to include the traditional model used for SAR intensity under its natural coverage. Consequently, the main beneficial implication of the proposed approach is that it provides a consistent theory unifying the currently disconnected proposals for SAR and POLSAR discrimination measures which enables the adaptation of many existing SAR data processing techniques for POLSAR data.

Index Terms

Polarimetric Synthetic Aperture Radar, Electromagnetic Modeling, Multidimensional Signal Processing

I. INTRODUCTION

During the past decades, exponential growth in computing power has allowed the once computationally-demanding Synthetic Aperture Radar (SAR) technology to become a feasible and preferred technique for earth observation applications. SAR technology has since been extended in a few directions, one of which is the polarimetric SAR (POLSAR). POLSAR extends SAR by

Thanh-Hai Le and Chan-Hua Vun are with School of Computer Engineering, Nanyang Technological University, Singapore. Ian McLoughlin is with School of Information Science and Technology, University of Science and Technology of China.

Manuscript received ?, 2013; revised ?.

exploiting the natural polarization property of Electro-Magnetic (EM) waves, leading to the availability of multi-channel POLSAR data, compared to traditional one-channel SAR data.

Similar to the SAR data, the POLSAR data is stochastic. Moreover, it is multi-dimensional making it even harder to interpret. Under this context, it is therefore important to establish a simple and intuitive understanding of the data and statistical models are undoubtedly crucial in understand the stochastic its nature. While several multidimensional models have been proposed for the POLSAR data, unfortunately, due to the multidimensional nature, they are complex and not very intuitive. Practical POLSAR data processing, however, makes heavily use of scalar discrimination measures, which should be based on statistically consistent models for scalar and representative observables of the multidimensional data. It is thus important to establish a scalar and representative observable for this multi-dimensional POLSAR data.

There has been a few scalar observables with accompanying statistical models proposal for POLSAR [1–7], but none of them is able to provide meaningful scalar discrimination measures. As such no observable has been widely accepted as being highly representative of the multi-dimensional data, which severely limits their applicability in practical data processing applications. From an alternative approach, there are also a few POLSAR discrimination measures proposed, but all of them are based on the likelihood ratio statistics. This statistical test should be based on an exact and consistent statistical distribution but so far has only shown to be based on an asymptotic distribution.

This article hence present a scalar and representative observable, and its associated generic statistical model, for the multi-dimensional POLSAR data that provides a consistent foundation for the derivation of POLSAR discrimination measures. The POLSAR data have been statistically modelled as following the complex Wishart distribution. Consequently, the generic statistical model for the covariance matrix determinant, being just a scalar projection of the multidimensional data, is presented. This model is then used to derive several scalar and consistent statistical models suggesting that their associated observables are capable of being used as discrimination measures for POLSAR data. The specific two and three dimensional models of these models, which also include the derived discrimination measures, are originally designed against both partial and full polarimetric SAR data.

Interestingly the paper also shows that the specific one dimensional versions of these models proposed match perfectly with the traditional statistical model used for SAR intensity. This

effectively puts the one dimensional SAR theory to stay within the proposed scalar approach for multi dimensional POLSAR. With this new insight the different discrimination proposals for SAR and POLSAR are reviewed and it is shown that the approach proposed in this paper provides a strong, unifying and consistent foundation relating them all together. The applicability of these theoretical models is illustrated by experiments where the specific one, two and three dimensional versions of the proposed generic models are validated against practical data.

Thus, the rest of this article is organized as follows. After the next section reviews existing discrimination measures as well as scalar observables models for POLSAR, section III derives the generic statistical model for the determinant of the POLSAR covariance matrix and proposes several new discrimination measures for the multi-dimensional data. After section IV illustrates the match between the specific model of one dimensional and the traditional model for SAR intensity, section V links the disparate discrimination measure proposals for the traditional SAR and the more recent POLSAR data. The applicability of these models against practical data is illustrated in section VI. Section VII presents a high-level discussion of the results presented before section VIII finally concludes this paper.

II. RELATED WORK IN LITERATURE

Section II-A reviews various statistical models used for different scalar observables for POLSAR, where it is shown that none of these is able to provide statistically consistent discrimination measures. Section II-B further strengthens those findings by discussing discrimination measures that have been proposed for POLSAR, and it demonstrates that almost all of them are based on the likelihood statistical test for complex Wishart distribution. While an exact statistical distribution is expected for the test, only asymptotic distribution is used in the original work [1].

A. *Scalar Observables for POLSAR Data and their Statistical Models*

Different target decomposition theorems have identified many possible scalar observables for complex POLSAR data. In [2], the performance of different scalar POLSAR observables is evaluated for classification purposes. While many scalar observables for POLSAR were presented, their corresponding statistical models and classifiers were not available. Furthermore, at its conclusion, the paper indicated that it is impossible to identify a single best representation.

Although, to be fair, the observables were identified for describing a decomposed portion of the complex POLSAR data, rather than providing a unified representation of the POLSAR data.

Using a different approach, given that the joint distribution for POLSAR is known to be the multi-variate complex Wishart distribution, it is possible to derive the scalar statistical models for some univariate POLSAR observables. However, such derivations are not trivial tasks, and so far, only a handful of such statistical models have been proposed, including the following:

- 1) cross-pol ratio $r_{HV/HH} = |S_{HV}|^2/|S_{HH}|^2$ [3],
- 2) co-pol ratio $r_{VV/HH} = |S_{VV}|^2/|S_{HH}|^2$ [3],
- 3) co-pol phase difference $\phi_{VV/HH} = \arg(S_{VV}S_{HH}^*)$ [3] [4],
- 4) magnitude $g = |\text{avg}(S_{pq}S_{rs}^*)|$ [4],
- 5) normalized magnitude $\xi = \frac{|\text{avg}(S_{pq}S_{rs}^*)|}{\sqrt{\text{avg}(|S_{pq}|^2)\text{avg}(|S_{rs}|^2)}}$ [4],
- 6) intensity ratio $w = \text{avg}(|S_{pq}|^2)/\text{avg}(|S_{rs}|^2)$ [4],
- 7) and the Stokes parameters $S_i, 0 \leq i \leq 3$ [5].

More recently, statistical models for each element of the POLSAR covariance matrix, i.e. $S_{pq}S_{rs}^*$, [6] as well as for the largest eigen-value of the covariance matrix λ_1 [7] have been proposed.

While these models undoubtedly help to further our understandings of the POLSAR data, none of the underlying observables have been shown to meet the dual criteria of 1) resulting in statistically consistent discrimination measures and thus 2) being representative of the complex POLSAR data.

B. POLSAR Discrimination Measures

The commonly used measure of distance for matrices are either the Euclidean or Manhattan distances, defined respectively as:

$$d(C_x, C_y) = \sum_{i,j} |\Re(C_x - C_y)_{i,j}| + \sum_{i,j} |\Im(C_x - C_y)_{i,j}| \quad (1)$$

$$d(C_x, C_y) = \sqrt{\sum_{i,j} |C_x - C_y|_{i,j}^2} \quad (2)$$

where $C_{i,j}$ denotes the (i,j) elements of the POLSAR covariance matrix C , $||$ denotes absolute values and \Re, \Im denote the real and imaginary parts respectively. However, in the context of POLSAR, these dissimilarity measures are not widely used mainly due to the multiplicative nature of the noisy data.

In the field of POLSAR, the Wishart distance is probably the most widely used, as part of the well-known Wishart classifier [8]. It is defined [9] as:

$$d(C_x, C_y) = \ln |C_y| + \text{tr}(C_x C_y^{-1}) \quad (3)$$

, where $\text{tr}(C)$ denotes the trace of the matrix C . As a measure of distance, its main disadvantage is that $d(C_y, C_y) = \ln |C_y| \neq 0$.

Recent works have suggested alternative dissimilarity measures including the symmetric and asymmetric refined Wishart distance [10],

$$d(C_x, C_y) = \frac{1}{2} \text{tr}(C_x^{-1} C_y + C_y^{-1} C_x) - d \quad (4)$$

$$d(C_x, C_y) = \ln |C_x| - \ln |C_y| + \text{tr}(C_x C_y^{-1}) - d \quad (5)$$

the Bartlett distance [11],

$$d(C_x, C_y) = 2 \ln |C_{x+y}| - \ln |C_x| - \ln |C_y| - 2d \ln 2 \quad (6)$$

the Bhattacharyya distance [12],

$$r(C_x, C_y) = \frac{|C_x|^{1/2} |C_y|^{1/2}}{|(C_x + C_y)/2|} \quad (7)$$

and the Wishart Statistical test distance [13]

$$d(C_x, C_y) = (L_x + L_y) \ln |C| - L_x \ln |C_x| - L_y \ln |C_y| \quad (8)$$

Closer examination of these dissimilarity measures reveals that most of them are related in some ways. The Bhattacharyya distance is easily shown to be related to the Barlett distance. At the same time the Barlett distance can be considered a special case of the Wishart Statistical Test distance, when the two data sets have the same number of looks, i.e. $L_x = L_y$. The close relation among the measures is further supported by the fact that all of their publications referred the same statistical model developed in [1] as the foundation. In [1], to determine if the two scaled multi-look POLSAR covariance matrixes Z_x and Z_y , which have L_x and L_y as the corresponding number of looks, come from the same underlying stochastic process, the likelihood ratio statistics for POLSAR covariance matrix is considered:

$$Q = \frac{(L_x + L_y)^{d \cdot (L_x + L_y)}}{L_x^{d \cdot L_x} L_y^{d \cdot L_y}} \frac{|Z_x|^{L_x} |Z_y|^{L_y}}{|Z_x + Z_y|^{(L_x + L_y)}}$$

Taking the log-transformation of the above equation, and denote $C_{vx} = Z_x/L_x$, $C_{vy} = Z_y/L_y$ and $C_{vxy} = (Z_x + Z_y)/(L_x + L_y)$ then:

$$Q = \frac{|C_{vx}|^{L_x} \cdot |C_{vy}|^{L_y}}{|C_{vxy}|^{L_x+L_y}} \quad (9)$$

$$\ln Q = L_x \ln |C_{vx}| + L_y \ln |C_{vy}| - (L_x + L_y) \ln |C_{vxy}| \quad (10)$$

To detect changes, a test statistic is developed for this discrimination measure. This means a distribution is to be derived for the dissimilarity measure. However, in the original work [1], only an asymptotic distribution was used. This paper proposes a statistical model for the determinant of the POLSAR covariance matrix $|C_v|$ which is capable of providing an exact distribution for the test.

III. THE GENERIC SCALAR STATISTICAL MODEL FOR POLSAR

In this section, after the basic foundations of POLSAR statistical analysis is introduced, the generic scalar statistical model for the multi-dimensional POLSAR is presented.

In this paper, the POLSAR scattering vector is denoted as s . For partial polarimetric SAR (single polarization in transmit and dual polarization in receipt), the vector is two-dimensional ($d = 2$) and is normally written as:

$$s_{part} = \begin{bmatrix} S_h \\ S_v \end{bmatrix} \quad (11)$$

For full and monostatic POLSAR data, the vector is three-dimensional ($d = 3$) and is presented as:

$$s_{full} = \begin{bmatrix} S_{hh} \\ \sqrt{2}S_{hv} \\ S_{vv} \end{bmatrix} \quad (12)$$

Let $\Sigma = E[ss^{*T}]$ denote the population expected value of the POLSAR covariance matrix, where s^{*T} is the complex conjugate transpose of s . Assuming s is jointly circular complex Gaussian with the expected covariance matrix Σ , then the probability density function (PDF) of s can be written as:

$$pdf(s; \Sigma) = \frac{1}{\pi^d |\Sigma|} e^{-s^{*T} \Sigma^{-1} s} \quad (13)$$

where $||$ denotes the matrix determinant.

The sample POLSAR covariance matrix is formed as the mean of Hermitian outer product of independent single-look scattering vectors,

$$C_v = \langle s s^{*T} \rangle = \frac{1}{L} \sum_{i=1}^L s_i s_i^{*T} \quad (14)$$

where L is the number of looks and s_i denotes the single-look scattering vector, which equals s_{part} for the case of partial POLSAR or s_{full} for the case of full polarimetry.

Complex Wishart distribution statistics are normally used for the scaled covariance matrix $Z = LC_v$, whose PDF is given as:

$$pdf(Z; d, \Sigma, L) = \frac{|Z|^{L-d}}{|\Sigma|^L \Gamma_d(L)} e^{-tr(\Sigma^{-1}Z)} \quad (15)$$

with $\Gamma_d(L) = \pi^{d(d-1)/2} \prod_{i=0}^{d-1} \Gamma(L-i)$ and d is the dimensional number of the POLSAR covariance matrix.

The approach taken in this paper differs by applying the homoskedastic log transformation on a less-than-well-known result for the determinant of the covariance matrix. Goodman [14] proved that the ratio between the observable and expected values of the sample covariance matrix determinants behaves like a product of d chi-squared random variables with different degrees of freedom:

$$\chi_L^d = (2L)^d \frac{|C_v|}{|\Sigma_v|} \sim \prod_{i=0}^{d-1} \chi(2L-2i) \quad (16)$$

This result is used here to develop the generic scalar statistical model for POLSAR. From Eqn. 16 we have:

$$|C_v| \sim |\Sigma_v| \cdot \frac{1}{(2L)^d} \cdot \prod_{i=0}^{d-1} \chi(2L-2i) \quad (17)$$

In a given homogeneous POLSAR area, the parameters Σ_v , d and L can be considered constant. Thus Eqn. 17 indicates that in the original POLSAR domain, a multiplicative speckle noise pattern is present.

Moreover, since the average and variance of these chi-squared distributions are known to be constant, i.e. $avg[\chi(2L)] = 2L$ and $var[\chi(2L)] = 4L$, the product and summation of these random variables also has fixed summary statistics. Specifically:

$$\begin{aligned} avg \left[\prod_{i=0}^{d-1} \chi(2L-2i) \right] &= 2^d \cdot \prod_{i=0}^{d-1} (L-i), \\ var \left[\prod_{i=0}^{d-1} \chi(2L-2i) \right] &= \prod_{i=0}^{d-1} 4(L-i)(L-i+1) - \prod_{i=0}^{d-1} 4(L-i)^2, \end{aligned}$$

Combining these results with Eqn. 17 , we have:

$$avg [|C_v|] = \frac{|\Sigma_v|}{L^d} \prod_{i=0}^{d-1} (L - i) \quad (18)$$

$$var [|C_v|] = \frac{|\Sigma_v|^2 \left[\prod_{i=0}^{d-1} (L - i)(L - i + 1) - \prod_{i=0}^{d-1} (L - i)^2 \right]}{L^{2d}} \quad (19)$$

For a real world captured image, while the parameters d and L do not change for the whole image, the underlying Σ_v is expected to differ from one region to the next. Thus over a heterogeneous scene, the stochastic process for $|C_v|$ and $\ln |C_v|$ vary depending on the underlying signal Σ_v . In such context, Eqn. 19 implies that the variance of $|C_v|$ also differs depending on the underlying signal Σ_v (i.e. it is heteroskedastic).

Similar to the way intensity-ratio is proposed as the discrimination measure for the multiplicative and heteroskedastic SAR intensity [15], in this paper, the determinant-ratio and the change-ratio are also proposed as a discrimination measure for the POLSAR data. which is shown above to also suffer from the multiplicative and heteroskedastic phenomena.

For cases where the true value of the underlying signal Σ_v is known *a priori*, then the determinant-ratio of the signal random variable (\mathbb{R}_Σ) is defined as:

$$\mathbb{R}_\Sigma = \frac{|C_v|}{|\Sigma_v|} \quad (20)$$

For cases where the POLSAR data is known to have come from a homogeneous area, but the true value of the underlying signal Σ_v is *unknown*, then random variable called the change-ratio (\mathbb{R}_C) is defined as:

$$\mathbb{R}_C = \frac{|C_1|}{|C_2|} \quad (21)$$

where C_1 and C_2 are samples of the covariance matrix determinant in an assumed homogeneous area.

Using the results from Eqn. 17, we have

$$\mathbb{R}_\Sigma \sim \frac{1}{(2L)^d} \cdot \prod_{i=0}^{d-1} \chi(2L - 2i) \quad (22)$$

$$\mathbb{R}_C \sim \prod_{i=0}^{d-1} \frac{\chi(2L - 2i)}{\chi(2L - 2i)} \quad (23)$$

Since each elementary component follows fixed distributions (i.e. $\chi^2(2L)$), it is natural that this variable also follows fixed distributions. Moreover, it is independent of the underlying signal Σ_v , indicating its statistically consistent properties and hence its applicability as a POLSAR discrimination measure. This claim is extrapolated from the widely-used intensity-ratio as SAR discrimination measure and will be further discussed in the next sections.

IV. SAR AS A ONE-DIMENSIONAL CASE OF POLSAR

This section shows that the proposed generic model is equally applicable to the 1-dimensional ($d = 1$) case, which is physically equivalent to collapsing the multi-dimensional POLSAR dataset into conventional single dimensional SAR data. Mathematically, the sample covariance matrix C_v is reduced to the sample variance while the determinant $|C_v|$ is equivalent to the scalar variance value. As variance is equal to intensity I for SAR data, our result is consistent with previous results for SAR intensity data. Hence this derivation can be considered both as cross-validation for the proposed POLSAR model as well as evidence showing that the proposed generic model for POLSAR also includes the traditional model for SAR intensity in its natural coverage.

The results for our models can be summarized using the following equations:

$$\begin{aligned} |C_v| &\sim |\Sigma_v| \cdot \frac{1}{(2L)^d} \cdot \prod_{i=0}^{d-1} \chi(2L - 2i) \\ \mathbb{R}_\Sigma &= \frac{|C_v|}{|\Sigma_v|} \sim \frac{1}{(2L)^d} \prod_{i=0}^{d-1} \chi(2L - 2i) \\ \mathbb{R}_C &= \frac{C_1}{C_2} \sim \prod_{i=0}^{d-1} \frac{\chi(2L - 2i)}{\chi(2L - 2i)} \end{aligned}$$

Upon setting $d = 1$ into the above equations, the equations become:

$$\begin{aligned} |C_v| &\sim \frac{|\Sigma_v|}{(2L)} \cdot \chi(2L) \\ \mathbb{R}_\Sigma &= \frac{|C_v|}{|\Sigma_v|} \sim \frac{1}{(2L)} \cdot \chi(2L) \\ \mathbb{R}_C &= \frac{C_1}{C_2} \sim \prod_{i=0}^{d-1} \frac{\chi(2L - 2i)}{\chi(2L - 2i)} \end{aligned}$$

Since the PDF of chi-squared distribution can be written as:

$$\chi(2L) \sim pdf \left[\frac{\chi^{L-1} e^{-\chi/2}}{2^L \Gamma(L)} \right]$$

, applying variable change theorem into the above equations results in:

$$\begin{aligned} |C_v| &\sim pdf \left[\frac{L^L x^{L-1} e^{-Lx/|\Sigma_v|}}{\Gamma(L)|\Sigma_v|^L} \right] \\ \mathbb{R}_\Sigma = \frac{|C_v|}{|\Sigma_v|} &\sim pdf \left[\frac{L^L x^{L-1} e^{-Lx}}{\Gamma(L)} \right] \\ \mathbb{R}_C = \frac{C_1}{C_2} &\sim pdf \left[\frac{\Gamma(2L-1)x^{L-1}}{\Gamma^2(L-1)(1+x)^{2L}} \right] \end{aligned}$$

These equations match exactly with the following traditional model for multi-look SAR intensity:

$$I \sim pdf \left[\frac{L^L x^{L-1} e^{-Lx/\bar{I}}}{\Gamma(L)\bar{I}^L} \right] \quad (24)$$

$$\mathbb{R}_{\bar{I}} = \frac{I}{\bar{I}} \sim pdf \left[\frac{L^L x^{L-1} e^{-Lx}}{\Gamma(L)} \right] \quad (25)$$

$$\mathbb{R}_I = \frac{I_1}{I_2} \sim pdf \left[\frac{\Gamma(2L-1)x^{L-1}}{\Gamma^2(L-1)(1+x)^{2L}} \right] \quad (26)$$

considering that $|C_v| \mapsto I$ and $|\Sigma_v| \mapsto \bar{I}$ as multi-dimensional POLSAR collapses into single-dimensional SAR.

V. UNIFYING THE DIFFERENT DISCRIMINATION MEASURES PROPOSALS FOR BOTH SAR AND POLSAR

Statistical models have always been the foundation to derive discrimination measures. This is true for both fields of SAR and POLSAR. For SAR, where the field is more or less settled on the issue, the statistical model for SAR intensity has been used to derive the most widely used intensity-ratio discrimination measure [15]. For POLSAR, where the field is much less matured, the same case should apply, except that so far only asymptotic distribution has been derived for the most common foundation, i.e. the likelihood test statistics [1].

With the insight gained from section IV, this section presents a few results. First, similar to the way that the statistical model for SAR intensity has been used to provide the foundation for SAR discrimination measures, e.g. intensity-ratio [15], the statistical model proposed for POLSAR covariance matrix determinant is to be reviewed as providing the foundation for POLSAR discrimination measures, i.e. the likelihood test statistics. And second, new discrimination measures for POLSAR may be derived by learning from the existing SAR discrimination measures.

As for the first matter, in view of the models given in Eqn 17, the likelihood test statistics presented in [1] and rewritten in Eqns 9 & 10 can be expressed as:

$$\ln Q \sim k + L_x \Lambda_{L_x}^d + L_y \Lambda_{L_y}^d - (L_x + L_y) \Lambda_{(L_x+L_y)}^d$$

$$Q \sim e^k \frac{(\chi_{L_x}^d)^{L_x} \cdot (\chi_{L_y}^d)^{L_y}}{(\chi_{L_x+L_y}^d)^{L_x+L_y}}$$

where $k = d[(L_x + L_y) \ln(L_x + L_y) - L_x \ln L_x - L_y \ln L_y]$. This, in essence, derives an exact statistical distribution for the likelihood test statistics, as opposed to the asymptotic distribution derived in [1].

As a by-product of this exact derivation, several discrimination measures for the common case of $L_x = L_y$ are further proposed. They are the determinant-ratio and the change-ratio presented in Section III. Compared to existing discrimination measures for POLSAR reviewed in Section II, the proposed dissimilarity measures are simpler in both concept and computation. They, at the same time, are multi-dimensional extensions of the intensity-ratio discrimination measure widely used in SAR.

In short, with the newly gained insight, the approach proposed in this paper provides a new bridge between the two fields of SAR and POLSAR. This invites the adaptation of many established SAR data processing techniques towards POLSAR data, where this Section has just described a few examples. However, before further discussion ensues, it is necessary to validate the presented theoretical model with real-life practical data.

VI. VALIDATING THE PROPOSED MODELS AGAINST REAL-LIFE DATA

This section aims to verify the theoretical models given in Eqns. 17, 22 and 23 against practical data. All of these models require the estimation of two parameters from the captured data. These are (i) the dimensional number d and (ii) the look number L . The dimensional number d is related to the type of (POL)SAR data captured, with $d = 1, 2, 3$ corresponding to the cases of SAR, partial and full polarimetric SAR data, respectively. The look-number L can be taken either as nominally stated by the data provider, or it can be estimated using the technique proposed by [16].

To show the robustness of the proposed models, their validations are also put to test on different types of POLSAR sensors. Specifically in this paper, the air-borne four-look ($L = 4$) AIRSAR image of Flevoland and a fine-quad single-look complex RADARSAT2 image ($L = 1$)

are used. Since the determinant of the covariance matrix is only significant on multi-look data, nine-look processing is first applied on the single-look RADARSAT2 data ($L = 9$).

A. The Traditional case of SAR ($d = 1$)

Since it has been shown in Section IV that the model for the case of $d = 1$ matches exactly with the traditional model for SAR intensity, the validation of the proposed $d = 1$ model is quite straightforward. Fig. 1 presents the results of a test performed for the stated purpose. In the test, the intensity of single-channel SAR data (HH) for a homogeneous area in both the AIRSAR Flevoland dataset is extracted. The histograms for both the intensity and intensity ratio are then plotted against the theoretical PDF given above. Also presented are the plots of the same test applied on a homogeneous patch inside the nine-look processed RADARSAT2 image. The plots are obtained with ENL set an estimated value obtained using the technique presented in [16]. In all cases, good visual match is apparent, which hence validates the model proposed above.

B. The Multi-dimensional case of POLSAR ($d = 2, 3$)

The remainder of this section now focuses on validating the models for both partial ($d = 2$) and full ($d = 3$) POLSAR. Similar tests are carried out for both types of polarimetric SAR on both AIRSAR and RADARSAT2 datasets. The look-number is estimated for each type of data and for each image using the technique described in [16]. From the plots, their visual matches again validates the proposed model.

VII. DISCUSSION

To begin with a few theoretical properties of the proposed statistical models are discussed. First, the use of covariance matrix log-determinant may be related to the standard eigen-decomposition method of the POLSAR covariance matrices. In fact, the log-determinant can also be computed as the sum of log-eigenvalues. Specifically $\ln |M| = \sum \ln \lambda_M$ where λ_M denotes all the eigenvalues of M . Thus similar to other eigenvalue based approach (e.g. entropy/anisotropy, ...), the models presented here are invariant to polarization basis transformations.

Second, the models are developed for the POLSAR covariance matrix. However, since the POLSAR coherency matrix is related to the covariance matrix via a unitary transformation which preserves the determinant, the model should also be applicable to the coherency matrix.




Fig. 1: The specific $d=1$ models are validated on both practical RADARSAT2 and AIRSAR datasets

It should be noted that the models are far from complete. While it is desirable to reduce the multi-dimensional POLSAR data to a scalar value for many applications, such a reduction is unlikely to be lossless. Thus the use of this technique could be complemented with some high-dimensional POLSAR target-decomposition techniques, such as the Freeman Durden decomposition [17] or the entropy/anisotropy decomposition [18] or similar.

Nevertheless the proposed models are promising. Even though initially developed for partial and monostatic POLSAR data, it was shown to be applicable to traditional SAR data as well. Since the models assumptions are quite minimal, they may potentially applicable to bi-static and interferometric data, although that would require further study.

Intuitively, the concept of the determinant being representative of the POLSAR data can be




Fig. 2: The specific $d=2$ models are validated on both practical RADARSAT2 and AIRSAR datasets

thought of as being similar to the concept of using the magnitude as being representative of a complex number (in a sense this is also why intensity is widely considered representative of the SAR data complex). Similar to the determinant, the magnitude observable is also scalar and is not lossless. To fully represent the complex number, another observable is needed besides the magnitude. The magnitude however is much more widely used, as it can be shown that it is invariant to a change in the reference frame. Similarly, the determinant of the POLSAR covariance matrix is also scalar and invariant to a change of polarization basis. Thus while it is not fully representative of the POLSAR data, it can be said to be highly representative of it. The important thing is that, it can be used to give a scalar comparison of the multi-dimensional data.




Fig. 3: The specific $d=3$ models are validated on both practical RADARSAT2 and AIRSAR datasets

This is also similar to the most commonly used magnitude to compare two or more complex numbers.

Besides the above properties, the theoretical models may also provide an alternative derivation for the likelihood test statistics widely used in POLSAR. Similar to the way that different measures of distance can be used to derive POLSAR classifiers [8], change detectors [1], edge detectors [19] or other clustering and speckle filtering techniques [20] [21], new detection, classification, clustering or speckle filtering algorithms can be derived using the models presented in this paper. These applications however are not included in this paper, but rather left as a

suggestion for future investigations.

VIII. CONCLUSION

In this paper, based on the commonly used complex Wishart distribution for the POLSAR target vector, a generic statistical model has been proposed for the POLSAR covariance matrix determinant. Then the specific ($d=2$) and ($d=3$) versions of the derived model are respectively validated against real-life partial and full polarimetric SAR data. The paper also establishes a few POLSAR discrimination measures: the determinant-ratio and the change-ratio, which essentially are the generic version of the SAR intensity-ratio. At the same time, it also shows a perfect match between the ($d=1$) specific models and the traditional model for SAR intensity, effectively encompassing the existing SAR theories within the proposed generic models' natural coverage.

The main result of this paper is the proposal to consider the POLSAR covariance matrix determinant as the scalar and representative observable for the multi-dimensional POLSAR data. Compared to other published scalar observables, the proposed determinant is highly representative of the multi-dimensional data. Its representative power is justified by the following reasons. Firstly, the covariance matrix determinant, when collapsed into one-dimensional case, transforms neatly into the representative SAR intensity. Secondly, the statistical model for this observable is shown to be the generic multi-dimensional extension of the traditional model for the one dimensional SAR intensity. And last but not least, this observable also leads to the establishment of the determinant-ratio discrimination measure for POLSAR data, which should increase the usability of this proposal.

There are several beneficial implications of the proposed approach presented in this paper. Currently while the field of SAR is much more developed than the field of POLSAR, the two fields however are quite separated. Hence many techniques applicable to SAR are not directly applicable to POLSAR. Since the scalar and representative model for POLSAR generalizes the traditional model for SAR intensity, its main benefit is that it enables the adaptation of many existing SAR data processing techniques for POLSAR data. One example has already been presented in this paper, where existing SAR discrimination measures are extended towards POLSAR. At the same time, it provides a consistent theory unifying the seemingly disparate discrimination measure proposals for SAR and POLSAR data.

The models proposed in this article also has its limitations. Firstly, they are based on the

complex Wishart distribution. which is only guaranteed to work for homogeneous area. Secondly, while it is desirable for a large number of applications to reduce the multi-dimensional POLSAR data to a scalar value, such a reduction is unlikely to be lossless.

In this paper, the theoretical model has only been validated for partial ($d=2$) and full ($d=3$) monostatic POLSAR data, leaving the test of its validity for different types of dataset like bistatic POLSAR data ($d=4$) or interferometric POLSAR data ($d=6$) a possible future work. Other possible future work includes an investigation on the applicability of this observables into POLSAR heterogeneous areas, as well as further studies to explore the use of these generic models in conjunction with other target decomposition techniques such as the Freeman-Durden decomposition [17] or the entropy/anisotropy decomposition [18], etc.

REFERENCES

- [1] K. Conradsen, A. Nielsen, J. Schou, and H. Skriver, "A test statistic in the complex Wishart distribution and its application to change detection in polarimetric SAR data," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 41, no. 1, pp. 4 – 19, Jan 2003.
- [2] V. Alberga, G. Satalino, and D. K. Staykova, "Comparison of polarimetric SAR observables in terms of classification performance," *International Journal of Remote Sensing*, vol. 29, no. 14, pp. 4129–4150, 2008. [Online]. Available: <http://www.tandfonline.com/doi/abs/10.1080/01431160701840182>
- [3] I. Joughin, D. Winebrenner, and D. Percival, "Probability density functions for multilook polarimetric signatures," *Geoscience and Remote Sensing, IEEE Transactions on*, vol. 32, no. 3, pp. 562 –574, may 1994.
- [4] J.-S. Lee, K. Hoppel, S. Mango, and A. Miller, "Intensity and phase statistics of multilook polarimetric and interferometric SAR imagery," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 32, no. 5, pp. 1017 –1028, Sep 1994.
- [5] R. Touzi and A. Lopes, "Statistics of the stokes parameters and of the complex coherence parameters in one-look and multilook speckle fields," *Geoscience and Remote Sensing, IEEE Transactions on*, vol. 34, no. 2, pp. 519 –531, mar 1996.
- [6] C. Lopez-Martinez and X. Fabregas, "Polarimetric SAR speckle noise model," *Geoscience and Remote Sensing, IEEE Transactions on*, vol. 41, no. 10, pp. 2232–2242, 2003.
- [7] E. Erten, "The performance analysis based on SAR sample covariance matrix," *Sensors (Basel)*, vol. 12, no. 3, pp. 2766–2786, 2012.
- [8] J.-S. Lee, M. Grunes, and G. de Grandi, "Polarimetric SAR speckle filtering and its implication for classification," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 37, no. 5, pp. 2363 –2373, Sep 1999.
- [9] J. S. Lee, M. R. Grunes, and R. Kwok, "Classification of multi-look polarimetric SAR imagery based on complex Wishart distribution," *International Journal of Remote Sensing*, vol. 15, no. 11, pp. 2299–2311, 1994.
- [10] S. N. Anfinsen, R. Jenssen, and T. Eltoft, "Spectral clustering of polarimetric SAR data with Wishart-derived distance measures," in *3rd International Workshop on Science and Applications of SAR Polarimetry and Polarimetric Interferometry*, vol. 3, Jan 2007.

- [11] P. Kersten, J. S. Lee, and T. Ainsworth, "Unsupervised classification of polarimetric synthetic aperture radar images using fuzzy clustering and EM clustering," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 43, no. 3, pp. 519 – 527, Mar 2005.
- [12] K. Y. Lee and T. Bretschneider, "Derivation of separability measures based on central complex Gaussian and Wishart distributions," in *IEEE International Geoscience and Remote Sensing Symposium (IGARSS), 2011*, Jul 2011, pp. 3740 –3743.
- [13] F. Cao, W. Hong, Y. Wu, and E. Pottier, "An unsupervised segmentation with an adaptive number of clusters using the SPAN/H/ α /A space and the complex Wishart clustering for fully polarimetric SAR data analysis," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 45, no. 11, pp. 3454 –3467, Nov 2007.
- [14] N. R. Goodman, "The distribution of the determinant of a complex Wishart distributed matrix," *Annals of Mathematical Statistics*, vol. 34, no. 1, pp. 178–180, 1963.
- [15] E. Rignot and J. van Zyl, "Change detection techniques for ERS-1 SAR data," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 31, no. 4, pp. 896 –906, Jul 1993.
- [16] S. Anfinson, A. Doulgeris, and T. Eltoft, "Estimation of the equivalent number of looks in polarimetric synthetic aperture radar imagery," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 47, no. 11, pp. 3795 –3809, Nov 2009.
- [17] A. Freeman and S. Durden, "A three-component scattering model for polarimetric SAR data," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 36, no. 3, pp. 963 –973, May 1998.
- [18] S. Cloude and E. Pottier, "An entropy based classification scheme for land applications of polarimetric SAR," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 35, no. 1, pp. 68 –78, Jan 1997.
- [19] J. Schou, H. Skriver, A. Nielsen, and K. Conradsen, "CFAR edge detector for polarimetric SAR images," *IEEE Transactions on Geoscience and Remote Sensing*, vol. 41, no. 1, pp. 20 – 32, Jan 2003.
- [20] T. H. Le, I. V. McLoughlin, K. Y. Lee, and T. Brestchneider, "SLC SAR speckle filtering using homoskedastic features of logarithmic transformation," in *Proceedings of the 31st Asian Conference on Remote Sensing (ACRS)*, Hanoi, Vietnam, Nov 2010.
- [21] T. H. Le and I. V. McLoughlin, "SAR Fuzzy-MLE speckle filter using the distance consistency property in homoskedastic log-transformed domain," in *Proceedings of the 32nd Asian Conference on Remote Sensing (ACRS)*, Taipei, Taiwan, Nov 2011.