

---

# Supplement of “Causal Shapley Values: Exploiting Causal Knowledge to Explain Individual Predictions of Complex Models”

---

Anonymous Author(s)

Affiliation

Address

email

## 1 Do-calculus for cyclic graphs

For completeness, we here repeat the rules of *do*-calculus for cyclic graphs, in the notation of the generalized ID algorithm of [2], which generalizes [5]. We are given a causal graph  $G$ . To each node  $X_i$  which is intervened upon, we add an ‘intervention node’  $I_{X_i}$ , with a directed edge from  $I_{X_i}$  to  $X_i$  that we clamp to the value  $x_i$ . The corresponding graph is called  $\hat{G}^+$ .  $\hat{G}_{do(\mathbf{W})}$  is now obtained by removing from  $\hat{G}^+$  all incoming edges to variables that are part of  $\mathbf{W}$ , except those from the corresponding intervention nodes  $I_{\mathbf{W}}$ . We use shorthand

$$\mathbf{Y} \overset{\sigma}{\perp\!\!\!\perp}_G \mathbf{X} \mid \mathbf{Z}, do(\mathbf{W})$$

to indicate that  $\mathbf{Y}$  and  $\mathbf{X}$  are  $\sigma$ -separated by  $\mathbf{Z}$  in the graph  $\hat{G}_{do(\mathbf{W})}$ .  $\sigma$ -separation is a generalization of standard d-separation (see [2] for details).

*Do*-calculus now consists of the following three inference rules that can be used to map interventional and observational distributions.

1. Insertion/deletion of observation:

$$P(\mathbf{Y} \mid \mathbf{X}, \mathbf{Z}, do(\mathbf{W})) = P(\mathbf{Y} \mid \mathbf{Z}, do(\mathbf{W})) \text{ if } \mathbf{Y} \overset{\sigma}{\perp\!\!\!\perp}_G \mathbf{X} \mid \mathbf{Z}, do(\mathbf{W}).$$

2. Action/observation exchange:

$$P(\mathbf{Y} \mid do(\mathbf{X}), \mathbf{Z}, do(\mathbf{W})) = P(\mathbf{Y} \mid \mathbf{X}, \mathbf{Z}, do(\mathbf{W})) \text{ if } \mathbf{Y} \overset{\sigma}{\perp\!\!\!\perp}_G I_{\mathbf{X}} \mid \mathbf{X}, \mathbf{Z}, do(\mathbf{W}).$$

3. Insertion/deletion of actions:

$$P(\mathbf{Y} \mid do(\mathbf{X}), \mathbf{Z}, do(\mathbf{W})) = P(\mathbf{Y} \mid \mathbf{Z}, do(\mathbf{W})) \text{ if } \mathbf{Y} \overset{\sigma}{\perp\!\!\!\perp}_G I_{\mathbf{X}} \mid \mathbf{Z}, do(\mathbf{W}).$$

Through consecutive application of these rules, we can try to turn any interventional probability of interest into an observational probability.

## 2 Shapley values for linear models

We will make use of the *do*-calculus rules above to derive the causal Shapley values for the four different models in Figure 1 in the main text. To this end, we consider the three models in Figure 1 that predict  $f(x_1, x_2) = \beta_1 x_1 + \beta_2 x_2$  for general values of  $\beta_1$  and  $\beta_2$ . All three models have the same observational probability distribution, with  $\mathbb{E}[X_i] = \bar{x}_i$  and  $\mathbb{E}[X_{3-i}] = \alpha_i x_i$ , for  $i = 1, 2$ , yet different




Figure 1: Three causal models with the same observational distribution over features, yet a different causal structure. To connect to the models in the main text, we set  $\beta_1 = 0$  and  $\beta_2 = \beta$ , except that for the ‘fork’ we set  $\beta_2 = 0$ ,  $\beta_1 = \beta$ , and then swap the indices.

causal structures. We will arrive at the main text’s results for the ‘chain’, ‘confounder’, and ‘cycle’ by setting  $\beta_1 = 0$ , whereas for the ‘fork’ we set  $\beta_2 = 0$  and swap the two indices. We then further need to take  $\bar{x}_1 = \bar{x}_2 = 0$ , and  $\alpha = \alpha_2$ .

Following the definitions in the main text, the contribution of feature  $i$  given permutation  $\pi$  is the difference in value function before and after setting the feature to its value:

$$\phi_i(\pi) = v(\{j : j \leq_\pi i\}) - v(\{j : j <_\pi i\}), \quad (1)$$

with value function

$$v(S) = \mathbb{E}[f(\mathbf{X}) | do(\mathbf{X}_S = \mathbf{x}_S)] = \int d\mathbf{X}_{\bar{S}} P(\mathbf{X}_{\bar{S}} | \hat{\mathbf{x}}_S) f(\mathbf{X}_{\bar{S}}, \mathbf{x}_S), \quad (2)$$

where we use shorthand  $\hat{\mathbf{x}}$  for  $do(\mathbf{X} = \mathbf{x})$ . Combining these two definitions and substituting  $f(\mathbf{x}) = \sum_i \beta_i x_i$ , we obtain

$$\phi_i(\pi) = \beta_i (x_i - \mathbb{E}[X_i | \hat{\mathbf{x}}_{j:j <_\pi i}]) + \sum_{k >_\pi i} \beta_k (\mathbb{E}[X_k | \hat{\mathbf{x}}_{j:j \leq_\pi i}] - \mathbb{E}[X_k | \hat{\mathbf{x}}_{j:j <_\pi i}]).$$

The first term corresponds to the direct effect, the second one to the indirect effect. Symmetric causal Shapley values will follow by averaging these contributions for the two possible permutations  $\pi = (1, 2)$  and  $\pi = (2, 1)$ . Conditional Shapley values result when replacing conditioning by intervention with conventional conditioning by observation, marginal Shapley values by not conditioning at all.

To start with the latter, we immediately see that for *marginal Shapley values* the indirect effect vanishes and the direct effect simplifies to

$$\phi_i = \phi_i(\pi) = \beta_i (x_i - \mathbb{E}[X_i]) = \beta_i (x_i - \bar{x}_i),$$

as also derived in [1].

For symmetric conditional Shapley values, we do get different contributions for the two different permutations, but by definition still the same for the three different models:

$$\begin{aligned} \phi_1(1, 2) &= \beta_1 (x_1 - \mathbb{E}[X_1]) + \beta_2 (\mathbb{E}[X_2 | x_1] - \mathbb{E}[X_2]) = \beta_1 (x_1 - \bar{x}_1) + \beta_2 \alpha_1 (x_1 - \bar{x}_1) \\ \phi_2(1, 2) &= \beta_2 (x_2 - \mathbb{E}[X_2 | x_1]) = \beta_2 (x_2 - \bar{x}_2) - \beta_2 \alpha_1 (x_1 - \bar{x}_1). \end{aligned} \quad (3)$$

Here the first term in the contribution for the first feature corresponds to the direct effect and the second term to the indirect effect. The contribution for the second feature only consists of a direct effect. The contributions for the other permutation follow by swapping the indices and the final Shapley values by averaging to arrive at the *symmetric conditional Shapley values*

$$\begin{aligned} \phi_1 &= \beta_1 (x_1 - \bar{x}_1) - \frac{1}{2} \beta_1 \alpha_2 (x_2 - \bar{x}_2) + \frac{1}{2} \beta_2 \alpha_1 (x_1 - \bar{x}_1) \\ \phi_2 &= \beta_2 (x_2 - \bar{x}_2) - \frac{1}{2} \beta_2 \alpha_1 (x_1 - \bar{x}_1) + \frac{1}{2} \beta_1 \alpha_2 (x_2 - \bar{x}_2), \end{aligned} \quad (4)$$

where now the first two terms constitute the direct effect and the third term the indirect effect.

expectation	chain	confounder	cycle
$\mathbb{E}[X_1 \hat{x}_2]$	$\mathbb{E}[X_1]$	$\mathbb{E}[X_1]$	$\mathbb{E}[X_1 x_2]$
$\mathbb{E}[X_2 \hat{x}_1]$	$\mathbb{E}[X_2 x_1]$	$\mathbb{E}[X_2]$	$\mathbb{E}[X_2 x_1]$

Table 1: Turning expectations under conditioning by intervention into expectations under conventional conditioning by observation for the three models in Figure 1.

44 The *asymmetric conditional Shapley values* consider both permutations for the confounder and the  
 45 cycle, and hence are equivalent to the symmetric Shapley values for those models. Yet for the chain,  
 46 they only consider the permutation  $\pi(1, 2)$  and thus yield  $\phi = \phi(1, 2)$  from (5).

47 To go from the symmetric conditional Shapley values to the causal symmetric Shapley values, we  
 48 follow the same line of reasoning, but have to replace  $\mathbb{E}[X_2|x_1]$  by  $\mathbb{E}[X_2|\hat{x}_1]$  and  $\mathbb{E}[X_1|x_2]$  by  $\mathbb{E}[X_1|\hat{x}_2]$ .  
 49 Table 1 tells whether the expectations under conditioning by intervention reduce to expectations  
 50 under conditioning by observation (because of the second rule of *do*-calculus above) or to marginal  
 51 expectations (because of the third rule). For the chain we have

$$P(X_2|\hat{x}_1) = P(X_2|x_1) \text{ since } X_2 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_1} | X_1 \text{ (rule 2), yet } P(X_1|\hat{x}_2) = P(X_1) \text{ since } X_1 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_2} \text{ (rule 3),}$$

52 for the confounder

$$P(X_2|\hat{x}_1) = P(X_2) \text{ since } X_2 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_1} \text{ and } P(X_1|\hat{x}_2) = P(X_1) \text{ since } X_1 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_2} \text{ (rule 3),}$$

53 and for the cycle

$$P(X_2|\hat{x}_1) = P(X_2|x_1) \text{ since } X_2 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_1} | X_1 \text{ and } P(X_1|\hat{x}_2) = P(X_1|x_2) \text{ since } X_1 \stackrel{\sigma}{\perp\!\!\!\perp}_G I_{X_2} | X_2 \text{ (rule 2).}$$

54 Consequently, for the confounder the *symmetric* and *asymmetric causal Shapley values* coincide with  
 55 the marginal Shapley values (consistent with [4]) and for the cycle with the symmetric conditional  
 56 Shapley values from (4). For the chain, the causal symmetric Shapley values become

$$\begin{aligned} \phi_1(1, 2) &= \beta_1(x_1 - \bar{x}_1) + \frac{1}{2}\beta_2\alpha_1(x_1 - \bar{x}_1) \\ \phi_2(1, 2) &= \beta_2(x_2 - \bar{x}_2) - \frac{1}{2}\beta_2\alpha_1(x_1 - \bar{x}_1), \end{aligned} \tag{5}$$

57 where the asymmetric causal Shapley values coincides with the asymmetric conditional Shapley  
 58 values from (5).

59 Collecting all results and setting  $\bar{x}_1 = \bar{x}_2 = \beta_1 = 0, \beta_2 = \beta$ , and  $\alpha_1 = \alpha$  (after swapping the indices for  
 60 the ‘fork’), we arrive at the Shapley values reported in Figure 1 in the main text. Note that for most  
 61 Shapley values, the indirect effect for the second feature vanishes because we chose to set  $\beta_1 = 0$ .  
 62 The exceptions, apart from the marginal Shapley values, are the causal Shapley values for the chain  
 63 and the confounder, as well as the asymmetric conditional Shapley values for the chain: these show  
 64 no indirect effect for the second feature even for nonzero  $\beta_1$ .

### 65 3 Proofs and corollaries on causal chain graphs

66 In this section we expand on the proof of Theorem 1 in the main text and add some corollaries to link  
 67 back to other approaches for computing Shapley values.

68 The probability distribution for a causal chain graph boils down to a directed acyclic graph of chain  
 69 components:

$$P(\mathbf{X}) = \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_\tau | \mathbf{X}_{pa(\tau)}). \tag{6}$$

70 For each (fully connected) chain component, we further need to specify whether (surplus) depen-  
 71 dencies within the component are due to confounding or due to mutual interactions. Given this  
 72 information, we can turn any causal query into an observational distribution with the following  
 73 interventional formula.

74 **Theorem 1.** *For causal chain graphs, we have the interventional formula*

$$P(\mathbf{X}_S | do(\mathbf{X}_S = \mathbf{x}_S)) = \prod_{\tau \in \mathcal{T}_{\text{confounding}}} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}) \times \prod_{\tau \in \overline{\mathcal{T}_{\text{confounding}}}} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{\tau \cap \bar{S}}). \quad (7)$$

75 *Proof.* Plugging in (6) and using shorthand  $\hat{\mathbf{x}} = do(\mathbf{X} = \mathbf{x})$ , we obtain

$$P(\mathbf{X}_{\bar{S}} | \hat{\mathbf{x}}_S) = P(\mathbf{X} | \hat{\mathbf{x}}_S) = \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_{\tau} | \mathbf{X}_{\tau' <_G \tau}, \hat{\mathbf{x}}_S) \stackrel{(1)}{=} \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_{\tau} | \mathbf{X}_{pa(\tau)}, \hat{\mathbf{x}}_S) = \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_S),$$

76 where in the second step we made use of *do*-calculus rule (1): the conditional independencies in the  
77 causal chain graph  $G$  are preserved when we intervene on some of the variables. Now rule (3) tells us  
78 that we can ignore any interventions from nodes in components further down the causal chain graph  
79 as well as those from higher up that are shielded by the direct parents:

$$P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_S) \stackrel{(3)}{=} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{\tau \cap \bar{S}}).$$

80 Rule (2) then states that conditioning by intervention upon variables higher up in the causal chain  
81 graph is equivalent to conditioning by observation:

$$P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{\tau \cap \bar{S}}) \stackrel{(2)}{=} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{\tau \cap \bar{S}}).$$

82 For a chain component with dependencies induced by a common confounder, rule (3) applies once  
83 more and makes that we can ignore the interventions:

$$P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{\tau \cap \bar{S}}) = P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}).$$

84 For a chain component with dependencies induced by mutual interactions, rule (2) again applies and  
85 allows us to replace conditioning by intervention with conditioning by observation:

$$P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}, \hat{\mathbf{x}}_{\tau \cap \bar{S}}) = P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{\tau \cap \bar{S}}).$$

86 □

87 Algorithm 1 provides pseudocode on how to estimate the value function  $v(S)$  by drawing samples  
88 from the interventional probability (7). It assumes that a user has specified a partial causal ordering  
89 of the features, which is translated to a chain graph with components  $\mathcal{T}$ , and for each (non-singleton)  
90 component  $\tau$  whether or not surplus dependencies are the result of confounding. Other prerequisites  
91 include access to the model  $f()$ , the feature vector  $\mathbf{x}$ , (a procedure to sample from) the observational  
92 probability distribution  $P(\mathbf{X})$ , and the number of samples  $N_{\text{samples}}$ .

93 Theorem 1 connects to observations made and algorithms proposed in recent papers.

94 **Corollary 1.** *With all features combined in a single component and all dependencies induced by*  
95 *confounding, as in [4], causal Shapley values are equivalent to marginal Shapley values.*

96 *Proof.* With just a single confounded component  $\tau$ ,  $pa(\tau) = \emptyset$  and (7) reduces to  $P(\mathbf{X}_{\bar{S}})$ . □

97 **Corollary 2.** *With all features combined in a single component and all dependencies induced by*  
98 *mutual interactions, causal Shapley values are equivalent to conditional Shapley values as proposed*  
99 *in [1].*

100 *Proof.* With just a single non-confounded component  $\tau$ ,  $pa(\tau) = \emptyset$  and (7) reduces to  $P(\mathbf{X}_{\bar{S}} | \mathbf{x}_S)$ . □

101 **Corollary 3.** *When we only take into account permutations that match the causal ordering and*  
102 *assume that all dependencies within chain components are induced by mutual interactions, the*  
103 *resulting asymmetric causal Shapley values are equivalent to the asymmetric conditional Shapley*  
104 *values as defined in [3].*

---

**Algorithm 1** Compute the value function  $v(S)$  under conditioning by intervention.

---

```

1: function VALUEFUNCTION( $S$ )
2:   for  $k \leftarrow 1$  to  $N_{\text{samples}}$  do
3:     for all  $j \leftarrow 1$  to  $|\mathcal{T}|$  do ▷ run over all chain components in their causal order
4:       if confounding( $\tau_j$ ) then
5:         for all  $i \in \tau_j \cap \bar{S}$  do
6:           Sample  $\tilde{x}_i^{(k)} \sim P(X_i | \tilde{\mathbf{x}}_{pa(\tau_j) \cap \bar{S}}, \mathbf{x}_{pa(\tau_j) \cap S})$  ▷ can be drawn independently
7:         end for
8:       else
9:         Sample  $\tilde{\mathbf{x}}_{\tau_j \cap \bar{S}}^{(k)} \sim P(\mathbf{X}_{\tau_j \cap \bar{S}} | \tilde{\mathbf{x}}_{pa(\tau_j) \cap \bar{S}}, \mathbf{x}_{pa(\tau_j) \cap S}, \mathbf{x}_{\tau_j \cap S})$  ▷ e.g., Gibbs sampling
10:      end if
11:    end for
12:  end for
13:   $v \leftarrow \frac{1}{N_{\text{samples}}} \sum_{k=1}^{N_{\text{samples}}} f(\mathbf{x}_S, \tilde{\mathbf{x}}_{\bar{S}}^{(k)})$ 
14:  return  $v$ 
15: end function

```

---

105 *Proof.* Following [3], asymmetric Shapley values only include those permutations  $\pi$  for which  $i <_{\pi} j$   
 106 for all known ancestors  $i$  of descendants  $j$  in the causal graph. For those permutations, we are  
 107 guaranteed to have  $\tau <_G \tau'$  for all  $\tau, \tau' \in \mathcal{T}$  such that  $\tau \cap S \neq \emptyset, \tau' \cap \bar{S} \neq \emptyset$ . That is, the chain  
 108 components that contain features from  $S$  are never later in the causal ordering of the chain graph  $G$   
 109 than those that contain features from  $\bar{S}$ . We then have

$$P(\mathbf{X}_{\bar{S}} | \mathbf{x}_S) = \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_S) = \prod_{\tau \in \mathcal{T}} P(\mathbf{X}_{\tau \cap \bar{S}} | \mathbf{X}_{pa(\tau) \cap \bar{S}}, \mathbf{x}_{pa(\tau) \cap S}, \mathbf{x}_{\tau \cap S}) = P(\mathbf{X}_{\bar{S}} | \hat{\mathbf{x}}_S),$$

110 where in the last step we used interventional formula (7) in combination with the fact that  $\mathcal{T}_{\text{confounding}} =$   
 111  $\emptyset$ . □

## 112 4 Additional illustrations on the bike rental data

113 Figure 2 shows sina plots for asymmetric conditional Shapley values (left) and marginal Shapley  
 114 values (right), to be compared with the sina plots for symmetric causal Shapley values in Figure 3  
 115 of the main text. In this case, the sina plots for asymmetric causal Shapley values are virtually  
 116 indistinguishable from those for the asymmetric conditional Shapley values.

117 It can be seen that the marginal Shapley values strongly focus on temperature, largely ignoring the  
 118 seasonal variables. The asymmetric Shapley values do the opposite: they focus on the seasonal  
 119 variables, in particular *cosyear* and put much less emphasis on the temperature variables.

## 120 References

- 121 [1] Kjersti Aas, Martin Jullum, and Anders Løland. Explaining individual predictions when  
 122 features are dependent: More accurate approximations to Shapley values. *arXiv preprint*  
 123 *arXiv:1903.10464*, 2019.
- 124 [2] Patrick Forré and Joris M Mooij. Causal calculus in the presence of cycles, latent confounders  
 125 and selection bias. In *Proceedings of the 35th Annual Conference on Uncertainty in Artificial*  
 126 *Intelligence*, 2019.
- 127 [3] Christopher Frye, Ilya Feige, and Colin Rowat. Asymmetric Shapley values: incorporating causal  
 128 knowledge into model-agnostic explainability. *arXiv preprint arXiv:1910.06358*, 2019.
- 129 [4] Dominik Janzing, Lenon Minorics, and Patrick Blöbaum. Feature relevance quantification in  
 130 explainable AI: A causality problem. *arXiv preprint arXiv:1910.13413*, 2019.
- 131 [5] Judea Pearl. The *do*-calculus revisited. *arXiv preprint arXiv:1210.4852*, 2012.




Figure 2: Sina plots of asymmetric (conditional) Shapley values (left) and marginal Shapley values (right). See Figure 3 in the main text for further details.