

Computing Abelian varieties over finite fields

Marseglia Stefano

Utrecht University

04 June 2019
Lightning Talk
ICERM

Motivation

up to isogeny: Honda-Tate theory:

Motivation

up to isogeny: Honda-Tate theory:

- the association

$A/\mathbb{F}_q \mapsto \text{char.poly.}(\text{Frob}_A : T_l A \rightarrow T_l A)$ which is a q -Weil poly.

allows us to **enumerate** all abelian varieties up to isogeny.

up to isomorphism (setup):

Motivation

up to isogeny: Honda-Tate theory:

- the association

$A/\mathbb{F}_q \mapsto \text{char.poly.}(\text{Frob}_A : T_l A \rightarrow T_l A)$ which is a q -Weil poly.

allows us to **enumerate** all abelian varieties up to isogeny.

up to isomorphism (setup):

- Let h be a squarefree ordinary q -Weil polynomial or a squarefree p -Weil polynomial without real roots.

Motivation

up to isogeny: Honda-Tate theory:

- the association

$A/\mathbb{F}_q \mapsto \text{char.poly.}(\text{Frob}_A : T_1 A \rightarrow T_1 A)$ which is a q -Weil poly.

allows us to **enumerate** all abelian varieties up to isogeny.

up to isomorphism (setup):

- Let h be a squarefree ordinary q -Weil polynomial or a squarefree p -Weil polynomial without real roots.
- Put $K = \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$.

Motivation

up to isogeny: Honda-Tate theory:

- the association

$A/\mathbb{F}_q \mapsto \text{char.poly.}(\text{Frob}_A : T_1 A \rightarrow T_1 A)$ which is a q -Weil poly.

allows us to **enumerate** all abelian varieties up to isogeny.

up to isomorphism (setup):

- Let h be a squarefree ordinary q -Weil polynomial or a squarefree p -Weil polynomial without real roots.
- Put $K = \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$.
- Denote by \mathcal{C}_h the isogeny class of abelian varieties with char. poly. of Frobenius h .

Main result

Theorem (M.)

$$\begin{array}{c} \{ \text{abelian varieties in } \mathcal{C}_h \} / \simeq \\ \updownarrow \\ \{ \text{fractional ideals of } \mathbb{Z}[F, q/F] \subset K \} / \simeq \end{array}$$

Main result

Theorem (M.)

$$\begin{array}{c} \{ \text{abelian varieties in } \mathcal{C}_h \} / \simeq \\ \updownarrow \\ \{ \text{fractional ideals of } \mathbb{Z}[F, q/F] \subset K \} / \simeq \end{array} =: \text{ICM}(\mathbb{Z}[F, q/F])$$

ideal class monoid

Main result

Theorem (M.)

$$\begin{array}{c} \{abelian\ varieties\ in\ \mathcal{C}_h\} / \simeq \\ \updownarrow \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, q/F] \subset K\} / \simeq \end{array} =: \text{ICM}(\mathbb{Z}[F, q/F])$$

ideal class monoid

- Problem: $\mathbb{Z}[F, q/F]$ might not be maximal \Rightarrow **non-invertible** ideals.

Main result

Theorem (M.)

$$\begin{array}{c} \{abelian\ varieties\ in\ \mathcal{C}_h\} / \simeq \\ \updownarrow \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, q/F] \subset K\} / \simeq \end{array} =: \text{ICM}(\mathbb{Z}[F, q/F])$$

ideal class monoid

- Problem: $\mathbb{Z}[F, q/F]$ might not be maximal \Rightarrow **non-invertible** ideals.
- Solution: attack the problem locally (weak equivalence) then use the action of the Pic's.

Main result

Theorem (M.)

$$\begin{array}{c} \{abelian\ varieties\ in\ \mathcal{C}_h\} / \simeq \\ \updownarrow \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, q/F] \subset K\} / \simeq \end{array} =: \text{ICM}(\mathbb{Z}[F, q/F])$$

ideal class monoid

- Problem: $\mathbb{Z}[F, q/F]$ might not be maximal \Rightarrow **non-invertible** ideals.
- Solution: attack the problem locally (weak equivalence) then use the action of the Pic's.
- Extra 1: in the ordinary case we can compute also **polarizations**.

Main result

Theorem (M.)

$$\begin{array}{c} \{ \text{abelian varieties in } \mathcal{C}_h \} / \simeq \\ \updownarrow \\ \{ \text{fractional ideals of } \mathbb{Z}[F, q/F] \subset K \} / \simeq \end{array} =: \text{ICM}(\mathbb{Z}[F, q/F])$$

ideal class monoid

- Problem: $\mathbb{Z}[F, q/F]$ might not be maximal \Rightarrow **non-invertible** ideals.
- Solution: attack the problem locally (weak equivalence) then use the action of the Pic's.
- Extra 1: in the ordinary case we can compute also **polarizations**.
- Extra 2: similar results for isogeny classes \mathcal{C}_{h^r} (when $\mathbb{Z}[F, q/F]$ is Bass)

Coming soon on the LMFDB.

Thank you!