

Polarizations of abelian varieties over finite fields via canonical liftings

Stefano Marseglia

Utrecht University

UGC Seminar - 29 March 2022

Polarizations of abelian varieties over finite fields via canonical liftings

Stefano Marseglia

Utrecht University

UGC Seminar - 29 March 2022

joint work with

Jonas Bergström and **Valentijn Karemaker**.

Abelian Varieties

- An **abelian variety** A over a field k is a projective geometrically connected group variety over k .

Abelian Varieties

- An **abelian variety** A over a field k is a projective geometrically connected group variety over k .
We have **morphisms** $\oplus : A \times A \rightarrow A$, $\ominus : A \rightarrow A$ and a k -rational point $e \in A(k)$ such that (A, \oplus, \ominus, e) is a group object in the category of projective geom. connected varieties over k .

Abelian Varieties

- An **abelian variety** A over a field k is a projective geometrically connected group variety over k .
We have **morphisms** $\oplus : A \times A \rightarrow A$, $\ominus : A \rightarrow A$ and a k -rational point $e \in A(k)$ such that (A, \oplus, \ominus, e) is a group object in the category of projective geom. connected varieties over k .
- In practice, we have diagrams \rightsquigarrow “**natural**” group structure on $A(\bar{k})$.

Abelian Varieties

- An **abelian variety** A over a field k is a projective geometrically connected group variety over k .
We have **morphisms** $\oplus : A \times A \rightarrow A$, $\ominus : A \rightarrow A$ and a k -rational point $e \in A(k)$ such that (A, \oplus, \ominus, e) is a group object in the category of projective geom. connected varieties over k .
- In practice, we have diagrams \rightsquigarrow “**natural**” group structure on $A(\bar{k})$.
- eg. (\ominus is the “inverse” morphism)

$$\begin{array}{ccc} A \times_k A & \xrightarrow{(\ominus, \text{id})} & A \times_k A \\ \uparrow \Delta & & \downarrow \oplus \\ A & \xrightarrow{\quad} \text{Spec}(k) \xrightarrow{e} & A \end{array}$$

Abelian Varieties

- An **abelian variety** A over a field k is a projective geometrically connected group variety over k .
We have **morphisms** $\oplus : A \times A \rightarrow A$, $\ominus : A \rightarrow A$ and a k -rational point $e \in A(k)$ such that (A, \oplus, \ominus, e) is a group object in the category of projective geom. connected varieties over k .
- In practice, we have diagrams \rightsquigarrow “**natural**” group structure on $A(\bar{k})$.
- eg. (\ominus is the “inverse” morphism)

$$\begin{array}{ccc}
 A \times_k A & \xrightarrow{(\ominus, \text{id})} & A \times_k A \\
 \uparrow \Delta & & \downarrow \oplus \\
 A & \xrightarrow{\quad} \text{Spec}(k) \xrightarrow{e} & A
 \end{array}$$

$$\begin{array}{ccc}
 A \times_k A & \xrightarrow{(\text{id}, \ominus)} & A \times_k A \\
 \uparrow \Delta & & \downarrow \oplus \\
 A & \xrightarrow{\quad} \text{Spec}(k) \xrightarrow{e} & A
 \end{array}$$

Example : $\dim A = 1$ elliptic curves

- AVs of dimension 1 are called **elliptic curves**.

Example : $\dim A = 1$ elliptic curves

- AVs of dimension 1 are called **elliptic curves**.
- They admit a plane model: if $\text{char } k \neq 2, 3$

$$Y^2Z = X^3 + AXZ^2 + BZ^3 \quad A, B \in k \text{ and } e = [0 : 1 : 0]$$

Example : $\dim A = 1$ elliptic curves

- AVs of dimension 1 are called **elliptic curves**.
- They admit a plane model: if $\text{char } k \neq 2, 3$

$$Y^2Z = X^3 + AXZ^2 + BZ^3 \quad A, B \in k \text{ and } e = [0 : 1 : 0]$$

- The groups law is explicit:
if $P = (x_P, y_P)$ then $\ominus P = (x_P, -y_P)$ and

Example : $\dim A = 1$ elliptic curves

- AVs of dimension 1 are called **elliptic curves**.
- They admit a plane model: if $\text{char } k \neq 2, 3$

$$Y^2Z = X^3 + AXZ^2 + BZ^3 \quad A, B \in k \text{ and } e = [0 : 1 : 0]$$

- The groups law is explicit:
if $P = (x_P, y_P)$ then $\ominus P = (x_P, -y_P)$ and
if $Q = (x_Q, y_Q) \neq \ominus P$ then $P \oplus Q = (x_R, y_R)$ where

$$x_R = \lambda^2 - x_P - x_Q, \quad y_R = y_P + \lambda(x_R - x_P),$$

where


$$\lambda = \begin{cases} \frac{3x_P^2 + B}{2A} & \text{if } P = Q \\ \frac{y_P - y_Q}{x_P - x_Q} & \text{if } P \neq Q \end{cases}$$

Example : EC over \mathbb{R}

Example : EC over \mathbb{R}

Over \mathbb{R} :
consider the abelian variety:

$$y^2 = x^3 - x + 1$$




Example : EC over \mathbb{R}

Over \mathbb{R} :
consider the abelian variety:

$$y^2 = x^3 - x + 1$$

Addition law: $P, Q \rightsquigarrow P \oplus Q$



Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .
- An isogeny $\mu : A \rightarrow A^\vee$ (over k) is called a **polarization** if there are an $k \subseteq k'$ and an ample line bundle \mathcal{L} such that (on points)

$$\varphi_{k'} : x \mapsto [t_x^* \mathcal{L} \otimes \mathcal{L}^{-1}].$$

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .
- An isogeny $\mu : A \rightarrow A^\vee$ (over k) is called a **polarization** if there are an $k \subseteq k'$ and an ample line bundle \mathcal{L} such that (on points)

$$\varphi_{k'} : x \mapsto [t_x^* \mathcal{L} \otimes \mathcal{L}^{-1}].$$

- A polarization μ is **principal** if $\deg \mu = 1 \iff \mu$ is an isomorphism.

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .
- An isogeny $\mu : A \rightarrow A^\vee$ (over k) is called a **polarization** if there are an $k \subseteq k'$ and an ample line bundle \mathcal{L} such that (on points)

$$\varphi_{k'} : x \mapsto [t_x^* \mathcal{L} \otimes \mathcal{L}^{-1}].$$

- A polarization μ is **principal** if $\deg \mu = 1 \iff \mu$ is an isomorphism.
- **Why** do we care about polarizations?

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .
- An isogeny $\mu : A \rightarrow A^\vee$ (over k) is called a **polarization** if there are an $k \subseteq k'$ and an ample line bundle \mathcal{L} such that (on points)

$$\varphi_{k'} : x \mapsto [t_x^* \mathcal{L} \otimes \mathcal{L}^{-1}].$$

- A polarization μ is **principal** if $\deg \mu = 1 \iff \mu$ is an isomorphism.
- **Why** do we care about polarizations?
 - ① $\text{Aut}(A, \mu)$ is finite \rightsquigarrow moduli space $\mathcal{A}_{g,d}$

Duals and Polarizations

- A hom. $\varphi : A \rightarrow B$ is an **isogeny** if $\dim A = \dim B$ and φ is surjective.
- Isogenies have finite kernel: $\deg \varphi = \text{rank}(\ker(\varphi))$
- Pic_A^0 is also an AV, called the **dual** of A and denoted A^\vee .
- An isogeny $\mu : A \rightarrow A^\vee$ (over k) is called a **polarization** if there are an $k \subseteq k'$ and an ample line bundle \mathcal{L} such that (on points)

$$\varphi_{k'} : x \mapsto [t_x^* \mathcal{L} \otimes \mathcal{L}^{-1}].$$

- A polarization μ is **principal** if $\deg \mu = 1 \iff \mu$ is an isomorphism.
- **Why** do we care about polarizations?
 - 1 $\text{Aut}(A, \mu)$ is finite \rightsquigarrow moduli space $\mathcal{A}_{g,d}$
 - 2 proper smooth curve $C/k \rightsquigarrow \text{Pic}_C^0 =: \text{Jac}(C)$ a PPAV.

- Pick A/\mathbb{C} of dimension g .

- Pick A/\mathbb{C} of dimension g .
- $A(\mathbb{C}) \simeq V := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$. It is a torus.

- Pick A/\mathbb{C} of dimension g .
- $A(\mathbb{C}) \simeq V := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$. It is a torus.
- V admits a non-degenerate **Riemann form** \longleftrightarrow polarization.

- Pick A/\mathbb{C} of dimension g .
- $A(\mathbb{C}) \simeq V := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$. It is a torus.
- V admits a non-degenerate **Riemann form** \longleftrightarrow polarization.
- Actually,

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting} \right. \\ \left. \text{a Riemann form} \right\}$$

induced by $A \mapsto A(\mathbb{C})$ is an **equivalence** of categories.

- Pick A/\mathbb{C} of dimension g .
- $A(\mathbb{C}) \simeq V := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$. It is a torus.
- V admits a non-degenerate **Riemann form** \longleftrightarrow polarization.
- Actually,

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting a Riemann form} \right\}$$

induced by $A \mapsto A(\mathbb{C})$ is an **equivalence** of categories.

- In char. $p > 0$ such an equivalence cannot exist : there are (supersingular) elliptic curves with quaternionic endomorphism algebras.

Canonical Liftings

- Let A_0 be an abelian variety over \mathbb{F}_q of dim g .

Canonical Liftings

- Let A_0 be an abelian variety over \mathbb{F}_q of dim g .

Definition

A **canonical lifting** of A_0 is an abelian scheme over a normal local domain \mathcal{R} of characteristic zero with residue field \mathbb{F}_q with:

- 1 special fiber A_0 , and
- 2 general fiber \mathcal{A}_{can} satisfying $\text{End}(\mathcal{A}_{\text{can}}) = \text{End}(A_0)$.

Canonical Liftings

- Let A_0 be an abelian variety over \mathbb{F}_q of dim g .

Definition

A **canonical lifting** of A_0 is an abelian scheme over a normal local domain \mathcal{R} of characteristic zero with residue field \mathbb{F}_q with:

- 1 special fiber A_0 , and
 - 2 general fiber \mathcal{A}_{can} satisfying $\text{End}(\mathcal{A}_{\text{can}}) = \text{End}(A_0)$.
- A_0 comes with a Frobenius endomorphism induced by $x \mapsto x^q$ on coordinates rings (we are in $\text{char}(\mathbb{F}_q) = p > 0!$)

Canonical Liftings

- Let A_0 be an abelian variety over \mathbb{F}_q of dim g .

Definition

A **canonical lifting** of A_0 is an abelian scheme over a normal local domain \mathcal{R} of characteristic zero with residue field \mathbb{F}_q with:

- 1 special fiber A_0 , and
 - 2 general fiber \mathcal{A}_{can} satisfying $\text{End}(\mathcal{A}_{\text{can}}) = \text{End}(A_0)$.
- A_0 comes with a Frobenius endomorphism induced by $x \mapsto x^q$ on coordinates rings (we are in $\text{char}(\mathbb{F}_q) = p > 0$!)
 - Example: ordinary abelian variety; almost-ordinary abelian variety.

Canonical Liftings

- Let A_0 be an abelian variety over \mathbb{F}_q of dim g .

Definition

A **canonical lifting** of A_0 is an abelian scheme over a normal local domain \mathcal{R} of characteristic zero with residue field \mathbb{F}_q with:

- 1 special fiber A_0 , and
 - 2 general fiber \mathcal{A}_{can} satisfying $\text{End}(\mathcal{A}_{\text{can}}) = \text{End}(A_0)$.
- A_0 comes with a Frobenius endomorphism induced by $x \mapsto x^q$ on coordinates rings (we are in $\text{char}(\mathbb{F}_q) = p > 0!$)
 - Example: ordinary abelian variety; almost-ordinary abelian variety.
 - Non-example: supersingular EC (quaternions).

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.
- A_{can} has morphisms F (and $V = \frac{q}{F}$) reducing to Frobenius (and Verschiebung).

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.
- A_{can} has morphisms F (and $V = \frac{q}{F}$) reducing to Frobenius (and Verschiebung).
- By **complex uniformization**:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I)$$

- I : a fractional $\mathbb{Z}[F, V]$ -ideal in $L := \mathbb{Q}[F]$,
- Φ : a **CM-type** of L (g maps $L \rightarrow \mathbb{C}$, one per conjugate pair).

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.
- A_{can} has morphisms F (and $V = \frac{q}{F}$) reducing to Frobenius (and Verschiebung).
- By **complex uniformization**:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I)$$

- I : a fractional $\mathbb{Z}[F, V]$ -ideal in $L := \mathbb{Q}[F]$,
- Φ : a **CM-type** of L (g maps $L \rightarrow \mathbb{C}$, one per conjugate pair).

- Define $\mathcal{H}(A_{\text{can}}) := I$.

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.
- A_{can} has morphisms F (and $V = \frac{q}{F}$) reducing to Frobenius (and Verschiebung).
- By **complex uniformization**:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I)$$

- I : a fractional $\mathbb{Z}[F, V]$ -ideal in $L := \mathbb{Q}[F]$,
 - Φ : a **CM-type** of L (g maps $L \rightarrow \mathbb{C}$, one per conjugate pair).

- Define $\mathcal{H}(A_{\text{can}}) := I$.
- By the same construction:

$$\text{char.0:} \quad \mathbb{F}_q : A_0^V \xrightarrow{\quad \otimes \mathbb{C} \quad} \mathcal{A}_{\text{can}}^V \xrightarrow{\quad \mathcal{H} \quad} \bar{I}^t = \{\bar{x} : \text{Tr}_{L/\mathbb{Q}}(xI) \subseteq \mathbb{Z}\}$$

Complex Uniformization

- Assume that A_0 admits a canonical lifting \mathcal{A}_{can} .
- Fix $\mathcal{R} \hookrightarrow \mathbb{C}$ and put $A_{\text{can}} := \mathcal{A}_{\text{can}} \otimes \mathbb{C}$.
- A_{can} has morphisms F (and $V = \frac{q}{F}$) reducing to Frobenius (and Verschiebung).
- By **complex uniformization**:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I) \quad \begin{array}{l} - I : \text{a fractional } \mathbb{Z}[F, V]\text{-ideal in } L := \mathbb{Q}[F], \\ - \Phi : \text{a \textbf{CM-type} of } L \text{ (} g \text{ maps } L \rightarrow \mathbb{C}, \text{ one} \\ \text{per conjugate pair).} \end{array}$$

- Define $\mathcal{H}(A_{\text{can}}) := I$.
- By the same construction:

$$\text{char.0:} \quad \mathcal{A}_{\text{can}}^V \xrightarrow{\otimes \mathbb{C}} A_{\text{can}}^V \xrightarrow{\mathcal{H}} \bar{I}^t = \{\bar{x} : \text{Tr}_{L/\mathbb{Q}}(xI) \subseteq \mathbb{Z}\}$$

$$\mathbb{F}_q : A_0^V$$

- In particular: $\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^V)) = (\bar{I}^t : I) = \{x \in L : xI \subseteq \bar{I}^t\}$.

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

- What about **polarizations**? We understand them over \mathbb{C} !

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

- What about **polarizations**? We understand them over \mathbb{C} !
- Let $\mu : A_{\text{can}} \rightarrow A_{\text{can}}^{\vee}$ an isogeny. Then μ is a polarization if and only if $\lambda := \mathcal{H}(\mu) \in (\bar{I}^t : I)$ satisfies

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

- What about **polarizations**? We understand them over \mathbb{C} !
- Let $\mu : A_{\text{can}} \rightarrow A_{\text{can}}^{\vee}$ an isogeny. Then μ is a polarization if and only if $\lambda := \mathcal{H}(\mu) \in (\bar{I}^t : I)$ satisfies
 - 1 $\lambda = -\bar{\lambda}$ (totally imaginary), and

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

- What about **polarizations**? We understand them over \mathbb{C} !
- Let $\mu : A_{\text{can}} \rightarrow A_{\text{can}}^{\vee}$ an isogeny. Then μ is a polarization if and only if $\lambda := \mathcal{H}(\mu) \in (\bar{I}^t : I)$ satisfies
 - 1 $\lambda = -\bar{\lambda}$ (**totally imaginary**), and
 - 2 for every $\varphi \in \Phi$ we have $\text{Im}(\varphi(\lambda)) > 0$ (**Φ -positive**).

Complex Uniformization : Polarizations

- We have:

$$A_{\text{can}}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(I), \quad A_{\text{can}}^{\vee}(\mathbb{C}) \simeq \mathbb{C}^g / \Phi(\bar{I}^t),$$

$$\mathcal{H}(\text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee})) = (\bar{I}^t : I).$$

- What about **polarizations**? We understand them over \mathbb{C} !
- Let $\mu : A_{\text{can}} \rightarrow A_{\text{can}}^{\vee}$ an isogeny. Then μ is a polarization if and only if $\lambda := \mathcal{H}(\mu) \in (\bar{I}^t : I)$ satisfies
 - 1 $\lambda = -\bar{\lambda}$ (**totally imaginary**), and
 - 2 for every $\varphi \in \Phi$ we have $\text{Im}(\varphi(\lambda)) > 0$ (**Φ -positive**).

Isogeny classification over \mathbb{F}_q

- The Frobenius endomorphism A/\mathbb{F}_q comes

Isogeny classification over \mathbb{F}_q

- The Frobenius endomorphism A/\mathbb{F}_q comes induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

Isogeny classification over \mathbb{F}_q

- The Frobenius endomorphism A/\mathbb{F}_q comes induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a q -**Weil** polynomial and isogeny invariant.

Isogeny classification over \mathbb{F}_q

- The Frobenius endomorphism A/\mathbb{F}_q comes induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a q -**Weil** polynomial and isogeny invariant.
- By **Honda-Tate** theory, the association

$$\text{isogeny class of } A \longmapsto h_A(x)$$

is injective and allows us to **list** all isogeny classes.

Isogeny classification over \mathbb{F}_q

- The Frobenius endomorphism A/\mathbb{F}_q comes induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a q -**Weil** polynomial and isogeny invariant.
- By **Honda-Tate** theory, the association

$$\text{isogeny class of } A \longmapsto h_A(x)$$

is injective and allows us to **list** all isogeny classes.

- One can prove that $h_A(x)$ is squarefree $\iff \text{End}(A)$ is commutative.

Isomorphism classification over \mathbb{F}_p

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

- Let A_h be an AV in $AV_h(p)$ with $\text{End}(A_h) = \mathbb{Z}[F, V]$.

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

- Let A_h be an AV in $AV_h(p)$ with $\text{End}(A_h) = \mathbb{Z}[F, V]$.
- The functor $\mathcal{G}(-) := \text{Hom}(-, A_h)$ induces the equivalence.

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

- Let A_h be an AV in $AV_h(p)$ with $\text{End}(A_h) = \mathbb{Z}[F, V]$.
- The functor $\mathcal{G}(-) := \text{Hom}(-, A_h)$ induces the equivalence.
- We can **choose** A_h so that for every $B_0 \in AV_h(p)$:

$$\mathcal{G}(B_0^\vee) = \overline{\mathcal{G}(B_0)}^t$$

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

- Let A_h be an AV in $AV_h(p)$ with $\text{End}(A_h) = \mathbb{Z}[F, V]$.
- The functor $\mathcal{G}(-) := \text{Hom}(-, A_h)$ induces the equivalence.
- We can **choose** A_h so that for every $B_0 \in AV_h(p)$:

$$\mathcal{G}(B_0^\vee) = \overline{\mathcal{G}(B_0)}^t \text{ and } \mathcal{G}(f^\vee) = \overline{\mathcal{G}(f)}, \text{ for any } f : B_0 \rightarrow B'_0 \text{ in } AV_h(p).$$

Isomorphism classification over \mathbb{F}_p

Theorem (Centeleghe-Stix)

Let $AV_h(p)$ be the isogeny class over the **prime field** \mathbb{F}_p determined by a **squarefree** characteristic polynomial of Frobenius h .

Let $L = \mathbb{Q}[x]/h = \mathbb{Q}[F]$ be the endomorphism algebra, and put $V = p/F$.
There is an **equivalence** of categories:

$$AV_h(p) \xrightarrow{\mathcal{G}} \{\text{fractional } \mathbb{Z}[F, V]\text{-ideals in } L\}.$$

- Let A_h be an AV in $AV_h(p)$ with $\text{End}(A_h) = \mathbb{Z}[F, V]$.
- The functor $\mathcal{G}(-) := \text{Hom}(-, A_h)$ induces the equivalence.
- We can **choose** A_h so that for every $B_0 \in AV_h(p)$:

$$\mathcal{G}(B_0^\vee) = \overline{\mathcal{G}(B_0)}^t \text{ and } \mathcal{G}(f^\vee) = \overline{\mathcal{G}(f)}, \text{ for any } f: B_0 \rightarrow B_0' \text{ in } AV_h(p).$$

- In particular:

$$\mathcal{G}(\text{Hom}(B_0, B_0^\vee)) = (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)}^t).$$

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.

$$\begin{array}{ccc} \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\ \downarrow \text{red} & \searrow \text{complex unif.} & \\ \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \end{array}$$

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.

$$\begin{array}{ccc} \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\ \downarrow \text{red} \searrow \text{complex unif.} & & \\ \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\ \downarrow \mathcal{G} & & \downarrow \\ (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & \equiv & (\bar{I}^t : I) \end{array}$$

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.

$$\begin{array}{ccc}
 \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 \downarrow \text{red} & \searrow \text{complex unif.} & \\
 \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \cdot \alpha \begin{array}{l} \text{tot. real } (\alpha = \bar{\alpha}) \\ \text{unit in End}(A_0) \end{array} \\
 (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & \equiv & (\bar{I}^t : I)
 \end{array}$$

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.
- Let $f : A_0 \rightarrow B_0$ be an isogeny.

$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow \text{complex unif.} & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \begin{array}{l} \text{tot. real } (\alpha = \bar{\alpha}) \\ \text{unit in End}(A_0) \end{array} \\
 & & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & = & (\bar{I}^t : I)
 \end{array}$$

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.
- Let $f : A_0 \rightarrow B_0$ be an isogeny.

$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow \text{complex unif.} & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ - \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \begin{array}{l} \text{tot. real } (\alpha = \bar{\alpha}) \\ \text{unit in End}(A_0) \end{array} \\
 (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)}^t) & \xrightarrow{\mathcal{G}(f^*)} & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & = & (\bar{I}^t : I)
 \end{array}$$

- f^* sends polarizations to polarizations.

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.
- Let $f : A_0 \rightarrow B_0$ be an isogeny.

$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow \text{complex unif.} & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \text{ tot. real } (\alpha = \bar{\alpha}) \\
 (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)})^t & \xrightarrow{\mathcal{G}(f^*)} & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)})^t & = & \downarrow \text{unit in End}(A_0) \\
 & & & & (\bar{I}^t : I)
 \end{array}$$

- f^* sends polarizations to polarizations.
- $\mathcal{G}(f^*) = \mathcal{G}(f)\mathcal{G}(f)$ is a totally positive element:

Comparison

- Assume that A_0 admits a canonical lifting A_{can} .
- We have two description using fractional ideals. Let's compare them.
- Let $f : A_0 \rightarrow B_0$ be an isogeny.

$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow \text{complex unif.} & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \text{ tot. real } (\alpha = \bar{\alpha}) \\
 (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)})^t & \xrightarrow{\mathcal{G}(f^*)} & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)})^t & = & (\bar{I}^t : I) \\
 & & & & \downarrow \text{unit in End}(A_0)
 \end{array}$$

- f^* sends polarizations to polarizations.
- $\mathcal{G}(f^*) = \mathcal{G}(f)\mathcal{G}(f)$ is a totally positive element:
it sends totally imaginary elements to totally imaginary elements and Φ -positive elements to Φ -positive elements.

Comparison : Polarizations

$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ - \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \\
 (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)}^t) & \xrightarrow{\mathcal{G}(f^*)} & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & = & (\bar{I}^t : I)
 \end{array}$$


Comparison : Polarizations

we WANT to understand pols. here

we DO understand pols. here


$$\begin{array}{ccccc}
 & & \text{Hom}(A_{\text{can}}, A_{\text{can}}^{\vee}) & & \\
 & & \downarrow \text{red} & \searrow & \\
 \text{Hom}(B_0, B_0^{\vee}) & \xrightarrow{f^* := f^{\vee} \circ - \circ f} & \text{Hom}(A_0, A_0^{\vee}) & & (\bar{I}^t : I) \\
 \downarrow \mathcal{G} & & \downarrow \mathcal{G} & & \downarrow \cdot \alpha \\
 (\mathcal{G}(B_0) : \overline{\mathcal{G}(B_0)}^t) & \xrightarrow{\mathcal{G}(f^*)} & (\mathcal{G}(A_0) : \overline{\mathcal{G}(A_0)}^t) & = & (\bar{I}^t : I)
 \end{array}$$

Comparison : Polarizations



By chasing the diagram, we get:

Comparison : Polarizations



By chasing the diagram, we get:

Theorem ("lift and spread")

Let $\mu : B_0 \rightarrow B_0^{\vee}$ be an isogeny. Then

μ is a **polarization** $\iff \alpha^{-1} \mathcal{G}(\mu)$ is **totally imaginary** and Φ -positive

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.
- Assume that $B_0 \simeq B_0^\vee$, i.e. $J = i_0 \overline{J}^t$ for some $i_0 \in L^*$.

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.
- Assume that $B_0 \simeq B_0^\vee$, i.e. $J = i_0 \bar{J}^t$ for some $i_0 \in L^*$.
- If μ and μ' are principal polarizations of B_0 then $(B_0, \mu) \simeq (B_0, \mu')$ (as PPAVs) if and only if there is $v \in T^*$ such that $\mathcal{G}(\mu) = v \bar{v} \mathcal{G}(\mu')$.

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.
- Assume that $B_0 \simeq B_0^\vee$, i.e. $J = i_0 \bar{J}^t$ for some $i_0 \in L^*$.
- If μ and μ' are principal polarizations of B_0 then $(B_0, \mu) \simeq (B_0, \mu')$ (as PPAVs) if and only if there is $v \in T^*$ such that $\mathcal{G}(\mu) = v \bar{v} \mathcal{G}(\mu')$.
- Let \mathcal{T} be a transversal of $T^* / \langle v \bar{v} : v \in T^* \rangle$.

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.
- Assume that $B_0 \simeq B_0^\vee$, i.e. $J = i_0 \bar{J}^t$ for some $i_0 \in L^*$.
- If μ and μ' are principal polarizations of B_0 then $(B_0, \mu) \simeq (B_0, \mu')$ (as PPAVs) if and only if there is $v \in T^*$ such that $\mathcal{G}(\mu) = v \bar{v} \mathcal{G}(\mu')$.
- Let \mathcal{T} be a transversal of $T^* / \langle v \bar{v} : v \in T^* \rangle$.
- Then

$$\mathcal{P}_\Phi^\alpha(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

is a set of representatives of the PPs of B_0 up to isomorphism.

Principal Polarizations up to isomorphism

- Let $B_0 \in AV_h(p)$. Put $T = \text{End}(B_0)$ and $\mathcal{G}(B_0) = J$.
- Assume that $B_0 \simeq B_0^\vee$, i.e. $J = i_0 \bar{J}^t$ for some $i_0 \in L^*$.
- If μ and μ' are principal polarizations of B_0 then $(B_0, \mu) \simeq (B_0, \mu')$ (as PPAVs) if and only if there is $v \in T^*$ such that $\mathcal{G}(\mu) = v \bar{v} \mathcal{G}(\mu')$.
- Let \mathcal{T} be a transversal of $T^* / \langle v \bar{v} : v \in T^* \rangle$.
- Then

$$\mathcal{P}_\Phi^\alpha(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

is a set of representatives of the PPs of B_0 up to isomorphism.

- It depends on α !

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Theorem (1)

Denote by $S_{\mathbb{R}}^$ (resp. $T_{\mathbb{R}}^*$) the group of totally real units of S (resp. T).*

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Theorem (1)

Denote by $S_{\mathbb{R}}^*$ (resp. $T_{\mathbb{R}}^*$) the group of totally real units of S (resp. T).

If $S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$, then the set

$$\mathcal{P}_{\Phi}^{\alpha}(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Theorem (1)

Denote by $S_{\mathbb{R}}^*$ (resp. $T_{\mathbb{R}}^*$) the group of totally real units of S (resp. T).

If $S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$, then the set

$$\mathcal{P}_{\Phi}^{\alpha}(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

is in bijection with the set (which does not depend on α !)

$$\mathcal{P}_{\Phi}^1(J) = \{i_0 \cdot u : u \in \mathcal{T} \text{ such that } i_0 u \text{ is totally imaginary and } \Phi\text{-positive}\}.$$

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Theorem (1)

Denote by $S_{\mathbb{R}}^*$ (resp. $T_{\mathbb{R}}^*$) the group of totally real units of S (resp. T).
If $S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$, then the set

$$\mathcal{P}_{\Phi}^{\alpha}(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

is in bijection with the set (which does not depend on α !)

$$\mathcal{P}_{\Phi}^1(J) = \{i_0 \cdot u : u \in \mathcal{T} \text{ such that } i_0 u \text{ is totally imaginary and } \Phi\text{-positive}\}.$$

Corollary

If $S = \mathbb{Z}[F, V]$ (eg. $AV_h(p)$ is ordinary or almost-ordinary) then we can ignore α .

Effective Results : when can we ignore α ?

Assume A_0 admits a canonical lifting. Put $S := \text{End}(A_0)$

Let B_0 be isogenous to A_0 . Put $T = \text{End}(B_0)$.

Theorem (1)

Denote by $S_{\mathbb{R}}^*$ (resp. $T_{\mathbb{R}}^*$) the group of totally real units of S (resp. T).
If $S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$, then the set

$$\mathcal{P}_{\Phi}^{\alpha}(J) := \{i_0 \cdot u : u \in \mathcal{T} \text{ s.t. } \alpha^{-1} i_0 u \text{ is tot. imaginary and } \Phi\text{-positive}\}$$

is in bijection with the set (which does not depend on α !)

$$\mathcal{P}_{\Phi}^1(J) = \{i_0 \cdot u : u \in \mathcal{T} \text{ such that } i_0 u \text{ is totally imaginary and } \Phi\text{-positive}\}.$$

Corollary

If $S = \mathbb{Z}[F, V]$ (eg. $AV_h(p)$ is ordinary or almost-ordinary) then we can ignore α . *We recover Deligne+Howe and Oswal-Shankar*

Effective Results II

Theorem (2)

Assume that there are r isomorphism classes of abelian varieties in $AV_h(p)$ with endomorphism ring T , represented under \mathcal{G} by the fractional ideals I_1, \dots, I_r . For any CM-type Φ' , we put

$$\mathcal{P}_{\Phi'}^1(I_i) = \{i_0 \cdot u : u \in \mathcal{T} \text{ such that } i_0 u \text{ is totally imaginary and } \Phi'\text{-positive} \}.$$

If there exists a non-negative integer N such that for every CM-type Φ' we have

$$|\mathcal{P}_{\Phi'}^1(I_1)| + \dots + |\mathcal{P}_{\Phi'}^1(I_r)| = N$$

then there are exactly N isomorphism classes of principally polarized abelian varieties with endomorphism ring T .

Proof.

- Consider the association $\Phi' \mapsto b$ where $b \in L^*$ is tot. imaginary and Φ' -positive.
- We can go back: for every b tot. imaginary there exists a unique CM-type Φ_b s.t. b is Φ_b -positive.
- Hence the totally real elements of L^* acts on the set of CM-types.
- If $\Phi = \Phi_b$ is the CM-type for which we have a canonical lift (as before) then $\mathcal{P}_{\Phi_b}^\alpha(l_i) \longleftrightarrow \mathcal{P}_{\Phi_{ab}}^1(l_i)$.
- If the we get the 'same sum' (over the l_i 's) for every CM-type we know that the result must be the correct one!



Note: even if the sum is not the same for all Φ' 's then we know that one of the outputs is the correct one!

When can we lift up to isogeny?

Definition (Chai-Conrad-Oort)

Let Φ be a p -adic CM-type for a CM-field $L = \mathbb{Q}(F)$.

When can we lift up to isogeny?

Definition (Chai-Conrad-Oort)

Let Φ be a p -adic CM-type for a CM-field $L = \mathbb{Q}(F)$. The pair (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F if the following conditions are met:

When can we lift up to isogeny?

Definition (Chai-Conrad-Oort)

Let Φ be a p -adic CM-type for a CM-field $L = \mathbb{Q}(F)$. The pair (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F if the following conditions are met:

1. The **Shimura-Taniyama formula** holds for F : for every place v of L above p , we have

$$\frac{\text{ord}_v(F)}{\text{ord}_v(q)} = \frac{\#\{\varphi \in \Phi \text{ s.t. } \varphi \text{ induces } v\}}{[L_v : \mathbb{Q}_p]}.$$

When can we lift up to isogeny?

Definition (Chai-Conrad-Oort)

Let Φ be a p -adic CM-type for a CM-field $L = \mathbb{Q}(F)$. The pair (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F if the following conditions are met:

1. The **Shimura-Taniyama formula** holds for F : for every place v of L above p , we have

$$\frac{\text{ord}_v(F)}{\text{ord}_v(q)} = \frac{\#\{\varphi \in \Phi \text{ s.t. } \varphi \text{ induces } v\}}{[L_v : \mathbb{Q}_p]}.$$

2. Let E be the reflex field attached to (L, Φ) , and let v be the induced p -adic place of E . Then the **residue field** k_v of $\mathcal{O}_{E,v}$ can be realized as a **subfield** of \mathbb{F}_q .

When can we lift up to isogeny?

Theorem (Chai-Conrad-Oort)

Assume that (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F , that is,

- 1 Φ satisfies the Shimura-Taniyama formula for F , and
- 2 the reflex field E has residue field $k_E \subseteq \mathbb{F}_q$.

When can we lift up to isogeny?

Theorem (Chai-Conrad-Oort)

Assume that (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F , that is,

- 1 Φ satisfies the Shimura-Taniyama formula for F , and
- 2 the reflex field E has residue field $k_E \subseteq \mathbb{F}_q$.

Then we can **canonically lift** an abelian variety A_0 with $\mathcal{O}_L = \text{End}(A_0)$.

When can we lift up to isogeny?

Theorem (Chai-Conrad-Oort)

Assume that (L, Φ) satisfies the **Residual Reflex Condition** w.r.t. F , that is,

- 1 Φ satisfies the Shimura-Taniyama formula for F , and
- 2 the reflex field E has residue field $k_E \subseteq \mathbb{F}_q$.

Then we can **canonically lift** an abelian variety A_0 with $\mathcal{O}_L = \text{End}(A_0)$.

- If there is a separable isogeny $A_0 \rightarrow A'_0$ then A'_0 admits a canonical lifting (useful in combination with Thm 1).

We run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

We run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4. For example:

squarefree dimension 3			$p = 2$	$p = 3$	$p = 5$	$p = 7$
total			185	621	2863	7847
ordinary			82	390	2280	6700
almost ordinary			58	170	474	996
p -rank 1	no RRC		0	0	0	0
	yes RRC	Thm 1 yes	20	26	76	118
		Thm 1 no	4	16	12	8
p -rank 0	no RRC		0	3	2	1
	yes RRC	Thm 1 yes	20	15	17	23
		Thm 1 no	1	1	2	1

We run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4. For example:

squarefree dimension 3			$p = 2$	$p = 3$	$p = 5$	$p = 7$
total			185	621	2863	7847
ordinary			82	390	2280	6700
almost ordinary			58	170	474	996
p -rank 1	no RRC		0	0	0	0
	yes RRC	Thm 1 yes	20	26	76	118
		Thm 1 no	4	16	12	8
p -rank 0	no RRC		0	3	2	1
	yes RRC	Thm 1 yes	20	15	17	23
		Thm 1 no	1	1	2	1

Among the 45 isogeny classes which we cannot 'handle' with Thm 1, we can compute the number of PPAV for 32 of them using Thm 2. For the remaining 13 (all over \mathbb{F}_2 and \mathbb{F}_3) we only get partial info.

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

squarefree dimension 4			$p = 2$	$p = 3$
total			1431	10453
ordinary			656	6742
almost ordinary			392	2506
p -rank 2	no RRC		0	0
	yes RRC	Thm 1 yes	149	500
		Thm 1 no	49	312
p -rank 1	no RRC		6	36
	yes RRC	Thm 1 yes	80	184
		Thm 1 no	14	40
p -rank 0	no RRC		3	6
	yes RRC	Thm 1 yes	73	88
		Thm 1 no	9	39

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

squarefree dimension 4			$p = 2$	$p = 3$
total			1431	10453
ordinary			656	6742
almost ordinary			392	2506
p -rank 2	no RRC		0	0
	yes RRC	Thm 1 yes	149	500
		Thm 1 no	49	312
p -rank 1	no RRC		6	36
	yes RRC	Thm 1 yes	80	184
		Thm 1 no	14	40
p -rank 0	no RRC		3	6
	yes RRC	Thm 1 yes	73	88
		Thm 1 no	9	39

Thm 1 ($S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$) doesn't handle $72/\mathbb{F}_2$ and $391/\mathbb{F}_3$.

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

squarefree dimension 4			$p = 2$	$p = 3$
total			1431	10453
ordinary			656	6742
almost ordinary			392	2506
p -rank 2	no RRC		0	0
	yes RRC	Thm 1 yes	149	500
		Thm 1 no	49	312
p -rank 1	no RRC		6	36
	yes RRC	Thm 1 yes	80	184
		Thm 1 no	14	40
p -rank 0	no RRC		3	6
	yes RRC	Thm 1 yes	73	88
		Thm 1 no	9	39

Thm 1 ($S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$) doesn't handle $72/\mathbb{F}_2$ and $391/\mathbb{F}_3$. Out of these, we can use Thm 2 for $20/\mathbb{F}_2$ and $214/\mathbb{F}_3$.

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

squarefree dimension 4			$p = 2$	$p = 3$
total			1431	10453
ordinary			656	6742
almost ordinary			392	2506
p -rank 2	no RRC		0	0
	yes RRC	Thm 1 yes	149	500
		Thm 1 no	49	312
p -rank 1	no RRC		6	36
	yes RRC	Thm 1 yes	80	184
		Thm 1 no	14	40
p -rank 0	no RRC		3	6
	yes RRC	Thm 1 yes	73	88
		Thm 1 no	9	39

Thm 1 ($S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$) doesn't handle $72/\mathbb{F}_2$ and $391/\mathbb{F}_3$. Out of these, we can use Thm 2 for $20/\mathbb{F}_2$ and $214/\mathbb{F}_3$. For the remaining $52/\mathbb{F}_2$ and $171/\mathbb{F}_3$ we can only get information about certain endomorphism rings (723 out of 946 and 3481 out of 4636, respectively).

We have run computations over all squarefree isogeny classes over small prime fields of dim 2,3 and 4.

squarefree dimension 4			$p = 2$	$p = 3$
total			1431	10453
ordinary			656	6742
almost ordinary			392	2506
p -rank 2	no RRC		0	0
	yes RRC	Thm 1 yes	149	500
		Thm 1 no	49	312
p -rank 1	no RRC		6	36
	yes RRC	Thm 1 yes	80	184
		Thm 1 no	14	40
p -rank 0	no RRC		3	6
	yes RRC	Thm 1 yes	73	88
		Thm 1 no	9	39

Thm 1 ($S_{\mathbb{R}}^* \subseteq T_{\mathbb{R}}^*$) doesn't handle $72/\mathbb{F}_2$ and $391/\mathbb{F}_3$. Out of these, we can use Thm 2 for $20/\mathbb{F}_2$ and $214/\mathbb{F}_3$. For the remaining $52/\mathbb{F}_2$ and $171/\mathbb{F}_3$ we can only get information about certain endomorphism rings (723 out of 946 and 3481 out of 4636, respectively). Also there are $9/\mathbb{F}_3$ for which the computations of the isomorphism classes of unpolarized abelian varieties is not over yet.

Thank you!