


Abelian varieties over finite fields

Stefano Marseglia


UPF - Gaati Lab

14/02/2024


What do I do for a living?




What do I do for a living?




What do I do for a living?



What do I do for a living?



What do I do for a living?



Abelian varieties: what are they ?


Abelian varieties are connected projective group varieties.

Abelian varieties: what are they ?

Abelian varieties are connected projective group varieties.

Abelian varieties of dim. 1
are called **elliptic curves**.

Eg: over \mathbb{R} , $y^2 = x^3 - x + 1$



Abelian varieties: what are they ?


Abelian varieties are connected projective group varieties.

Abelian varieties of dim. 1
are called **elliptic curves**.

Eg: over \mathbb{R} , $y^2 = x^3 - x + 1$

We can add points:

$P, Q \rightsquigarrow P \oplus Q$



Abelian varieties: what are they ?

Abelian varieties are connected projective group varieties.

Abelian varieties of dim. 1
are called **elliptic curves**.


Eg: over \mathbb{R} , $y^2 = x^3 - x + 1$

We can add points:

$P, Q \rightsquigarrow P \oplus Q$

Equations are impractical in
dim ≥ 2 .

We need a better way to
represent them...



Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.
- T admits a non-degenerate Riemann form \longleftrightarrow polarization.

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.
- T admits a non-degenerate Riemann form \longleftrightarrow polarization.
- In fact, $A \mapsto A(\mathbb{C})$ induces an **equivalence** of categories:

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting} \right. \\ \left. \text{a Riemann form} \right\}.$$

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.
- T admits a non-degenerate Riemann form \longleftrightarrow polarization.
- In fact, $A \mapsto A(\mathbb{C})$ induces an **equivalence** of categories:

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting a Riemann form} \right\}.$$

- In **char. $p > 0$** such an equivalence **cannot exist**: there are (supersingular) elliptic curves with quaternionic endomorphism algebras.

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.
- T admits a non-degenerate Riemann form \longleftrightarrow polarization.
- In fact, $A \mapsto A(\mathbb{C})$ induces an **equivalence** of categories:

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting a Riemann form} \right\}.$$

- In **char. $p > 0$** such an equivalence **cannot exist**: there are (supersingular) elliptic curves with quaternionic endomorphism algebras.
- Nevertheless, over finite fields, we obtain analogous results if we restrict ourselves to certain **subcategories** of AVs...

Abelian varieties over \mathbb{C} vs \mathbb{F}_q

- Let A/\mathbb{C} be an abelian variety of dimension g .
- Then $A(\mathbb{C})$ is a **torus**: $T := \mathbb{C}^g / \Lambda$, where $\Lambda \simeq_{\mathbb{Z}} \mathbb{Z}^{2g}$.
- T admits a non-degenerate Riemann form \longleftrightarrow polarization.
- In fact, $A \mapsto A(\mathbb{C})$ induces an **equivalence** of categories:

$$\{\text{abelian varieties } / \mathbb{C}\} \longleftrightarrow \left\{ \mathbb{C}^g / \Lambda \text{ with } \Lambda \simeq \mathbb{Z}^{2g} \text{ admitting a Riemann form} \right\}.$$

- In **char. $p > 0$** such an equivalence **cannot exist**: there are (supersingular) elliptic curves with quaternionic endomorphism algebras.
- Nevertheless, over finite fields, we obtain analogous results if we restrict ourselves to certain **subcategories** of AVs...
- ... which we are going to use to **classify the AVs up to isomorphism**.

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.
- A/\mathbb{F}_q comes with a **Frobenius** endomorphism,

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.
- A/\mathbb{F}_q comes with a **Frobenius** endomorphism, that induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.
- A/\mathbb{F}_q comes with a **Frobenius** endomorphism, that induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a **q -Weil** polynomial and **isogeny invariant**.

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.
- A/\mathbb{F}_q comes with a **Frobenius** endomorphism, that induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a **q -Weil** polynomial and **isogeny invariant**.
- By **Honda-Tate** theory, the association

$$\text{isogeny class of } A \longmapsto h_A(x)$$

is injective and allows us to **enumerate** all AVs up to isogeny.

Isogeny classification over \mathbb{F}_q

- An **isogeny** $A \rightarrow B$ is a surjective morphism with finite kernel.
- A/\mathbb{F}_q comes with a **Frobenius** endomorphism, that induces an action

$$\text{Frob}_A : T_\ell A \rightarrow T_\ell A \text{ for any } \ell \neq p,$$

where $T_\ell(A) = \varprojlim_\ell A[\ell^n] \simeq \mathbb{Z}_\ell^{2g}$.

- $h_A(x) := \text{char}(\text{Frob}_A)$ is a **q -Weil** polynomial and **isogeny invariant**.
- By **Honda-Tate** theory, the association

$$\text{isogeny class of } A \mapsto h_A(x)$$

is injective and allows us to **enumerate** all AVs up to isogeny.

- Also, $h_A(x)$ is squarefree $\iff \text{End}(A)$ is commutative.

Deligne's equivalence

Recall: A/\mathbb{F}_q is **ordinary** if half of the p -adic roots of h_A are units.

Deligne's equivalence

Recall: A/\mathbb{F}_q is **ordinary** if half of the p -adic roots of h_A are units.

Theorem (Deligne '69)

Let $q = p^r$, with p a prime. There is an *equivalence* of categories:

$$\{ \text{Ordinary abelian varieties over } \mathbb{F}_q \} \quad A$$

Deligne's equivalence

Recall: A/\mathbb{F}_q is **ordinary** if half of the p -adic roots of h_A are units.

Theorem (Deligne '69)

Let $q = p^r$, with p a prime. There is an *equivalence* of categories:

$$\begin{array}{ccc} \{ \text{Ordinary abelian varieties over } \mathbb{F}_q \} & & A \\ \updownarrow & & \downarrow \\ \left\{ \text{pairs } (T, F), \text{ where } T \simeq_{\mathbb{Z}} \mathbb{Z}^{2g} \text{ and } T \xrightarrow{F} T \text{ s.t.} \right. & & \end{array}$$

Deligne's equivalence

Recall: A/\mathbb{F}_q is **ordinary** if half of the p -adic roots of h_A are units.

Theorem (Deligne '69)

Let $q = p^r$, with p a prime. There is an *equivalence* of categories:

$$\begin{array}{ccc} \{ \text{Ordinary abelian varieties over } \mathbb{F}_q \} & & A \\ \downarrow & & \downarrow \\ \left\{ \begin{array}{l} \text{pairs } (T, F), \text{ where } T \simeq_{\mathbb{Z}} \mathbb{Z}^{2g} \text{ and } T \xrightarrow{F} T \text{ s.t.} \\ - F \otimes \mathbb{Q} \text{ is semisimple} \\ - \text{the roots of } \text{char}_{F \otimes \mathbb{Q}}(x) \text{ have abs. value } \sqrt{q} \\ - \text{half of them are } p\text{-adic units} \\ - \exists V : T \rightarrow T \text{ such that } FV = VF = q \end{array} \right\} & & (T(A), F(A)) \end{array}$$

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.
- Put $V = q/F$. Deligne's equivalence induces:

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.
- Put $V = q/F$. Deligne's equivalence induces:

Theorem

$$\begin{array}{c} \{abelian\ varieties\ over\ \mathbb{F}_q\ in\ \mathcal{C}_h\} / \\ \updownarrow \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, V] \subset K\} / \end{array}$$

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.
- Put $V = q/F$. Deligne's equivalence induces:

Theorem

$$\begin{array}{c} \{abelian\ varieties\ over\ \mathbb{F}_q\ in\ \mathcal{C}_h\} / \simeq \\ \updownarrow \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, V] \subset K\} / \simeq \end{array}$$

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.
- Put $V = q/F$. Deligne's equivalence induces:

Theorem

$$\begin{array}{ccc} \{abelian\ varieties\ over\ \mathbb{F}_q\ in\ \mathcal{C}_h\} / \simeq & & \\ \updownarrow & & \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, V] \subset K\} / \simeq & =: & \text{ICM}(\mathbb{Z}[F, V]) \\ & & \text{ideal class monoid} \end{array}$$

Squarefree case

- Fix an **ordinary squarefree** q -Weil polynomial h :
- \rightsquigarrow an isogeny class $\mathcal{C}_h/\mathbb{F}_q$.
- Put $K := \mathbb{Q}[x]/(h) = \mathbb{Q}[F]$, an étale algebra = product of number fields.
- Put $V = q/F$. Deligne's equivalence induces:

Theorem

$$\begin{array}{ccc} \{abelian\ varieties\ over\ \mathbb{F}_q\ in\ \mathcal{C}_h\} / \simeq & & \\ \updownarrow & & \\ \{fractional\ ideals\ of\ \mathbb{Z}[F, V] \subset K\} / \simeq & =: & \text{ICM}(\mathbb{Z}[F, V]) \\ & & \text{ideal class monoid} \end{array}$$

- **Problem:** $\mathbb{Z}[F, V]$ might not be maximal \rightsquigarrow **non-invertible** ideals.

ICM : Ideal Class Monoid

Let R be an **order** in an étale \mathbb{Q} -algebra K .

ICM : Ideal Class Monoid

Let R be an **order** in an étale \mathbb{Q} -algebra K .

- Recall: for **fractional R -ideals** I and J

$$I \simeq_R J \iff \exists x \in K^\times \text{ s.t. } xI = J$$

ICM : Ideal Class Monoid

Let R be an **order** in an étale \mathbb{Q} -algebra K .

- Recall: for **fractional R -ideals** I and J

$$I \simeq_R J \iff \exists x \in K^\times \text{ s.t. } xI = J$$

- We have

$$\mathrm{ICM}(R) \supseteq \mathrm{Pic}(R) = \{\text{invertible fractional } R\text{-ideals}\} / \simeq_R$$

with equality \Updownarrow iff $R = \mathcal{O}_K$

ICM : Ideal Class Monoid

Let R be an **order** in an étale \mathbb{Q} -algebra K .

- Recall: for **fractional R -ideals** I and J

$$I \simeq_R J \iff \exists x \in K^\times \text{ s.t. } xI = J$$

- We have

$$\mathrm{ICM}(R) \supseteq \mathrm{Pic}(R) = \{\text{invertible fractional } R\text{-ideals}\} / \simeq_R$$

with equality \Updownarrow iff $R = \mathcal{O}_K$

- ...and actually

$$\mathrm{ICM}(R) \supseteq \bigsqcup_{\substack{R \subseteq S \subseteq \mathcal{O}_K \\ \text{over-orders}}} \mathrm{Pic}(S) \quad \text{with equality iff } R \text{ is Bass}$$

ICM : Ideal Class Monoid

Let R be an **order** in an étale \mathbb{Q} -algebra K .

- Recall: for **fractional R -ideals** I and J

$$I \simeq_R J \iff \exists x \in K^\times \text{ s.t. } xI = J$$

- We have

$$\mathrm{ICM}(R) \supseteq \mathrm{Pic}(R) = \{\text{invertible fractional } R\text{-ideals}\} / \simeq_R$$

with equality \Updownarrow iff $R = \mathcal{O}_K$

- ...and actually

$$\mathrm{ICM}(R) \supseteq \bigsqcup_{\substack{R \subseteq S \subseteq \mathcal{O}_K \\ \text{over-orders}}} \mathrm{Pic}(S) \quad \text{with equality iff } R \text{ is Bass}$$

- Hofmann-Sircana '19: computation of over-orders.

simplify the problem

Study the isomorphism problem locally: (Dade, Taussky, Zassenhaus '62)

simplify the problem

Study the isomorphism problem locally: (Dade, Taussky, Zassenhaus '62)

- **weak equivalence:**

$$I_{\mathfrak{p}} \simeq_{R_{\mathfrak{p}}} J_{\mathfrak{p}} \text{ for every } \mathfrak{p} \in \mathrm{mSpec}(R)$$

simplify the problem

Study the isomorphism problem locally: (Dade, Taussky, Zassenhaus '62)

- **weak equivalence:**

$$I_{\mathfrak{p}} \simeq_{R_{\mathfrak{p}}} J_{\mathfrak{p}} \text{ for every } \mathfrak{p} \in \text{mSpec}(R)$$



$$1 \in (I : J)(J : I) \quad \text{easy to check!}$$

simplify the problem

Study the isomorphism problem locally: (Dade, Taussky, Zassenhaus '62)

- **weak equivalence:**

$$I_{\mathfrak{p}} \simeq_{R_{\mathfrak{p}}} J_{\mathfrak{p}} \text{ for every } \mathfrak{p} \in \mathrm{mSpec}(R)$$



$$1 \in (I : J)(J : I) \quad \text{easy to check!}$$

- Let $\mathcal{W}(R)$ be the set of weak eq. classes...

simplify the problem

Study the isomorphism problem locally: (Dade, Taussky, Zassenhaus '62)

- **weak equivalence:**

$$I_{\mathfrak{p}} \simeq_{R_{\mathfrak{p}}} J_{\mathfrak{p}} \text{ for every } \mathfrak{p} \in \mathrm{mSpec}(R)$$



$$1 \in (I : J)(J : I) \quad \text{easy to check!}$$

- Let $\mathcal{W}(R)$ be the set of weak eq. classes...
...whose representatives can be found in

$$\left\{ \text{sub-}R\text{-modules of } \mathcal{O}_K / \mathfrak{f}_R \right\} \quad \text{finite! and most of the time not-too-big ...}$$

where $\mathfrak{f}_R = (R : \mathcal{O}_K)$ is the conductor of R .

Compute $ICM(R)$

Compute $\text{ICM}(R)$

Partition w.r.t. the multiplier ring:

$$\mathcal{W}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \mathcal{W}_S(R)$$

$$\text{ICM}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \text{ICM}_S(R)$$

Compute $\text{ICM}(R)$

Partition w.r.t. the multiplier ring:

$$\mathcal{W}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \mathcal{W}_S(R)$$

$$\text{ICM}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \text{ICM}_S(R)$$

the “pedix” $-_S$ means
“only classes with multiplier ring S ”

Compute $\text{ICM}(R)$

Partition w.r.t. the multiplier ring:

$$\mathcal{W}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \mathcal{W}_S(R)$$

$$\text{ICM}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \text{ICM}_S(R)$$

the “pedix” $-_S$ means
“only classes with multiplier ring S ”

Theorem (M.)

For every over-order S of R , $\text{Pic}(S)$ acts *freely* on $\text{ICM}_S(R)$ and

$$\mathcal{W}_S(R) = \text{ICM}_S(R) / \text{Pic}(S)$$

Compute $\text{ICM}(R)$

Partition w.r.t. the multiplier ring:

$$\mathcal{W}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \mathcal{W}_S(R)$$

$$\text{ICM}(R) = \bigsqcup_{R \subseteq S \subseteq \mathcal{O}_K} \text{ICM}_S(R)$$

the “pedix” $-_S$ means
“only classes with multiplier ring S ”

Theorem (M.)

For every over-order S of R , $\text{Pic}(S)$ acts *freely* on $\text{ICM}_S(R)$ and

$$\mathcal{W}_S(R) = \text{ICM}_S(R) / \text{Pic}(S)$$

Repeat for every $R \subseteq S \subseteq \mathcal{O}_K$:

$$\rightsquigarrow \text{ICM}(R).$$

To sum up:

- To sum up:

To sum up:

- To sum up:
- Given a **ordinary squarefree** q -Weil polynomial $h \dots$

To sum up:

- To sum up:
- Given a **ordinary squarefree** q -Weil polynomial h ...
- ... \rightsquigarrow algorithm to **compute the isomorphism classes** of AVs in the isogeny class \mathcal{C}_h .

To sum up:

- To sum up:
- Given a **ordinary squarefree** q -Weil polynomial h ...
- ... \rightsquigarrow algorithm to **compute the isomorphism classes** of AVs in the isogeny class \mathcal{C}_h .

Remark

*Let \mathcal{C}_h be a **squarefree** isogeny classes over the **prime field** \mathbb{F}_p . Building on work by Centeleghe-Stix, we get a bijection between the isomorphism classes of AVs in \mathcal{C}_h and the ideal class monoid of $\mathbb{Z}[F, V]$, as above. But the functor is completely different! (eg. It is contravariant)*

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

- $A^\vee \leftrightarrow \bar{I}^t := \{\bar{x} \in K : \text{Tr}(xI) \subseteq \mathbb{Z}\}.$

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

- $A^\vee \leftrightarrow \bar{I}^t := \{\bar{x} \in K : \text{Tr}(xI) \subseteq \mathbb{Z}\}.$
- a polarization μ of A corresponds to a $\lambda \in K^\times$ such that
 - $\lambda I \subseteq \bar{I}^t$ (isogeny);
 - λ is **totally imaginary** ($\bar{\lambda} = -\lambda$);
 - λ is Φ -positive, where Φ is a CM-type of K satisf. the **Shimura-Taniyama** formula.

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

- $A^\vee \leftrightarrow \bar{I}^t := \{\bar{x} \in K : \text{Tr}(xI) \subseteq \mathbb{Z}\}.$
- a polarization μ of A corresponds to a $\lambda \in K^\times$ such that
 - $\lambda I \subseteq \bar{I}^t$ (isogeny);
 - λ is **totally imaginary** ($\bar{\lambda} = -\lambda$);
 - λ is Φ -positive, where Φ is a CM-type of K satisf. the **Shimura-Taniyama** formula.

Also: $\deg \mu = [\bar{I}^t : \lambda I].$

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

- $A^\vee \leftrightarrow \bar{I}^t := \{\bar{x} \in K : \text{Tr}(xI) \subseteq \mathbb{Z}\}.$
- a polarization μ of A corresponds to a $\lambda \in K^\times$ such that
 - $\lambda I \subseteq \bar{I}^t$ (isogeny);
 - λ is **totally imaginary** ($\bar{\lambda} = -\lambda$);
 - λ is Φ -positive, where Φ is a CM-type of K satisf. the **Shimura-Taniyama** formula.

Also: $\deg \mu = [\bar{I}^t : \lambda I].$

- if $(A, \mu) \leftrightarrow (I, \lambda)$ is a princ. polarized ab. var. and $S = (I : I)$ then

$$\left\{ \begin{array}{l} \text{non-isomorphic princ.} \\ \text{polarizations of } A \end{array} \right\} \longleftrightarrow \frac{\{\text{totally positive } u \in S^\times\}}{\{\bar{v}v : v \in S^\times\}},$$

Dual varieties and Polarizations

Howe described **dual** varieties and **polarizations** on Deligne modules.

Theorem

Let $A \in \mathcal{C}_h$ with h ordinary and squarefree. If $A \leftrightarrow I$, then:

- $A^\vee \leftrightarrow \bar{I}^t := \{\bar{x} \in K : \text{Tr}(xI) \subseteq \mathbb{Z}\}.$
- a polarization μ of A corresponds to a $\lambda \in K^\times$ such that
 - $\lambda I \subseteq \bar{I}^t$ (isogeny);
 - λ is **totally imaginary** ($\bar{\lambda} = -\lambda$);
 - λ is Φ -positive, where Φ is a CM-type of K satisf. the **Shimura-Taniyama** formula.

Also: $\deg \mu = [\bar{I}^t : \lambda I].$

- if $(A, \mu) \leftrightarrow (I, \lambda)$ is a princ. polarized ab. var. and $S = (I : I)$ then

$$\left\{ \begin{array}{l} \text{non-isomorphic princ.} \\ \text{polarizations of } A \end{array} \right\} \longleftrightarrow \frac{\{\text{totally positive } u \in S^\times\}}{\{v\bar{v} : v \in S^\times\}},$$

- and $\text{Aut}(A, \mu) = \{\text{torsion units of } S\}.$

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.
- 2 Loop over the representatives u of the finite quotient

$$\frac{S^\times}{\{v\bar{v} : v \in S^\times\}}.$$

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.
- 2 Loop over the representatives u of the finite quotient

$$\frac{S^\times}{\{v\bar{v} : v \in S^\times\}}.$$

- 3 If $\lambda := i_0 u$ is totally imaginary and Φ -positive ...

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.
- 2 Loop over the representatives u of the finite quotient

$$\frac{S^\times}{\{v\bar{v} : v \in S^\times\}}.$$

- 3 If $\lambda := i_0 u$ is totally imaginary and Φ -positive ...
- 4 ... then we have one principal polarization.

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.
- 2 Loop over the representatives u of the finite quotient

$$\frac{S^\times}{\{v\bar{v} : v \in S^\times\}}.$$

- 3 If $\lambda := i_0 u$ is totally imaginary and Φ -positive ...
- 4 ... then we have one principal polarization.
- 5 By the previous Theorem, we have all princ. polarizations up to isom.

Principal Polarizations

We have an **algorithm** to enumerate principal polarizations up to isomorphism:

- 1 Compute i_0 such that $i_0 I = \bar{I}^t$.
- 2 Loop over the representatives u of the finite quotient

$$\frac{S^\times}{\{v\bar{v} : v \in S^\times\}}.$$

- 3 If $\lambda := i_0 u$ is totally imaginary and Φ -positive ...
- 4 ... then we have one principal polarization.
- 5 By the previous Theorem, we have all princ. polarizations up to isom.

Can modify to compute polarizations of any degree.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.
- Let F be a root of $h(x)$ and put $R := \mathbb{Z}[F, 3/F] \subset \mathbb{Q}(F)$.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.
- Let F be a root of $h(x)$ and put $R := \mathbb{Z}[F, 3/F] \subset \mathbb{Q}(F)$.
- 8 over-orders of R : two of them are not Gorenstein.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.
- Let F be a root of $h(x)$ and put $R := \mathbb{Z}[F, 3/F] \subset \mathbb{Q}(F)$.
- 8 over-orders of R : two of them are not Gorenstein.
- $\# \text{ICM}(R) = 18 \rightsquigarrow 18$ isom. classes of AV in the isogeny class.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.
- Let F be a root of $h(x)$ and put $R := \mathbb{Z}[F, 3/F] \subset \mathbb{Q}(F)$.
- 8 over-orders of R : two of them are not Gorenstein.
- $\#\text{ICM}(R) = 18 \rightsquigarrow 18$ isom. classes of AV in the isogeny class.
- 5 are not invertible in their multiplier ring.

Example

- Let $h(x) = x^8 - 5x^7 + 13x^6 - 25x^5 + 44x^4 - 75x^3 + 117x^2 - 135x + 81$,
LMFDB label: 4.3.af_n_az_bs.
- \rightsquigarrow isogeny class of a simple ordinary abelian varieties over \mathbb{F}_3 of dimension 4.
- Let F be a root of $h(x)$ and put $R := \mathbb{Z}[F, 3/F] \subset \mathbb{Q}(F)$.
- 8 over-orders of R : two of them are not Gorenstein.
- $\#\text{ICM}(R) = 18 \rightsquigarrow 18$ isom. classes of AV in the isogeny class.
- 5 are not invertible in their multiplier ring.
- More info at
https://abvar.lmfdb.xyz/Variety/Abelian/Fq/4/3/af_n_az_bs

Example

Concretely:

$$\begin{aligned} I_1 = & 2645633792595191\mathbb{Z} \oplus (F + 836920075614551)\mathbb{Z} \oplus (F^2 + 1474295643839839)\mathbb{Z} \oplus \\ & \oplus (F^3 + 1372829830503387)\mathbb{Z} \oplus (F^4 + 1072904687510)\mathbb{Z} \oplus \\ & \oplus \frac{1}{3}(F^5 + F^4 + F^3 + 2F^2 + 2F + 6704806986143610)\mathbb{Z} \oplus \\ & \oplus \frac{1}{9}(F^6 + F^5 + F^4 + 8F^3 + 2F^2 + 2991665243621169)\mathbb{Z} \oplus \\ & \oplus \frac{1}{27}(F^7 + F^6 + F^5 + 17F^4 + 20F^3 + 9F^2 + 68015312518722201)\mathbb{Z} \end{aligned}$$

principal polarizations:

$$\begin{aligned} x_{1,1} = & \frac{1}{27}(-121922F^7 + 588604F^6 - 1422437F^5 + \\ & + 1464239F^4 + 1196576F^3 - 7570722F^2 + 15316479F - 12821193) \\ x_{1,2} = & \frac{1}{27}(3015467F^7 - 17689816F^6 + 35965592F^5 - \\ & - 64660346F^4 + 121230619F^3 - 191117052F^2 + 315021546F - 300025458) \end{aligned}$$

$$\text{End}(I_1) = R$$

$$\# \text{Aut}(I_1, x_{1,1}) = \# \text{Aut}(I_1, x_{1,2}) = 2$$

- The equivalence is not just useful to classify the AVs!

- The equivalence is not just useful to classify the AVs!
- It can be used to compute polarizations, isogenies, and ...

- The equivalence is not just useful to classify the AVs!
- It can be used to compute polarizations, isogenies, and ...
- ... group of \mathbb{F}_q -points.

- The equivalence is not just useful to classify the AVs!
- It can be used to compute polarizations, isogenies, and ...
- ... group of \mathbb{F}_q -points.
- In the rest of the talk, we will prove

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group.

- The equivalence is not just useful to classify the AVs!
- It can be used to compute polarizations, isogenies, and ...
- ... group of \mathbb{F}_q -points.
- In the rest of the talk, we will prove

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

$$A(\mathbb{F}_q) \simeq \frac{J}{(1-F)J}.$$

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

$$A(\mathbb{F}_q) \simeq \frac{J}{(1-F)J}.$$

Proposition (M.-Springer)

Every ordinary squarefree isogeny class contains a cyclic abelian variety.

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

$$A(\mathbb{F}_q) \simeq \frac{J}{(1-F)J}.$$

Proposition (M.-Springer)

Every ordinary squarefree isogeny class contains a cyclic abelian variety.

Proof: Take $A \longleftrightarrow J = \mathbb{Z}[F, q/F]$.

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

$$A(\mathbb{F}_q) \simeq \frac{J}{(1-F)J}.$$

Proposition (M.-Springer)

Every ordinary squarefree isogeny class contains a cyclic abelian variety.

Proof: Take $A \longleftrightarrow J = \mathbb{Z}[F, q/F]$.

$$A(\mathbb{F}_q) \simeq \frac{\mathbb{Z}[F, q/F]}{(1-F)} \simeq \frac{\mathbb{Z}[x, y]}{(h(x), xy - q, 1-x)}$$

Cyclic abelian varieties

Recall that we have an equivalence

$$\mathcal{C}_h \longleftrightarrow \{\text{fractional ideals of } \mathbb{Z}[F, q/F]\}.$$

Corollary

If A corresponds to the fractional $\mathbb{Z}[F, q/F]$ -ideal J then

$$A(\mathbb{F}_q) \simeq \frac{J}{(1-F)J}.$$

Proposition (M.-Springer)

Every ordinary squarefree isogeny class contains a cyclic abelian variety.

Proof: Take $A \longleftrightarrow J = \mathbb{Z}[F, q/F]$.

$$A(\mathbb{F}_q) \simeq \frac{\mathbb{Z}[F, q/F]}{(1-F)} \simeq \frac{\mathbb{Z}[x, y]}{(h(x), xy - q, 1-x)} \simeq \frac{\mathbb{Z}}{h(1)\mathbb{Z}}. \quad \square$$

Number of points

Theorem (Howe-Kedlaya)

Let $m \in \mathbb{Z}_{\geq 0}$. Then there is a squarefree ordinary A/\mathbb{F}_2 such that $\#A(\mathbb{F}_2) = m$.

Number of points

Theorem (Howe-Kedlaya)

Let $m \in \mathbb{Z}_{\geq 0}$. Then there is a squarefree ordinary A/\mathbb{F}_2 such that $\#A(\mathbb{F}_2) = m$.

Theorem (van Bommel-Costa-Li-Poonen-Smith)

Let $m \in \mathbb{Z}_{\geq 0}$ and k be \mathbb{F}_3 or \mathbb{F}_5 . Then there is a squarefree ordinary A/k such that $\#A(k) = m$.

Number of points

Theorem (Howe-Kedlaya)

Let $m \in \mathbb{Z}_{\geq 0}$. Then there is a squarefree ordinary A/\mathbb{F}_2 such that $\#A(\mathbb{F}_2) = m$.

Theorem (van Bommel-Costa-Li-Poonen-Smith)

Let $m \in \mathbb{Z}_{\geq 0}$ and k be \mathbb{F}_3 or \mathbb{F}_5 . Then there is a squarefree ordinary A/k such that $\#A(k) = m$.

They use extremely clever constructions that allows them to construct characteristic polynomials h_A such that $h_A(1) = m$.

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group.

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Proof: Write

$$G \simeq \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_s\mathbb{Z}}.$$

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Proof: Write

$$G \simeq \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_s\mathbb{Z}}.$$

By H-K or vBCLPS, for each i there is an isogeny class with m_i points.

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Proof: Write

$$G \simeq \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_s\mathbb{Z}}.$$

By H-K or vBCLPS, for each i there is an isogeny class with m_i points. By Proposition, within each of the isogeny classes, there is a cyclic A_i .

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Proof: Write

$$G \simeq \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_s\mathbb{Z}}.$$

By H-K or vBCLPS, for each i there is an isogeny class with m_i points. By Proposition, within each of the isogeny classes, there is a cyclic A_i . Take $A = \prod_i A_i$. □

Group of points

Theorem (M.-Springer)

Let k be \mathbb{F}_2 , \mathbb{F}_3 or \mathbb{F}_5 . Let G be a finite abelian group. Then there exists an ordinary A/k with $A(k) = G$.

Proof: Write

$$G \simeq \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_s\mathbb{Z}}.$$

By H-K or vBCLPS, for each i there is an isogeny class with m_i points. By Proposition, within each of the isogeny classes, there is a cyclic A_i . Take $A = \prod_i A_i$. □

Corollary

If G is cyclic we can take A to be ordinary and squarefree.

Further results (building on vBCLPS)

Further results (building on vBCLPS)

- Over \mathbb{F}_4 : for every abelian $G \neq 0$ there exists an ordinary or almost ordinary A/\mathbb{F}_4 such that $A(\mathbb{F}_4) \simeq G$.

Further results (building on vBCLPS)

- Over \mathbb{F}_4 : for every abelian $G \neq 0$ there exists an ordinary or almost ordinary A/\mathbb{F}_4 such that $A(\mathbb{F}_4) \simeq G$.
- Over \mathbb{F}_7 : for every cyclic $G \neq 0$ with $\#G \notin \{2, 8, 14, 16, 17, 73\}$ there exists a squarefree ordinary A/\mathbb{F}_7 such that $A(\mathbb{F}_7) \simeq G$.

Further results (building on vBCLPS)

- Over \mathbb{F}_4 : for every abelian $G \neq 0$ there exists an ordinary or almost ordinary A/\mathbb{F}_4 such that $A(\mathbb{F}_4) \simeq G$.
- Over \mathbb{F}_7 : for every cyclic $G \neq 0$ with $\#G \notin \{2, 8, 14, 16, 17, 73\}$ there exists a squarefree ordinary A/\mathbb{F}_7 such that $A(\mathbb{F}_7) \simeq G$.
- vBCLPS: For an arbitrary q , every integer $m \geq q^{3\sqrt{q}\log q}$ arises as $m = \#A(\mathbb{F}_q)$ for some ordinary squarefree A/\mathbb{F}_q .

Further results (building on vBCLPS)

- Over \mathbb{F}_4 : for every abelian $G \neq 0$ there exists an ordinary or almost ordinary A/\mathbb{F}_4 such that $A(\mathbb{F}_4) \simeq G$.
- Over \mathbb{F}_7 : for every cyclic $G \neq 0$ with $\#G \notin \{2, 8, 14, 16, 17, 73\}$ there exists a squarefree ordinary A/\mathbb{F}_7 such that $A(\mathbb{F}_7) \simeq G$.
- vBCLPS: For an arbitrary q , every integer $m \geq q^{3\sqrt{q}\log q}$ arises as $m = \#A(\mathbb{F}_q)$ for some ordinary squarefree A/\mathbb{F}_q .

Theorem (M.-Springer)

Let m_1, \dots, m_r be integers satisfying $m_i \geq q^{3\sqrt{q}\log q}$. Put

$$G = \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_r\mathbb{Z}}.$$

Further results (building on vBCLPS)

- Over \mathbb{F}_4 : for every abelian $G \neq 0$ there exists an ordinary or almost ordinary A/\mathbb{F}_4 such that $A(\mathbb{F}_4) \simeq G$.
- Over \mathbb{F}_7 : for every cyclic $G \neq 0$ with $\#G \notin \{2, 8, 14, 16, 17, 73\}$ there exists a squarefree ordinary A/\mathbb{F}_7 such that $A(\mathbb{F}_7) \simeq G$.
- vBCLPS: For an arbitrary q , every integer $m \geq q^{3\sqrt{q}\log q}$ arises as $m = \#A(\mathbb{F}_q)$ for some ordinary squarefree A/\mathbb{F}_q .

Theorem (M.-Springer)

Let m_1, \dots, m_r be integers satisfying $m_i \geq q^{3\sqrt{q}\log q}$. Put

$$G = \frac{\mathbb{Z}}{m_1\mathbb{Z}} \times \dots \times \frac{\mathbb{Z}}{m_r\mathbb{Z}}.$$

Then there is an ordinary A/\mathbb{F}_q such that $G = A(\mathbb{F}_q)$.

Thank you!