



Australian
National
University

Student Number:

u							
---	--	--	--	--	--	--	--

Mathematical Sciences Institute

EXAMINATION: Semester 2 — Mid-Semester, 2016

MATH1013 — Advanced Mathematics and Applications 1

Book A — Calculus

Exam Duration: 120 minutes.

Reading Time: 15 minutes.

Materials Permitted In The Exam Venue:

- One A4 page with hand written notes on both sides.
(This A4 page is to cover both Algebra and Calculus.)
- Unmarked English-to-foreign-language dictionary (no approval from MSI required).
- **No electronic aids are permitted e.g. laptops, phones, calculators.**

Materials To Be Supplied To Students:

- Scribble Paper.

Instructions To Students:

- Answer the Calculus questions in Book A, and the Algebra questions in Book B, in the spaces provided.
- The Algebra and Calculus sections are worth a total of 20 points each, with the value of each question as shown. There are 2 questions in this section.
- It is recommended that you spend equal time on the Calculus and the Algebra papers.
- A good strategy is not to spend too much time on any question. Read them through first and attack them in the order that allows you to make the most progress.
- *You must justify your answers. Please be neat.*

Q1	Q2
10	10

Total / 20

Question 1**10 pts**

- (a) Using the definition of $\sinh x$ in terms of exponentials, find the indefinite integral

$$\int \sinh x \, dx . \quad 2 \text{ pts}$$

Write your solution here

(b) Evaluate $\int_1^{\sqrt{3}} \frac{1}{(1+x^2)(\tan^{-1} x)^2} dx .$ 4 pts

Write your solution here

Answer Make the substitution $u = \tan^{-1} x$. Then $du = \frac{1}{1+x^2} dx$, so

$$\begin{aligned} \int_1^{\sqrt{3}} \frac{1}{(1+x^2)(\tan^{-1} x)^2} dx &= \int_{\pi/4}^{\pi/3} \frac{du}{u^2} \\ &= \left[-u^{-1} \right]_{\pi/4}^{\pi/3} \\ &= -\frac{3}{\pi} + \frac{4}{\pi} \\ &= \frac{1}{\pi} . \end{aligned}$$

Grading Scheme:

- 1 point for trying a substitution
- 1 point for correctly performing the change of variables
- 1 point for computing the resulting integral
- 1 point for the right answer!

(c) Evaluate $\int_1^2 (\ln x)^2 dx$.

4 pts

Write your solution here

Answer

Let $U_1 = (\ln x)^2$ and $dV_1 = dx$. Then $dU_1 = \frac{2}{x} \ln x dx$ and $V_1 = x$, so integration by parts gives

$$\int_1^2 (\ln x)^2 dx = \left[x(\ln x)^2 \right]_1^2 - \int_1^2 x \frac{2}{x} \ln x dx = 2(\ln 2)^2 - 2 \int_1^2 \ln x dx.$$

Now let $U_2 = \ln x$ and $dV_2 = dx$. Then $dU_2 = \frac{1}{x} dx$ and $V_2 = x$, so integration by parts gives

$$\begin{aligned} \int_1^2 \ln x dx &= \left[x \ln x \right]_1^2 - \int_1^2 x \frac{1}{x} dx \\ &= 2 \ln 2 - \int_1^2 dx \\ &= 2 \ln 2 - 1. \end{aligned}$$

Hence

$$\int_1^2 (\ln x)^2 dx = 2(\ln 2)^2 - 4 \ln 2 + 2.$$

Question 2**10 pts**

Suppose $\alpha > 1$, $K \geq 0$, and that $f : \mathbb{R} \rightarrow \mathbb{R}$ has the property that, for all $x, y \in \mathbb{R}$,

$$|f(x) - f(y)| \leq K|x - y|^\alpha.$$

Prove that f is constant.

Hints: for this question you may use the facts that

- for all $r > 0$, the mapping $t \mapsto t^r$ is continuous on $[0, \infty)$, and
- if $g : \mathbb{R} \setminus \{c\} \rightarrow \mathbb{R}$, then $\lim_{x \rightarrow c} g(x) = 0$ if and only if $\lim_{x \rightarrow c} |g(x)| = 0$.

Given $c \in \mathbb{R}$, consider $h : \mathbb{R} \setminus \{c\} \rightarrow \mathbb{R}$ defined by $h(x) = \frac{f(x) - f(c)}{x - c}$.

Finally, please note that $\alpha > 1$, NOT $\alpha > 0$.

Write your solution here

Extra space for previous question