

Quartic Analysis

Sina Tootoonian, Claude

August 4, 2026

We're given

- $\mathbf{X}$: an N cells $\times$ P odours matrix of observed input responses,
- $\mathbf{C}$: a P cells $\times$ P cells observed output covariance matrix,
- The regularization parameter $\lambda = \lambda_0 \frac{P^2}{N}$, where λ_0 is the value we sweep in the paper.
- The mean subtraction operation, $\mathbf{J} = \mathbf{I} - N^{-1}\mathbf{1}\mathbf{1}^T$.
- The prediction error $\mathbf{E}(\mathbf{z}) = \mathbf{X}^T[\mathbf{z}]\mathbf{J}[\mathbf{z}]\mathbf{X} - \mathbf{C}$

Our loss function is

$$L(\mathbf{z}) = \frac{1}{2}\|\mathbf{E}(\mathbf{z})\|_F^2 + \frac{\lambda}{2}\|\mathbf{z} - \mathbf{1}\|_2^2, \quad (1) \quad \{\text{eq:loss}\}$$

where $[\cdot]$ converts its argument to a diagonal matrix.

We want to view this from the perspective of a single unit z_i . To do that, we rewrite $\mathbf{E}(\mathbf{z})$ to isolate that unit. Examining its components,

$$\begin{aligned} \mathbf{X}^T[\mathbf{z}]\mathbf{J}[\mathbf{z}]\mathbf{X} &= \mathbf{X}^T[\mathbf{z}^2]\mathbf{X} - N^{-1}\mathbf{X}^T\mathbf{z}\mathbf{z}^T\mathbf{X} \\ \mathbf{X}^T[\mathbf{z}^2]\mathbf{X} &= \sum_{j=1}^N \mathbf{x}_j \mathbf{x}_j^T z_j^2 = \mathbf{x}_i \mathbf{x}_i^T z_i^2 + \sum_{j \neq i} \mathbf{x}_j \mathbf{x}_j^T z_j^2 \\ \mathbf{m} \triangleq \mathbf{X}^T \mathbf{z} &= \sum_j \mathbf{x}_j z_j = \mathbf{x}_i z_i + \sum_{j \neq i} \mathbf{x}_j z_j = \mathbf{x}_i z_i + \mathbf{m}_i \\ \mathbf{E}(\mathbf{z}) &= \mathbf{x}_i \mathbf{x}_i^T z_i^2 + \sum_{j \neq i} \mathbf{x}_j \mathbf{x}_j^T z_j^2 - N^{-1}(\mathbf{x}_i z_i + \mathbf{m}_i)(\mathbf{x}_i z_i + \mathbf{m}_i)^T - \mathbf{C} \\ &= \mathbf{x}_i \mathbf{x}_i^T z_i^2 - N^{-1}(\mathbf{x}_i \mathbf{m}_i^T z_i + \mathbf{m}_i \mathbf{x}_i^T z_i + \mathbf{x}_i \mathbf{x}_i^T z_i^2) + \sum_{j \neq i} \mathbf{x}_j \mathbf{x}_j^T z_j^2 - N^{-1}\mathbf{m}_i \mathbf{m}_i^T - \mathbf{C} \\ &= \alpha \mathbf{x}_i \mathbf{x}_i^T z_i^2 - N^{-1}(\mathbf{x}_i \mathbf{m}_i^T + \mathbf{m}_i \mathbf{x}_i^T) z_i + \mathbf{E}_i. \end{aligned}$$

Computing the goodness-of-fit term of the loss,

$$\begin{aligned} \text{tr}(\mathbf{E}(\mathbf{z})^T \mathbf{E}(\mathbf{z})) &= \alpha^2 \|\mathbf{x}_i\|_2^4 z_i^4 - 4\alpha N^{-1} \|\mathbf{x}_i\|_2^2 \mathbf{m}_i^T \mathbf{x}_i z_i^3 + 2\alpha \mathbf{x}_i^T \mathbf{E}_i \mathbf{x}_i z_i^2 \\ &\quad + 2N^{-2} z_i^2 (\|\mathbf{x}_i\|_2^2 \|\mathbf{m}_i\|_2^2 + (\mathbf{x}_i^T \mathbf{m}_i)^2) - 4N^{-1} z_i \mathbf{x}_i^T \mathbf{E}_i \mathbf{m}_i + \text{tr}(\mathbf{E}_i^T \mathbf{E}_i). \end{aligned}$$

Our loss function in terms of z_i is then

$$\begin{aligned}
L(z_i) = & \frac{1}{2}\alpha^2\|\mathbf{x}_i\|_2^4 z_i^4 \\
& - \frac{2\alpha}{N}\|\mathbf{x}_i\|_2^2 \mathbf{m}_i^T \mathbf{x}_i z_i^3 \\
& + \left(\alpha \mathbf{x}_i^T \mathbf{E}_i \mathbf{x}_i + \frac{\|\mathbf{x}_i\|_2^2 \|\mathbf{m}_i\|_2^2 + (\mathbf{x}_i^T \mathbf{m}_i)^2}{N^2} + \frac{\lambda}{2} \right) z_i^2 \\
& - \frac{2\mathbf{x}_i^T \mathbf{E}_i \mathbf{m}_i + N\lambda}{N} z_i \\
& + \frac{\text{tr}(\mathbf{E}_i^T \mathbf{E}_i) + \lambda(1 + \sum_{i \neq j} (z_j - 1)^2)}{2}.
\end{aligned}$$

The trailing constant doesn't matter for the minimization, so we'll drop it. We'll express the remaining coefficients in terms of a few geometric quantities:

- The raw power of the population mean $m \triangleq \mathbf{m}_i^T \mathbf{m}_i$;
- The raw channel power, $p_i \triangleq \mathbf{x}_i^T \mathbf{x}_i$;
- The residual channel power, $q_i \triangleq \mathbf{x}_i^T \mathbf{E}_i \mathbf{x}_i$;
- The raw alignment with the population mean, $a_i \triangleq \mathbf{x}_i^T \mathbf{m}_i$;
- The residual alignment with the population mean, $b_i \triangleq \mathbf{x}_i^T \mathbf{E}_i \mathbf{m}_i$;
- The curvature at the origin, $\kappa_i \triangleq \alpha q_i + \frac{p_i m + a_i^2}{N^2}$.

In these terms, and dropping the constant,

$$L(z_i) = \frac{1}{2}\alpha^2 p_i^2 z_i^4 - \frac{2\alpha}{N} p_i a_i z_i^3 + \left(\kappa_i + \frac{\lambda}{2} \right) z_i^2 - \left(\frac{2b_i}{N} + \lambda \right) z_i \quad (2) \quad \{\mathbf{eq:Lzi}\}$$

$$\triangleq A z_i^4 + B z_i^3 + C z_i^2 + D z_i. \quad (3)$$

To better understand this quartic, we will change coordinates. First, we'll rescale by A ,

$$\begin{aligned}
L(z) & \propto z^4 + \frac{B}{A} z^3 + \frac{C}{A} z^2 + \frac{D}{A} z \\
& \triangleq z^4 + b z^3 + c z^2 + d z.
\end{aligned}$$

Next, we'll shift to a coordinate $s = z - \bar{z}$ in which the cubic disappears. Expressing the loss as a function $\tilde{L}$ of s , we note that for a monic quartic with no cubic, the third derivative should be $24s$.

$$\begin{aligned}
\tilde{L}(s) &= (s + \bar{z})^4 + b(s + \bar{z})^3 + c(s + \bar{z})^2 + d(s + \bar{z}). \\
\tilde{L}'''(s) &= 24(s + \bar{z}) + 6b = 24s \implies \bar{z} = -\frac{b}{4}.
\end{aligned}$$

So, in the shifted and scaled coordinates, the loss function is

$$\tilde{L}(s) = \left(s - \frac{b}{4} \right)^4 + c \left(s - \frac{b}{4} \right)^2 + d \left(s - \frac{b}{4} \right)$$

We can expand out the first equation to get the coefficients. But an easier way is to Taylor expand our loss function $L(z)$ to get a polynomial in s :

$$\tilde{L}(s) = L(s + \bar{z}) = L(\bar{z}) + L'(\bar{z})s + \frac{1}{2}L''(\bar{z})s^2 + \frac{1}{24}L'''(\bar{z})s^4.$$

Then, $L''''(\bar{z}) = 24$. The other derivatives are

$$\begin{aligned} L'(\bar{z}) &= 4\bar{z}^3 + 3b\bar{z}^2 + 2c\bar{z} + d = -\frac{b^3}{16} + 3\frac{b^3}{16} - \frac{bc}{2} + d = \frac{b^3}{8} - \frac{bc}{2} + d \\ L''(\bar{z}) &= 12\bar{z}^2 + 6b\bar{z} + 2c = 12\frac{b^2}{16} - 6\frac{b^2}{4} + 2c = -\frac{3}{4}b^2 + 2c. \end{aligned}$$

So we arrive at

$$\begin{aligned} \tilde{L}(s) &= s^4 + \left(c - \frac{3}{8}b^2\right)s^2 + \left(d - \frac{bc}{2} + \frac{b^3}{8}\right)s + \text{const} \\ &\triangleq s^4 + \tilde{c}s^2 + \tilde{d}s + \text{const}. \end{aligned} \tag{4}$$

We can see that $\tilde{L}$ combines a convex quartic with a parabola. If the parabola is also convex, so $\tilde{c} \geq 0$, then $\tilde{L}$ will also be convex, and have only one root. If the parabola is concave (pointing down), so $\tilde{c} < 0$, then $\tilde{L}$ may have one or two roots, depending on the tilt imposed by the linear term.

To determine the location of the root(s), we set the derivative to 0:

$$0 = \tilde{L}'(s) = 4s^3 + 2\tilde{c}s + \tilde{d} = s^3 + \frac{\tilde{c}}{2}s + \frac{\tilde{d}}{4} \triangleq s^3 + gs + h.$$

To determine the locations of the roots, we use the triple angle formula:

$$4\cos^3(\theta) - 3\cos(\theta) - \cos(3\theta) = 0.$$

Let $s = K\cos(\theta)$. Then our derivative equation is

$$K^3\cos(\theta)^3 + Kg\cos(\theta) + h = 0.$$

Dividing out by $K^3/4$, we get

$$4\cos(\theta)^3 + \frac{4g}{K^2}\cos(\theta) + \frac{4h}{K^3} = 0.$$

Now if $g \leq 0$, we can set K such that

$$\frac{4g}{K^2} = -3 \implies K = 2\sqrt{|g|/3}.$$

Then we get

$$\cos(3\theta) = -\frac{4h}{K^3} = -\frac{4h}{K^2} \frac{1}{K} = -\frac{3h}{2|g|} \sqrt{\frac{3}{|g|}} = \frac{3h}{2g} \sqrt{\frac{3}{-g}} \triangleq u.$$

This equation will have solution if $|u| \leq 1$, equivalently

$$1 \geq u^2 = \frac{27h^2}{-4g^3} \iff 27h^2 \leq 4(-g)^3,$$

with equality determining the *cusp*. The solutions will satisfy

$$3\theta = \arccos(u) + 2\pi k \implies \theta = \frac{1}{3} \arccos(u) + \frac{2\pi k}{3}, \quad k \in \{0, 1, 2\},$$

corresponding to the two wells and one ridge.

Which k corresponds to the ridge, and which to the valleys? When $u = 0$, there is no tilt in the quartic and the wells will be symmetric around 0. In this case,

$$\theta = \frac{1}{3} \arccos(0) + \frac{2\pi k}{3} = \frac{\pi}{6} + \frac{2\pi k}{3} = \left\{ \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2} \right\}.$$

Then

$$\cos(\theta) = \{\sqrt{3}/2, -\sqrt{3}/2, 0\}.$$

So, assuming this ordering holds at non-zero u , $k = 0$ gives the solution when the tilt is negative, $k = 1$ gives the solution for positive tilt, and $k = 2$ gives the ridge.

Now we're interested in how large the cosine term can get. As the tilt becomes more negative, the minimum at $\sqrt{3}/2$ will move to the right. How far forward will it go? Negative tilt corresponds to negative h and hence positive u since g is negative in this regime. Hence the maximally negative tilt, occurring when the ridge is about to disappear, will correspond to the $u = 1$ case. At that point

$$\theta = \frac{1}{3} \arccos(1) + \frac{2\pi k}{3} = \left\{0, \frac{2\pi}{3}, \frac{4\pi}{3}\right\}$$

So then

$$\cos(\theta) = \{1, -1/2, -1/2\},$$

corresponding to a global minimum at 1 and an inflection point at $-1/2$.

So, when $|u| \leq 1$, we can represent the solution as

$$s^* = \frac{\sqrt{3}}{2} K \operatorname{sign}(u) f(|u|), \quad f \in \left[1, \frac{2}{\sqrt{3}}\right].$$

Plugging in our expression for K simplifies this to

$$s^* = \sqrt{|g|} \operatorname{sign}(u) f(|u|).$$

Now let's consider the case when $g > 0$. In that case, the relevant triple-angle formula is for $\sinh$,

$$4 \sinh^3(\theta) + 3 \sinh(\theta) - \sinh(3\theta) = 0.$$

If we set $s = K \sinh(\theta)$, our depressed cubic becomes

$$K^3 \sinh(\theta)^3 + Kg \sinh(\theta) + h = 0.$$

Dividing by $K^3/4$ this gives

$$4 \sinh(\theta)^3 + \frac{4g}{K^2} \sinh(\theta) + \frac{4h}{K^3} = 0.$$

Setting

$$\frac{4g}{K^2} = 3 \implies K = 2\sqrt{g/3}$$

gives

$$\sinh(3\theta) = -\frac{4h}{K^3} = -\frac{4h}{K^2} \frac{1}{K} = -3 \frac{4h}{4g} \frac{1}{K} = -\frac{3h}{2g} \sqrt{\frac{3}{g}} \triangleq v.$$

We then get $\theta = \frac{1}{3} \operatorname{arcsinh}(v)$, so

$$s^* = K \sinh(\theta) = 2\sqrt{\frac{g}{3}} \sinh\left(\frac{1}{3} \operatorname{arcsinh}(v)\right).$$

When v is small,

$$s^* \approx \frac{1}{3} K v = -\frac{h}{g}.$$

When v is large,

$$s^* \approx -h^{1/3}.$$

Complexifying

Rather than treating each of these cases separately, we can view them all in a unified manner by consider the roots of our depressed cubic in the complex plane.

In that setting, the triple angle formula is

$$w^3 + w^{-3} = (w + w^{-1})^3 - 3(w + w^{-1}).$$

We now parameterize

$$s = \frac{K}{2}(w + w^{-1}).$$

Our cubic equation then becomes

$$\frac{K^3}{8}(w + w^{-1})^3 + \frac{Kg}{2}(w + w^{-1}) + h = 0.$$

Normalizing,

$$(w + w^{-1})^3 + \frac{4g}{K^2}(w + w^{-1}) + \frac{8h}{K^3} = 0.$$

Setting K so that

$$\frac{4g}{K^2} = -3 \implies K = \sqrt{\frac{-4g}{3}} = 2\sqrt{\frac{-g}{3}},$$

Making that substitution gives

$$w^3 + w^{-3} = -\frac{8h}{K^3} \implies w^3 + w^{-3} + \frac{8h}{K^3} = 0.$$

Then letting $W = w^3$, we have

$$W + \frac{1}{W} + \frac{8h}{K^3} = 0.$$

Multiplying by W , we get

$$W^2 + 2\frac{4h}{K^3}W + 1 = 0.$$

Since the constant coefficient is 1, we have that that two roots of this equation are inverses of each other.

Defining

$$u \triangleq -\frac{4h}{K^3} = -\frac{4h}{K^2} \frac{1}{K} = \frac{3h}{2g} \sqrt{\frac{3}{-g}}, \quad u^2 = -\frac{27h^2}{4g^3},$$

we get

$$W = \frac{2u \pm \sqrt{4u^2 - 4}}{2} = u \pm \sqrt{u^2 - 1}.$$

Now each value of W will gives us three values of w . There are two values of W , so this will yield six total values of w . However, these can be put in three pairs, where each element of the pair is the inverse of the other. And s is invariant to inverses, so $s(w) = s(w^{-1})$, and we get three values of s as the roots of the cubic, as expected.

Let's consider the various cases.

First, let's assume $g < 0$, so K is real. In that case, $u^2 \geq 0$.

If $|u| \leq 1$, then the square-root is pure imaginary, so $|W| = 1$ and W is on the unit circle. So

$$W = e^{j(3\theta+2\pi k)} \implies w = e^{j\theta_k}, \quad \theta_k \triangleq \theta + \frac{2\pi k}{3}, \quad k \in \{0, 1, 2\}.$$

This in turn means three real roots

$$s = K \cos\left(\theta + \frac{2\pi k}{3}\right), \quad k \in \{0, 1, 2\}, \quad \theta = \frac{1}{3} \arccos(u).$$

If $|u| > 1$, then the square-root in the definition of W is real, so $\cos$ becomes $\cosh$, and u has a unique real cube root,

$$s = K \operatorname{sign}(u) \cosh(\theta), \quad \theta = \frac{1}{3} \operatorname{arccosh}|u|.$$

Next, let's assume $g > 0$, so K is imaginary. Then $u^2 < 0$, so u is imaginary, and W is pure imaginary.