

Linear Algebra: Final Extra Credit

Due on December 13, 2019

Professor MacArthur

Carson Storm

The inner product is essential for many of the concepts used in Chapter 6, but so far we have only defined the inner product for real number spaces, \mathbb{R}^n . In order to expand the notion of an inner product to other vector spaces we define a set of axioms that a space's inner product must satisfy, so that we can define an inner product for other vector spaces. A inner product is an operation that takes two vectors in the considered vectors space as operands and associated with them a real number. This inner product can be used to define what a vector's length is, distances in the space as well as orthogonality. For example, be defining a inner product for \mathbb{P}^n , we define what it means for two polynomials to be orthogonal. After defining the inner product we can utilize concepts such as the Gram-Schmidt process, or solving Least-Squares problems in spaces that are not \mathbb{R}^n .

Carson Storm

Section 6.7 # 1, 3, 4, 7, 10, 13, 18, 21, 25

$$2) a) \|\vec{x}\| = \sqrt{\langle \vec{x}, \vec{x} \rangle} = \sqrt{4x_1^2 + 5x_2^2} \quad \|\vec{y}\| = \sqrt{\langle \vec{y}, \vec{y} \rangle} = \sqrt{4y_1^2 + 5y_2^2} = \sqrt{100 + 5} = \sqrt{105}$$

$$= \sqrt{9} = 3 \quad \|\vec{y}\| = \sqrt{105}$$

$$\|\vec{x}\| = 3$$

$$|\langle \vec{x}, \vec{y} \rangle|^2 = |4x_1y_1 + 5x_2y_2|^2 = |20 - 5|^2 = |15|^2 = 225$$

$$|\langle \vec{x}, \vec{y} \rangle|^2 = 225$$

$$b) \langle \vec{z}, \vec{y} \rangle = 0 = 4z_1y_1 + 5z_2y_2 = 0 \quad z_1 = \frac{1}{4}z_2$$

$$= 20z_1 - 5z_2$$

$$0 = 4z_1 - z_2$$

All $\vec{z} \in \text{Span}\left\{\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right\}$ are orthogonal to \vec{y}

$$3) P(t) = 4 + t, \quad q(t) = 5 - 4t^2$$

$$\langle P, q \rangle = P(t_0)q(t_0) + P(t_1)q(t_1) + \dots + P(t_n)q(t_n)$$

$$\{t_n\} = \{-1, 0, 1\}$$

$$\langle P, q \rangle = P(-1)q(-1) + P(0)q(0) + P(1)q(1)$$

$$= 3(1) + 4(5) + 5(1)$$

$$= 3 + 20 + 5 = 28$$

$$\langle P, q \rangle = 28$$

$$4) P(t) = 3t - t^2, \quad q(t) = 3 + 2t^2$$

$$\langle P, q \rangle = P(-1)q(-1) + P(0)q(0) + P(1)q(1)$$

$$= -4(5) + 0(3) + 2(5)$$

$$= -20 + 10 = -10$$

$$\langle P, q \rangle = -10$$

$$7) \text{Proj}_P q = \frac{\langle P, q \rangle}{\|P\|^2} P = 28 \cdot \frac{4+t}{9+16+25} = 28 \cdot \frac{4+t}{50} = \frac{28(4+t)}{50} = \frac{14(4+t)}{25} = \frac{56+14t}{25}$$

$$\text{Proj}_P q(t) = \frac{56+14t}{25}$$

$$10) \{P_0, P_1, q\} = \left\{1, t, \frac{t^2-5}{4}\right\} \quad \text{Proj}_V P = \frac{\langle P, P_0 \rangle}{\langle P_0, P_0 \rangle} P_0 + \frac{\langle P, P_1 \rangle}{\langle P_1, P_1 \rangle} P_1 + \frac{\langle P, q \rangle}{\langle q, q \rangle} q$$

$$= \frac{0}{4} \cdot 1 + \frac{164}{20} \cdot t + \frac{0}{4} \cdot \frac{t^2-5}{4}$$

$$= \frac{41}{5}t$$

The best approximation of $P(t) = t^3$ in $\text{Span}\{P_0, P_1, q\}$ is

$$\text{Proj}_V P(t) = \frac{41}{5}t$$

$$13) \langle \vec{u}, \vec{v} \rangle = (A\vec{u}) \cdot (A\vec{v})$$

$$1. \langle \vec{u}, \vec{v} \rangle = (A\vec{u}) \cdot (A\vec{v}) = (A\vec{v}) \cdot (A\vec{u})$$

$$\langle \vec{u}, \vec{v} \rangle = \langle \vec{v}, \vec{u} \rangle$$

$$2. \langle \vec{u} + \vec{v}, \vec{w} \rangle = (A(\vec{u} + \vec{v})) \cdot (A\vec{w})$$

$$= (A\vec{u} + A\vec{v}) \cdot (A\vec{w})$$

$$= (A\vec{u}) \cdot (A\vec{w}) + (A\vec{v}) \cdot (A\vec{w})$$

$$= \langle \vec{u}, \vec{w} \rangle + \langle \vec{v}, \vec{w} \rangle$$

Therefore, $\langle u, v \rangle = (A\vec{u}) \cdot (A\vec{v})$ defines an inner product space because it satisfies all of its axioms.

$$3) \langle c\vec{u}, \vec{v} \rangle = (A(c\vec{u})) \cdot (A\vec{v})$$

$$= (cA\vec{u}) \cdot A\vec{v}$$

$$= c(A\vec{u}) \cdot (A\vec{v})$$

$$= c \langle \vec{u}, \vec{v} \rangle$$

$$4) \text{ let } (A\vec{u}) = [a_1, a_2, \dots, a_n]$$

$$\langle \vec{u}, \vec{u} \rangle = (A\vec{u})^T (A\vec{u}) = a_1^2 + a_2^2 + \dots + a_n^2$$

$$\langle \vec{u}, \vec{u} \rangle \geq 0 \text{ since } a_n^2 \geq 0 \forall n$$

$$\langle \vec{u}, \vec{u} \rangle = 0 \Rightarrow a_n = 0 \forall n \Rightarrow \vec{u} = 0 \text{ because } A \text{ is invertible}$$

$$18) \text{ Show that } \|\vec{u} + \vec{v}\|^2 + \|\vec{u} - \vec{v}\|^2 = 2\|\vec{u}\|^2 + 2\|\vec{v}\|^2$$

$$\|\vec{u} + \vec{v}\|^2 + \|\vec{u} - \vec{v}\|^2 = \langle \vec{u} + \vec{v}, \vec{u} + \vec{v} \rangle + \langle \vec{u} - \vec{v}, \vec{u} - \vec{v} \rangle$$

$$= \langle \vec{u}, \vec{u} + \vec{v} \rangle + \langle \vec{v}, \vec{u} + \vec{v} \rangle + \langle \vec{u}, \vec{u} - \vec{v} \rangle - \langle \vec{v}, \vec{u} - \vec{v} \rangle \quad (\text{Axiom 2})$$

$$= \langle \vec{u} + \vec{v}, \vec{u} \rangle + \langle \vec{u} + \vec{v}, \vec{v} \rangle + \langle \vec{u} - \vec{v}, \vec{u} \rangle - \langle \vec{u} - \vec{v}, \vec{v} \rangle \quad (\text{Axiom 1})$$

$$= \langle \vec{u}, \vec{u} \rangle + \langle \vec{v}, \vec{u} \rangle + \langle \vec{u}, \vec{v} \rangle + \langle \vec{v}, \vec{v} \rangle + \langle \vec{u}, \vec{u} \rangle - \langle \vec{u}, \vec{v} \rangle - \langle \vec{u}, \vec{v} \rangle + \langle \vec{v}, \vec{v} \rangle \quad (\text{Axiom 2})$$

$$= 2\langle \vec{u}, \vec{u} \rangle + 2\langle \vec{v}, \vec{v} \rangle$$

$$= 2\|\vec{u}\|^2 + 2\|\vec{v}\|^2$$

$$21) \text{ In } C[0,1] \langle f, g \rangle = \int_0^1 f(t)g(t)dt, \quad f(t) = 1 - 3t^2, \quad g(t) = t - t^3$$

$$\langle f, g \rangle = \int_0^1 (1 - 3t^2)(t - t^3)dt = \int_0^1 (3t^5 - 4t^3 + t)dt = \left[\frac{1}{2}t^6 - t^4 + \frac{1}{2}t^2 \right]_0^1 = \frac{1}{2} - 1 + \frac{1}{2} - 0 + 0 - 0$$

$$\langle f, g \rangle = 0$$

$$25) \{x_1, x_2, x_3\} = \{1, t, t^2\}, \quad \langle f, g \rangle = \int_{-1}^1 f(t)g(t)dt$$

$$v_1 = x_1, \quad v_2 = x_2, \quad v_3 = x_3 - \frac{\langle x_3, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 - \frac{\langle x_3, v_2 \rangle}{\langle v_2, v_2 \rangle} v_2$$

$$= t^2 - \frac{(2/3)}{2}(1) - \frac{(0)}{(2/3)}(t)$$

$$= t^2 - \frac{1}{3}$$

A orthogonal basis for $\text{span}\{1, t, t^2\}$ in $C[-1, 1]$ is $\{1, t, t^2 - \frac{1}{3}\}$