

Should we build a macro-scale theory for coevolution? Nope.

Timothée Poisot Daniel B. Stouffer

Oct. 31, 2014

1 Coevolutionary dynamics acting on both species and their interactions are
2 a key driving force behind the structure of ecological communities. The
3 Geographic Mosaic Theory of Coevolution (GMTC) provides a spatial per-
4 spective to these dynamics by proposing the existence of feedbacks between
5 local and regional scales. It remains unclear, however, how the structure
6 of communities at larger spatial scales either influences or is influenced
7 by local coevolutionary processes. Despite this, ecological networks are
8 known to have an ecological structure, which suggest that coevolution may
9 play a role in their evolutionary dynamics. Here we show that the coevo-
10 lutionary significance of individual interactions is maintained when mov-
11 ing between the local and regional scale. Importantly, this occurs despite
12 the fact that community variation at the local scale tends to weaken or re-
13 move community-wide coevolutionary signal. This apparent mismatch be-
14 tween our interaction-level and community-level results strongly suggests
15 that interactions provide the correct scale to study coevolution at small spa-
16 tial scales while communities are the relevant scale only at larger spatial

1 extents. We provide a new perspective on the interplay between coevolu-
2 tionary theory and community ecology, by establishing the organisational
3 scales at which the different theories have relevance. Although it has been
4 tempting so far to understand how coevolution relates to network structure,
5 our results suggest that the way forward is to understand how network
6 structure may affect coevolution over space instead.

7 Ecological interactions introduce selective pressures on the species involved.
8 At large organisational scales, this results in taxonomic boundaries delineating
9 groups of interacting species ¹, invariant structures in some ecological com-
10 munities ², and conservatism of both the distribution of community modules
11 ⁴ and the role of species occupy within them ⁵. Although the evolutionary dy-
12 namics for a pair of interacting species has been well described ^{???}, attempts
13 to understand how these mechanisms cascade up to generate species diversity
14 observed in large ecological networks have been unconvulsive ^{6,7}, despite the
15 well known effect of antagonistic coevolution on genomic diversification ^{8,9}.
16 The scales at which these diversities happen are hard to reconcile: coevolution
17 is expressed within patches connected by gene-flow ¹⁰⁻¹², whereas the species
18 diversity of complex networks is typically observed at spatial scales matching
19 the species distribution ^{13,14}. Because these scales differ by orders of magni-
20 tude, one must question the relevance of previous calls to scale the theory on
21 coevolution up to multi-species systems covering large spatial extents ^{???}.

22 Network-based approaches ^{15,16}, on the other hand, were designed to study
23 and describe species-rich systems. Previous empirical findings revealed the
24 impact of evolutionary dynamics on overall network structure in food webs

1 ^{1,17}, pollination networks ^{18,19}, and host-parasite networks ^{20,21}. Both micro
2 ^{3,22} and macro ^{23,24} evolutionary dynamics have been measured or modelled
3 in species-rich communities, but there is no understanding of how, or even of
4 whether, local/micro-evolutionary and regional/macro-evolutionary feedback
5 into one another. Another layer of complexity is that ecological networks
6 are known to vary in their structure over time and space ²⁵: the same two
7 species will not interact in a consistent way locally, either because of local
8 environmental contingencies, by chance, or because their phenologies do not
9 positively covary in space ²⁶. As a consequence, *locally*, the evolutionary signal
10 on network structure is expected to be buried under much ecological noise,
11 and the effect of coevolution can only be inferred *regionally*.

12 In multi-species systems that typically span a large taxonomic range, coevo-
13 lution is often measured as the matching between the phylogenies of two
14 sets of interacting organisms ^{20,27}. This build on the century-old ideas that
15 extant species interact in a way similar to the way their ancestors did ²⁸. “Co-
16 evolved” systems should (i) have approximately similar phylogenetic trees
17 and (ii) species at matching positions in either trees should interact. It is not
18 clear, however, how this idea relates to dynamics occurring at smaller scales ²⁹:
19 many ecological and evolutionary processes that occur locally, or over small
20 spatial scales, can disturb this expected structure. Notably, it has been shown
21 that species interactions are not consistent through space ^{25,30}. Local loss of
22 both interactions and species from the regional pool is most likely to result in
23 observed communities that do not appear to have been shaped by coevolution.

24 We use data on ectoparasites of rodents from Western to Eastern Europe ³¹

1 to test the following four hypotheses. First, local species assemblages do not
2 show evidence of coevolution even though the system as a whole does. Sec-
3 ond, interaction-level coevolutionary signal is conserved. Third, interaction-
4 level coevolutionary signal does not predict the spatial consistency of inter-
5 actions. We do so by coupling two novel methods: the *PACO* algorithm for
6 detection of phylogenetic congruence³², and a general framework for the vari-
7 ation of ecological networks³⁰.

8 Local observations on the 51 localities (*Supp. Mat. 1*) are aggregated into a re-
9 gional metanetwork³⁰. The two phylogenetic trees have been rendered ultra-
10 metric. We use *PACO*³² to measure the congruence between trees knowing the
11 matrix of interaction. *PACO* yields a network-level significance value for the
12 likelihood that hosts and parasites have coevolved. For each local network, we
13 measure the strength of coevolution using (i) only local observations and (ii)
14 all possible interactions between local species (as known from the regional ag-
15 gregation of all local networks). This allows us to separate the effect of species
16 sorting (regional interactions) and interaction sorting (local interactions). At
17 the regional scale, coevolutionary signal is extremely strong ($p \leq 10^{-4}$), as
18 established by previous analysis of this system³³. Most local networks, on
19 the other hand, show very little evidence of phylogenetic congruence. Out
20 of 51 local networks, 35 show no signal of coevolution, 11 show coevolution
21 when using the regional interactions, and 12 show coevolution using the local
22 interactions (see *Supp. Mat. 1* for network-level significance values). **Figure?**

23 This suggests that macro-evolutionary processes (such as co-diversification)
24 have consequences at the macro-ecological level³⁴, but may not be detected




Figure 1: figure1

1 at finer spatial scales due to a stronger effect of ecological processes locally.
2 *PACO* permits the analysis of *how strongly* each interaction contribute to co-
3 evolution, in a way that is as independent as possible from other interactions.
4 As interactions vary only insofar that there are some locations in which they
5 *do not* happen, we expect that the overall contribution of interactions will be
6 the same in the local and regional networks. For the 5 networks that show
7 evidences of coevolution accounting both for species and interactions sorting,
8 we measured the contribution of each interaction locally, and compared it to
9 its contribution to the regional network. Results are presented in **FIG**.

10 **Point 2 – Interactions respond locally to micro-processes**

- 11 • Review of mechanisms involved in interactions locally: neutrality and
12 trait-based mechanisms
- 13 • Key refs: Oikos paper, Canard, Olesen, Combes
- 14 • These mechanism can only *filter* from a pool of regional interactions
- 15 • Key refs: Biol Lett paper, Olito & Fox, Lounnas, ...
- 16 • Key points: i) even though they will change the network-level signal,
17 contribution of interactions should not vary locally vs. regionally ii) this
18 is the expectation because (a) past knowledge about motifs/evolution
19 and (b) interactions are the unit in which selection takes place, not net-
20 works

- 1 • Illustration: distribution of the significance in each interaction locally
- 2 (measured vs. metaweb) vs. in the regional network.




Figure 2: figure2

1 **Point 3 – The spatial replication of networks is im-** 2 **perfect**

- 3 • Ecological processes = sorting of interactions + species
- 4 • Key refs: Ecol Lett -div
- 5 • Key points: observation of interactions result from filtering through two
6 processes, and not clear they are related to past coevolution = we expect
7 a loss of signal locally
- 8 • Illustration: jackknife score vs. observed/possible : there is no relation-
9 ship between how important an interaction is (coevo) and how frequent
10 it will be. suggests that ecological >> evolutionary drivers of local net-
11 work structure
- 12 • This is caused by local ecological processes, but also influences local
13 evolutionary dynamics
- 14 • Key refs: Gomulkiewicz, Hochberg, Benkman, Siepilski
- 15 • Key points: i) Opportunity to integrate more network theory to under-
16 stand between-sites differences ii) Even though coevo theory recognized
17 variation in effects, very little attention to variation in presence/absence
18 of int and its implications




Figure 3: figure3

Conclusions

- Need to bridge ecological networks with co-evolution (understand the impact of ecology vs. evolution in the local observations)
- Main advance: coevolutionary dynamics are important at the continental/regional scale, much less at the local scale
- Focus on datasets that are spatially replicated, with as good as possible phylogenies. Ref Buckley, Harkel.
- Currently local variability is not accounted for by statistical methods, need to do a better job. See Desdevises chapter on ParaFit 2.0.

1. Eklof, A., Helmus, M. R., Moore, M., Allesina, S. & Eklöf, A. *et al.* Relevance of evolutionary history for food web structure. *Proc. R. Soc. B Biol. Sci.* **279**, 1588–1596 (2011).

2. Nuismer, S. L., Jordano, P. & Bascompte, J. *et al.* Coevolution and the architecture of mutualistic networks. *Evolution* **67**, 338–354 (2013).

3. Jordano, P., Bascompte, J. & Olesen, J. M. *et al.* Invariant properties in coevolutionary networks of plant-animal interactions. *Ecol. Lett.* **6**, 69–81 (2003).

4. Stouffer, D. B., Camacho, J., Guimera, R., Ng, C. A. & Amaral, L. A. N. *et al.* Quantitative patterns in the structure of model and empirical food webs. *Ecology* **86**, 1301–1311 (2005).

5. Stouffer, D. B., Sales-Pardo, M., Sirer, M. I. & Bascompte, J. *et al.* Evolutionary Conservation of Species' Roles in Food Webs. *Science* **335**, 1489–1492 (2012).

- 1 6.Hembry, D. H., Yoder, J. B. & Goodman, K. R.*et al.* Coevolution and the
2 Diversification of Life. *The American Naturalist* **184**, 425–438 (2014).
- 3 7.Yoder, J. B. & Nuismer, S. L. *et al.* When does coevolution promote diversifi-
4 cation? *Am. Nat.* **176**, 802–817 (2010).
- 5 8.Paterson, S. *et al.* Antagonistic coevolution accelerates molecular evolution.
6 *Nature* **464**, 275–278 (2010).
- 7 9.Buckling, A., Wei, Y., Massey, R. C., Brockhurst, M. A. & Hochberg, M. E.*et al.*
8 Antagonistic coevolution with parasites increases the cost of host deleterious
9 mutations. *Proc. R. Soc. B* **273**, 45–49 (2006).
- 10 10.Brockhurst, M. A., Buckling, A., Poullain, V. & Hochberg, M. E.*et al.* The
11 impact of migration from parasite-free patches on antagonistic host-parasite
12 coevolution. *Evolution* **61**, 1238–1243 (2007).
- 13 11.Nuismer, S. L., Thompson, J. N. & Gomulkiewicz, R.*et al.* Coevolution
14 between hosts and parasites with partially overlapping geographic ranges. *J.*
15 *Evol. Biol.* **16**, 1337–1345 (2003).
- 16 12.Benkman, C. W., Parchman, T. L., Favis, A. & Siepielski, A. M.*et al.* Re-
17 ciprocal selection causes a coevolutionary arms race between crossbills and
18 lodgepole pine. *Am. Nat.* **162**, 182–194 (2003).
- 19 13.Havens, K. *et al.* Scale and structure in natural food webs. *Science* **257**,
20 1107–1109 (1992).
- 21 14.Bascompte, J. *et al.* Disentangling the Web of Life. *Science* **325**, 416–419
22 (2009).
- 23 15.Proulx, S., Promislow, D. & Phillips, P.*et al.* Network thinking in ecology

- 1 and evolution. *Trends Ecol. Evol.* **20**, 345–353 (2005).
- 2 16.Dunne, J. A. *et al.* in *Ecological networks: Linking structure and dynamics* (eds.
3 Dunne, J. A. & Pascual, M.) 27–86 (Oxford University Press, 2006).
- 4 17.Stouffer, D. B., Camacho, J., Jiang, W. & Amaral, L. A. N.*et al.* Evidence for
5 the existence of a robust pattern of prey selection in food webs. *Proc. R. Soc. B*
6 *Biol. Sci.* **274**, 1931–40 (2007).
- 7 18.Chamberlain, S. A. *et al.* Traits and phylogenetic history contribute to net-
8 work structure across Canadian plant–pollinator communities. *Oecologia* 1–12
9 (2014).
- 10 19.Vamosi, J. C., Armbruster, W. S. & Renner, S. S.*et al.* Evolutionary ecology
11 of specialization: insights from phylogenetic analysis. *Proc. R. Soc. B* **281**,
12 20142004 (2014).
- 13 20.Desdevises, Y., Morand, S., Jousson, O. & Legendre, P.*et al.* Coevolution
14 between *Lamellodiscus* (Monogenea: Diplectanidae) and Sparidae (Teleostei):
15 the study of a complex host-parasite system. *Evolution* **56**, 2459–2471 (2002).
- 16 21.Hafner, M. S. & Page, R. D. M. *et al.* Molecular phylogenies and host-
17 parasite cospeciation: gophers and lice as a model system. *Philos. Trans. R.*
18 *Soc. B Biol. Sci.* **349**, 77–83 (1995).
- 19 22.Hall, A. R., Scanlan, P. D. & Buckling, A.*et al.* Bacteria-Phage coevolution
20 and the emergence of generalist pathogens. *Am. Nat.* **177**, 44–53 (2010).
- 21 23.Roopnarine, P. D. & Angielczyk, K. D. *et al.* The evolutionary palaeoecology
22 of species and the tragedy of the commons. *Biol. Lett.* **8**, 147–50 (2012).
- 23 24.Dunne, J. a, Williams, R. J., Martinez, N. D., Wood, R. a & Erwin, D. H.*et al.*

- 1 Compilation and network analyses of cambrian food webs. *PLoS Biol.* **6**, e102
2 (2008).
- 3 25.Poisot, T., Stouffer, D. B. & Gravel, D.*et al.* Beyond species: why ecological
4 interaction networks vary through space and time. *Oikos* (2014).
- 5 26.Devictor, V. *et al.* Differences in the climatic debts of birds and butterflies at
6 a continental scale. *Nat. Clim. Change* **2**, 121–124 (2012).
- 7 27.Legendre, P., Desdevises, Y. & Bazin, E.*et al.* A statistical test for host-
8 parasite coevolution. *Syst. Biol.* **51**, 217–234 (2002).
- 9 28.Fahrenholz, H. *et al.* Ectoparasiten und abstammungslehre. *Zool. Anz.* **41**,
10 371–374 (1913).
- 11 29.Poisot, T. *et al.* in *Evolutionary Ecology of Host-Parasite Systems* (eds. Morand,
12 S., Littlewood, D. T. & Poulin, R.) (Cambridge University Press, 2015).
- 13 30.Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D.*et al.* The
14 dissimilarity of species interaction networks. *Ecol. Lett.* **15**, 1353–1361 (2012).
- 15 31.Data from: Phylogenetic signal in module composition and species connec-
16 tivity in compartmentalized host-parasite networks. (2012).
- 17 32.Balbuena, J. A., Míguez-Lozano, R. & Blasco-Costa, I.*et al.* PACo: A Novel
18 Procrustes Application to Cophylogenetic Analysis. *PLoS ONE* **8**, e61048 (2013).
- 19 33.Krasnov, B. R. *et al.* Phylogenetic signal in module composition and species
20 connectivity in compartmentalized host-parasite networks. *Am. Nat.* **179**,
21 501–11 (2012).
- 22 34.Price, P. W. *et al.* *Macroevolutionary Theory on Macroecological Patterns*. (Cam-
23 bridge University Press, 2003).