

Coevolution leaves a stronger imprint on interactions than on community structure

Timothée Poisot and Daniel B. Stouffer

December 15, 2014

1 Coevolutionary dynamics act on both species and their interactions
2 in ways that shape ecological communities. It remains unclear, how-
3 ever, how the structure of communities at larger spatial scales either
4 influences or is influenced by local coevolutionary processes, and how
5 mechanisms acting at these different scales feedback onto one another.
6 Here we show that, though species interactions vary substantially over
7 a continental gradient, the coevolutionary significance of individual in-
8 teractions is maintained across different scales. Notably, this occurs
9 despite the fact that observed community variation at the local scale
10 frequently tends to weaken or remove community-wide coevolutionary
11 signal. When considered in terms of the interplay between community
12 ecology and coevolutionary theory, our results demonstrate that indi-
13 vidual interactions are capable and likely to show a consistent signature
14 of past coevolution even when woven into communities that do not.

15 Ecological interactions often exert important selective pressures on the species
16 involved. For example, the phenologies of lodgepole pines and red crossbills re-

1 spond spatially to the presence of squirrels ¹ and palm species undergo changes
2 in seed morphology in response to the extinction of bird dispersing their seeds
3 ². Given that interactions are distributed in similar ways across communities, at
4 both the large ³ or small ⁴ scale, it can be argued that much ecological structure
5 is the end result of evolutionary or coevolutionary dynamics between species ^{5,6}.
6 Unfortunately, while the coevolutionary dynamic of pairs of interacting species
7 has been well described at macro ⁷ and micro ⁸ evolutionary timescales, most
8 attempts to understand how they cascade up to the levels of diversity of both
9 species and interactions found within empirical communities have been inconclu-
10 sive ⁹. Moreover, because coevolutionary dynamics are often presented as a key
11 driving force behind ecological structure across both time and space ¹⁰, it is crucial
12 to determine the scale at which they are both relevant and quantifiable.

13 Historically, the evidence for coevolution in taxonomically diverse communities
14 is quantified as the degree of matching between the phylogenies of two sets of
15 interacting organisms ¹¹. This notion builds on the century-old idea that extant
16 species interact in a way similar to the way their ancestors did ¹². Yet it is consid-
17 erably more restrictive than just phylogenetic conservation of species' interactions
18 ¹³, as it accounts for higher-order constraints. More explicitly, it is thought that
19 communities that have assembled by successive divergence events should display
20 phylogenetic congruence, that is (i) have similar phylogenetic trees and (ii) have
21 species at matching positions in the trees that tend to interact ¹⁴. On the other
22 hand, many ecological and evolutionary processes that occur locally are expected
23 to blur community-wide coevolutionary signal ¹⁵. One possible explanation is
24 that interactions can display substantial turnover at ecologically relevant tem-
25 poral and spatial scales ¹⁶: the same two species can interact in different ways

1 under the effect of local environmental contingencies, spatial mismatch in species
2 phenologies, variations in population abundances, and chance events ¹⁷. It is un-
3 clear, however, whether these mechanisms influence how the coevolutionary signal
4 within individual interactions should vary across spatial scales.

5 To answer these questions, we study a dataset of interactions between rodents
6 and their ectoparasites from 51 sites across Western to Eastern Europe ¹⁸ (Meth-
7 ods Summary). This dataset is uniquely suited for this task as it represents
8 a paradigmatic system in which species-species interactions are thought to be
9 driven by macro-evolution and co-speciation events ¹⁹, and coevolutionary signal
10 is indeed significant at the continental level ²⁰ ($p \leq 10^{-4}$; Methods Summary).
11 Importantly, it also provides spatial replication and variability at a scale large
12 enough to capture macro-ecological processes.

13 As host-macroparasites interactions are hypothesized to be both ecologically con-
14 strained and evolutionary conserved ²¹, the congruence observed at the continen-
15 tal level sets the baseline for what would be expected in local communities. Of
16 course, if ecological mechanisms reduce coevolutionary signal, we should detect
17 coevolution at the continental scale but not locally. Noting that variation of in-
18 teractions can decrease congruence, we analyse the data at two different levels to
19 test these hypotheses: first, we use *regional* interaction data—which accounts for
20 different species composition across sites—and second, we use the *local* interaction
21 data—which also accounts for variation in the interactions between observed these
22 species (Methods Summary). Out of 51 sites, 35 show no signal of coevolution, 11
23 show significant coevolutionary signal when using the regional interactions, and
24 12 show significant coevolutionary signal using the local interactions (see *Supp.*

1 *Mat. 1* for network-level significance values).

2 These results would appear to support the idea that macro-evolutionary processes
3 such as co-diversification can have consequences at the macro-ecological level ²²
4 but may not in fact be detectable at finer spatial scales. This is particularly
5 apparent when comparing the degree of network-level coevolutionary signal ob-
6 served at the local and regional scales to the same at the continental scale (Fig.
7 2A). On the other hand, system-level differences say little about the behavior of
8 individual interactions, despite the fact most coevolutionary mechanisms act at
9 the interaction level ²³. Intriguingly, we observe here that the distribution of indi-
10 vidual interactions' contributions to coevolution is strongly conserved, regardless
11 of the scale at which the interactions are quantified (Fig. 2B). As such, networks
12 with interactions that are important for coevolution at the continental scale re-
13 tain more coevolutionary signal at the local and regional scales alike (Fig. 2C).
14 Because interactions differ in their total contribution to coevolution, their distri-
15 bution across networks is what actually drives differences in overall coevolutionary
16 signal. Network-level coevolutionary signal emerges directly from the properties
17 of interactions and is not a property of the network itself.

18 Beyond their contribution to coevolution, interactions also ultimately differ in
19 how frequently they vary when the species involved co-occur ^{24,25}. Once more, the
20 literature on host-parasite interactions usually assumes that the reason why some
21 interactions are more frequent is because they reflect a significant past history
22 of coevolution ²⁶. If this were true, we should observe a significant, positive
23 correlation between the probability of observing an interaction and the importance
24 of that interaction for coevolution at the continental scale (Methods Summary).

1 Surprisingly, we find that neither is true here since interactions that are important
2 for coevolution are not more conserved (Fig. 3).

3 Nonetheless, interactions that *are* observed at the local scale should retain their
4 signal of contribution to coevolution because they involve species at matching po-
5 sitions in the phylogenetic trees. Indeed, we find that interactions that contribute
6 strongly to coevolutionary signal at the continental scale *also* show a significant
7 tendency to contribute strongly at the local scale (Fig. 4). Remarkably, this result
8 implies that the remnants of coevolution are still locally detectable in *individual*
9 *interactions* even though it does not leave its imprint on most local networks.

10 Overall, the results of our analyses demonstrate that there is a sizeable gap be-
11 tween our current understanding of coevolution as the basis of multi-species in-
12 teractions and its applicability to ecological questions. Local networks show little
13 to no signal of coevolution and the strength of coevolution between two species
14 is a surprisingly poor predictor of how frequently they interact. In contrast to
15 the frequent assumption that phylogenetic structure is a key driver of community
16 structure ²⁷, these data reveal that this impact is actually minimal at ecologi-
17 cally relevant spatial scales. Despite all the above, individual interactions are
18 able to maintain their coevolutionary signal even when the community they are
19 woven into does not. Thinking more broadly, these discrepancies provide a clear
20 roadmap for bridging the aforementioned gap between our appreciation of the role
21 of coevolution and its empirically measurable outcomes. Network structure is the
22 most parsimonious *mechanism* by which coevolution proceeds, not the imprint
23 coevolution leaves on ecological communities.

1 Methods Summary

2 We use data on observations of interactions between 121 species of rodents and
3 205 species of parasitic fleas in 51 locations across Europe ¹⁸ to build 51 species-
4 species interaction networks. Interactions were measured by combing rodents for
5 fleas, a method that gives high quality data as it has a high power of detection. To
6 account for differential sampling effort and across site variations in abundance, we
7 only study the networks' incidence matrices (presence and absence of interactions).

8 In our study, we define three scales for the network data and analysis— conti-
9 nental, regional, and local. The continental scale is the aggregated “metanetwork”
10 which includes all potential interactions between co-occurring species ¹⁶ (*i.e.*, all
11 species and all their interactions across the 51 networks). Within each site, the
12 regional scale is given by the list of observed species and all their possible interac-
13 tions. Hence the regional networks are always a perfect subset of the continental
14 network. The local scale includes only the interactions that were actually observed
15 in the field at a given site. Therefore, the local and regional networks always in-
16 clude the same species, but the local network has only a subset (or, at most, an
17 exact match) of the interactions in the regional network. The spatial consistency
18 of every individual interaction is measured as the number of sites in which the
19 two species involved co-occur.

20 We quantified the coevolutionary signal in terms of the degree of matching be-
21 tween host and parasite phylogenies given knowledge of species interactions using
22 the *PACO* method ²⁸, which is robust to variations in number of species. *PACO*
23 provides measures of both the network-level congruence (*i.e.*, is the network co-
24 evolved?) and the interaction-level signal (*i.e.*, what is the contribution of each

1 interaction to the overall coevolutionary signal?). We measured the phylogenetic
2 dissimilarity between two sites for hosts and parasites using PCD ²⁹, a measure
3 that accounts for the dissimilarity of species, corrected for the phylogenetic dis-
4 tance between all species in the dataset. Since it is a requirement of the methods
5 we use here, the phylogenetic trees for hosts and parasites were rendered ultra-
6 metric (i.e., all species are at the same distance from the root).

7 References

8 **Acknowledgments.** We thank Juan Antonio Balbuena for discussions about the
9 *PACo* method, and members of the Stouffer and Tylianakis groups for comments
10 on an early draft of this manuscript. Funding to TP and DBS was provided
11 by a Marsden Fund Fast-Start grant (UOC-1101) and to DBS by a Rutherford
12 Discovery Fellowship, both administered by the Royal Society of New Zealand.

13 1.Benkman, C. W., Parchman, T. L., Favis, A. & Siepielski, A. M. *et al.* Reciprocal
14 selection causes a coevolutionary arms race between crossbills and lodgepole pine.
15 *Am. Nat.* **162**, 182–194 (2003).

16 2.Galetti, M. *et al.* Functional Extinction of Birds Drives Rapid Evolutionary
17 Changes in Seed Size. *Science* **340**, 1086–1090 (2013).

18 3.Jordano, P., Bascompte, J. & Olesen, J. M. *et al.* Invariant properties in coevo-
19 lutionary networks of plant-animal interactions. *Ecol. Lett.* **6**, 69–81 (2003).

20 4.Stouffer, D. B., Camacho, J., Jiang, W. & Amaral, L. A. N. *et al.* Evidence for
21 the existence of a robust pattern of prey selection in food webs. *Proc. R. Soc. B*
22 *Biol. Sci.* **274**, 1931–40 (2007).




Figure 1: Spatial distribution of coevolutionary signal across the 51 sites. For each location, we indicate whether or nor the structure of regional and local interaction networks is consistent with phylogenetic congruence. The color of the circle corresponds to regionally significant or nonsignificant (black and grey, respectively) while the color of the symbol within corresponds to locally significant or nonsignificant (black and grey, respectively).




Figure 2: Distribution of coevolutionary signal at the network and interaction level. **A** Networks that have lower coevolutionary signal are made of interactions that contribute little to coevolution at the continental scale. **B** Interactions in the local, regional, and continental scale have equal contributions to coevolution.




Figure 3: Lack of a relationship between the spatial consistency of an interaction and its importance for coevolution in the continental network. Spatial consistency is defined as the probability of observing an interaction between two species given that they were observed to co-occur.




Figure 4: Relationship between continental and local contribution to coevolution. Each line corresponds to a location.

- 1 5.Eklof, A., Helmus, M. R., Moore, M., Allesina, S. & Eklöf, A.*et al.* Relevance
2 of evolutionary history for food web structure. *Proc. R. Soc. B Biol. Sci.* **279**,
3 1588–1596 (2011).
- 4 6.Stouffer, D. B., Sales-Pardo, M., Sirer, M. I. & Bascompte, J.*et al.* Evolutionary
5 Conservation of Species’ Roles in Food Webs. *Science* **335**, 1489–1492 (2012).
- 6 7.Van Valen, L. *et al.* A new evolutionary law. *Evol. Theory* **1**, 1–30 (1973).
- 7 8.Gandon, S., Buckling, A., Decaestecker, E. & Day, T.*et al.* Host-parasite co-
8 evolution and patterns of adaptation across time and space. *J. Evol. Biol.* **21**,
9 1861–1866 (2008).
- 10 9.Hembry, D. H., Yoder, J. B. & Goodman, K. R.*et al.* Coevolution and the
11 Diversification of Life. *The American Naturalist* **184**, 425–438 (2014).
- 12 10.Thompson, J. N. *et al.* *The Geographic Mosaic of Coevolution*. (University Of
13 Chicago Press, 2005).
- 14 11.Legendre, P., Desdevises, Y. & Bazin, E.*et al.* A statistical test for host-
15 parasite coevolution. *Syst. Biol.* **51**, 217–234 (2002).
- 16 12.Fahrenholz, H. *et al.* Ectoparasiten und abstammungslehre. *Zool. Anz.* **41**,
17 371–374 (1913).
- 18 13.Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P. & Bascompte,
19 J.*et al.* Non-random coextinctions in phylogenetically structured mutualistic net-
20 works. *Nature* **448**, 925–8 (2007).
- 21 14.Page, R. D. M. *et al.* *Tangled trees: Phylogeny, cospeciation, and coevolution*.
22 (University of Chicago Press, 2003).

- 1 15. Poiset, T. *et al.* in *Evolutionary Ecology of Host-Parasite Systems* (eds. Morand,
2 S., Littlewood, D. T. & Poulin, R.) (Cambridge University Press, 2015).
- 3 16. Poiset, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. *et al.* The
4 dissimilarity of species interaction networks. *Ecol Lett* **15**, 1353–1361 (2012).
- 5 17. Poiset, T., Stouffer, D. B. & Gravel, D. *et al.* Beyond species: why ecological
6 interaction networks vary through space and time. *Oikos* n/a–n/a (2014).
- 7 18. Data from: Phylogenetic signal in module composition and species connectivity
8 in compartmentalized host-parasite networks. (2012).
- 9 19. Verneau, O., Du Preez, L. & Badets, M. *et al.* Lessons from parasitic flatworms
10 about evolution and historical biogeography of their vertebrate hosts. *C. R. Biol.*
11 **332**, 149–158 (2009).
- 12 20. Krasnov, B. R. *et al.* Phylogenetic Signal in Module Composition and Species
13 Connectivity in Compartmentalized Host-Parasite Networks. *Am. Nat.* **179**,
14 501–511 (2012).
- 15 21. Combes, C. *et al.* *Parasitism - The Ecology and Evolution of Intimate Inter-*
16 *actions*. (University Of Chicago Press, 2001).
- 17 22. Price, P. W. *et al.* *Macroevolutionary Theory on Macroecological Patterns*.
18 (Cambridge University Press, 2003).
- 19 23. Thompson, J. N. *et al.* The raw material for coevolution. *Oikos* **84**, 5–16
20 (1999).
- 21 24. Carstensen, D. W., Sabatino, M., Trøjelsgaard, K. & Morellato, L. P. C. *et al.*
22 Beta Diversity of Plant-Pollinator Networks and the Spatial Turnover of Pairwise
23 Interactions. *PLoS ONE* **9**, e112903 (2014).

- 1 25.Olito, C. & Fox, J. W. *et al.* Species traits and abundances predict metrics of
2 plant–pollinator network structure, but not pairwise interactions. *Oikos* n/a–n/a
3 (2014).
- 4 26.*Biogeography of host-parasite interactions*. (Oxford University Press, 2010).
- 5 27.Cavender-Bares, J., Kozak, K. H., Fine, P. V. A. & Kembel, S. W.*et al.* The
6 merging of community ecology and phylogenetic biology. *Ecol. Lett.* **12**, 693–715
7 (2009).
- 8 28.Balbuena, J. A., Míguez-Lozano, R. & Blasco-Costa, I.*et al.* PACo: A Novel
9 Procrustes Application to Cophylogenetic Analysis. *PLoS ONE* **8**, e61048 (2013).
- 10 29.Ives, A. R. & Helmus, M. R. *et al.* Phylogenetic Metrics of Community Simi-
11 larity. *The American Naturalist* **176**, E128–E142 (2010).