

Quantifying the manageability of pollination networks in an invasion context

E. Fernando Cagua^{1*}, Kate L. Wootton^{1,2§}, Daniel B. Stouffer^{1#}

¹ Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8041, New Zealand

² Current address: Department of Ecology, Swedish University of Agricultural Sciences, Box 7044, SE-750 07 Uppsala, Sweden

* efc29@uclive.ac.nz § kate.l.wootton@gmail.com # daniel.stouffer@canterbury.ac.nz

• **Running title:** Manageability of pollination networks.

• **Keywords:** Network control, ecological networks, species importance, driver species, food webs, mutualism, management interventions, food web, interaction asymmetry, mutual dependence

• **Article type:** Letter.

• **Number of words:** 148 abstract; approx. 4,974 main text.

• **Number of references:** 65

• **Number of displays:** 5 figures; 1 table.

• **Correspondence to:** Daniel Stouffer (daniel.stouffer@canterbury.ac.nz) - +64 3 364 2729 - Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

• **Authorship:** DBS conceived the idea; all authors contributed to the development of the theoretical framework. EFC performed all analysis. EFC and DBS wrote the manuscript. All authors contributed to its revision.

• **Data accessibility:** All data used in this manuscript have already been published by Lopezaraiza-Mikel *et al.* (2007), Bartomeus *et al.* (2008), and Ballantyne *et al.* (2015). The reader should refer to the original sources to access the data.

Abstract

Here, we leverage recent advances in control theory to assess the “manageability” of ten pairs of uninvaded and invaded plant-pollinator communities. We found that the networks’ manageability is most strongly determined by the ratio of plants to pollinator richness, which in turn constrains the networks’ degree distribution. We also characterise species’ suitability for inclusion in management interventions by exploring the entire space of control strategies. Specifically, we measure the extent to which species (i) are necessary to steer the state of the community and (ii) are able to affect the abundance of other species. We found that invasive plants have a dominant position in every invaded community and that this dominance is underpinned by high asymmetry in the dependence of their interaction partners. Our results highlight the advantages of the control-theoretic framework for ecological questions and provide novel insight into the design of informed management interventions.

Introduction

In a complex system, the whole is often greater than the sum of its parts (Jørgensen *et al.* 1998; Levin 1999; Montoya *et al.* 2006). Within community ecology, a complex-systems approach has led to the development of analytical and simulation tools with which to understand, for example, the role of species embedded in a network of interactions (Pascual & Dunne 2005; Bascompte & Stouffer 2009; Stouffer *et al.* 2012). The inherent complexity of nature, however, has regularly hindered—or at least complicated—our ability to find management solutions to many problems ecological communities face. To overcome this obstacle, we require a framework that allows us to explain, predict, and manage ecological communities, particularly when they are confronted with perturbations (Solé & Montoya 2001; Green *et al.* 2005). Ideally, such a framework is equipped to account for their complex structure and the dynamics that determine the species abundances and the state of the community.

Among the various possibilities, *control theory* appears to be a strong candidate (Isbell & Loreau 2013). Widely used in engineering to determine and supervise the behaviour of dynamical systems (Motter 2015), it is well equipped to deal with the many feedbacks present in ecological communities (Liu & Barabási 2016). Research in this area has established a strong link between the structure of complex networks and their *controllability*—the relative ability to manipulate network components to drive the system to a desired state (Liu *et al.* 2011; Cornelius *et al.* 2013; Ruths & Ruths 2014). These advances suggest that it is in principle possible to alter a whole ecological community’s composition by modifying the abundances of only a few species. Applications of control theory to ecological networks can also take into account the extent to which changes in the abundances of one species may ripple through the community (Cornelius *et al.* 2013). Therefore, control theory could also be harnessed to help identify which species are most relevant from a structural and dynamic perspective.

This information is valuable not only for basic ecology, but it might be also relevant to address more applied management and conservation challenges. This is particularly true in the context of biotic invasions, where identifying key players in the community is a prerequisite to informed attempts to alter the state of invaded ecosystems and maintain the state of uninvaded ones. Despite recent advances in network theory, practical challenges to the conservation of interaction networks persist (Tylianakis *et al.* 2010), and the link between the structure of complex networks and our ability to manage and conserve them is still ambiguous (Blüthgen 2010; Kaiser-Bunbury & Blüthgen 2015). To complicate things further, biotic invasions can induce dramatic changes in the patterns of interactions that determine the structure of ecological networks (Baxter *et al.* 2004; Tylianakis *et al.* 2008; Ehrenfeld 2010), in particular pollination (Olesen *et al.* 2002; Aizen *et al.* 2008; Bartomeus *et al.* 2008; Vilà *et al.* 2009; Traveset *et al.* 2013). Understanding how the differences in network structure before and after invasion impact our ability to manage the communities is thus a double challenge, but it is also the critical first step towards a fully

informed recovery. Despite the apparent overlap, the control-theoretic perspective has not been adopted in an invasion context.

To bridge this gap, we outline an approach to apply control theory in an ecological context and implement it using empirical data. Specifically, we use a set of ten pairs of uninvaded and invaded plant-pollinator communities to investigate the link between invasion, network structure and ecological management. While doing so, we focus on two particular questions. First, grounded in the difficulties usually involved with invasive-species eradication and ecosystem restoration (Woodford *et al.* 2016), we ask whether invaded networks have lower levels of “manageability” than their uninvaded counterparts; that is, whether they require a greater proportion of species to be managed to achieve the same level of control. Second, we ask whether some species are more important than others at driving the population dynamics of the community and which factors determine this importance.

Methods

Theoretical framework

Disregarding practical considerations, any network (community) could hypothetically be fully controlled if we control the state of every single node (species) individually. At the core of control theory of complex networks, however, rests the idea that the state of a node depends on the state of the nodes it interacts with, and the form of this dependence is determined by both the dynamic relationship among interacting nodes and the structure of the links in the network. This principle can therefore be harnessed to find a subset of *driver nodes* to which to apply external input signals which then drive the state of every other node in the network to a desired configuration (Liu & Barabási 2016).

Conveniently the information necessary to determine the minimum number of driver nodes D in a network with linear dynamics is fully contained in the network structure (Kalman 1963; Liu *et al.* 2011; Motter 2015). We therefore start here from the assumption that an ecological network near equilibrium points can be described by $\frac{dx}{dt} = Ax + Bu(t)$, where the change of its state over time ($\frac{dx}{dt}$) depends on its current state x (for example the species’ abundances), an external time-varying input $u(t)$, and two matrices A and B , which encapsulate information about the network structure and how the species respond to the external input, respectively. Even though linearity might be a strong assumption, it has been shown that fundamental insight about the control of complex systems can be gained without a detailed knowledge of the nonlinear dynamics and the system parameters (Liu & Barabási 2016), an important step towards ultimately understanding the controllability of systems with nonlinear dynamics (Liu *et al.* 2011). This means that structural controllability can be applied to a wide range of network representations

of ecological communities to provide an indication of our ability to control them. Moreover, we show here how quantitative data about species interactions can help us move past some of the limitations of structural controllability by better approximating the role of interspecific dynamic relationships without being overly dependent on the choices of how these dynamics are modelled or characterised.

Manageability

The number of driver nodes D provides a structural indication of how difficult a network’s control might be. This is because systems that require a large number of external input signals are intuitively more difficult or costly to control. In an ecological context, external inputs that modify the state of a node can be thought of as management interventions. Therefore, the density of driver nodes $n_D = \frac{D}{S}$, where S is the total number of species in the community, is a measure of the extent to which network structure can be harnessed for network control. For instance, a hypothetical “network” in which species do not interact would require direct interventions for every single species to achieve full control, whereas a linear food chain would require just one species to be directly controlled to harness cascading effects through its trophic levels. From this perspective, it is possible to use n_D as an index of the manageability of an ecological community, understood in the context of how difficult is to modify the abundances of species in the community using external interventions—a common theme in ecosystem management, conservation, and restoration.

It has been recently shown that calculating D is equivalent to finding a maximum matching in the network (Liu *et al.* 2011). In a directed network, a *matching* consists of a subset of links in which no two of them share a common starting or ending node (Figure 1, Supporting Information S1). A given matching has *maximal cardinality* if the number of matched links (also referred to as the *matching size*) is the largest possible. A maximal cardinality matching is then called a *maximum matching* if the sum of the weights of the matched links (also referred to as *matching weight*) is again the largest possible (West 2001).

Once we have the subset of links that constitute a matching, we can also classify the nodes in the network based on that matching (Figure 1). A node is called *matched* if it is at the end of a matched link and *unmatched* otherwise. A node is also called *superior* if it is at the start of a matched link. Note that a node cannot be superior if it has no outgoing links. Notably, these node categories are what helps us to link a maximum matching back to the concept of network controllability, as follows. Unmatched nodes are the *driver nodes* D because they have no superior in the network and must be directly controlled by an external input (Liu *et al.* 2011). Each matched node, on the other hand, can be controlled by its superior. Note that this framework requires a directed network in which the direction of the links corresponds to the direction of control. In the “Empirical application” section below, we explain our approach to determining the link direction in pollination networks.

Relative importance

While calculating n_D measures the manageability of an ecological community, it does not provide information about the identity of the species that compose the set of driver nodes. Ecologically, potential differences between species are relevant because management and conservation resources are limited, and therefore ecological interventions should be focused on the set of species that might provide the largest impact. Moreover, maximum matchings in a network are often not unique, and each maximum matching indicates unique paths that can potentially be used to control the network. We harness this property and use a network's complete set of maximum matchings to characterise each species' relative importance in driving the state of the community. One possibility is to characterise a species by the frequency f_D with which it is classified as a driver node within this set of matchings. However, the profile of our networks indicates that a large proportion of species were classified as driver nodes because of external interventions required to achieve **full** controllability and not because they influence the abundance of other species (Supporting Information S2). Furthermore, the precise role of driver nodes is more ambiguous when full control is unfeasible or undesired—often the case in ecological settings. We therefore also calculate the frequency f_S with which a species is classified as a superior node since this is the frequency with which they form part of possible control paths.

Most commonly, structural controllability assumes unweighted networks—links exist or not, and hence f_D and f_S can be calculated by computing all possible maximum-cardinality matchings. However, we take the link weights into account when calculating the matchings here because the weights can reveal significant ecological patterns and processes that might be undetectable in unweighted networks (Scotti *et al.* 2007; Tylanakis *et al.* 2007; Vázquez *et al.* 2007; Kaiser-Bunbury *et al.* 2010). Additionally, species A may interact with both species B and C but depends strongly on B and only weakly on C . Intuitively, a management intervention designed to indirectly modify the abundance of species A is more likely to succeed if the abundance of B , rather than C , is directly controlled. A complication of including the interaction weight when calculating the maximum matching, however, is that empirical interaction strengths are to some extent stochastic and depend on proximate factors such as sampling method and intensity (Gibson *et al.* 2011). We overcome this issue by calculating all maximal cardinality matchings and then ranking them by their matching weight. By following this approach, we effectively give priority to the species that participate in the pathways that potentially have the largest impact on the community while acknowledging the limitations associated with sampling and its potential restrictions (Jordano 2016).

Empirical application

We next applied the previously defined framework to ten paired pollination networks. Each network pair was composed of a community invaded by a plant and a community “free” of the invasive species. Four pairs were obtained from natural or semi-natural vegetation communities in the city of Bristol, UK (Lopezaraiza-Mikel *et al.* 2007). These networks are comprised of 19–87 species (mean 55), and non-invaded plots were obtained by experimentally removing all the flowers of the invasive species *Impatiens grandulifera*. The other six pairs were obtained from lower diversity Mediterranean shrublands in Cap de Creus National Park, Spain (Bartomeus *et al.* 2008). These networks are comprised of 30–57 species (mean 38); in contrast to the above, uninvaded communities were obtained from plots that had not yet been colonised by either of the invasive species *Carpobrotus affine acinaciformis* or *Opuntia stricta*. Further details about the empirical networks can be found in the Supporting Information S3.

We then specified the structure of all networks using pollinator visitation frequency, which has been shown to be an appropriate surrogate for interspecific effects in pollination networks (Vázquez *et al.* 2005; Bascompte *et al.* 2006). To further examine whether this decision would influence our results, we also evaluated the effect of using pollinator efficiency or pollinator importance as alternative measures in a different data set that lacked invasive species (Ne’Eman *et al.* 2010; Ballantyne *et al.* 2015), and we found quantitatively similar results for all three options (Supporting Information S4). Because our approach depends on the network topology, we also evaluated the robustness of our results to the undersampling of interactions. Specifically, we calculated n_D and species relative importance for 500 random subsamples of each empirical network in which the weakest links were more likely to be removed. Our sensitivity analysis indicated that, even in the absence of complete sampling, a control-theoretic approach can still be applied (Supporting Information S5).

Manageability

We began by quantifying the manageability of each network. To do so, we calculated the networks’ maximum matching and determined the minimum proportion of species n_D that need external input signals to fully control the species abundances in the community. Note that because all maximum matchings have the same matching size, it is only necessary to calculate one of them. To simplify the analysis, if a network had more than one component (two species are in different components if there exists no path between them and are hence independent of each other in terms of network control) we only considered the largest. Smaller components were present in eleven out of the twenty networks and were typically composed of just one plant and one pollinator. Their removal represented an average loss of 4.7% of the species and 2.7% of the interactions.

As we noted earlier, the maximum-matching approach requires a directed network in which a link from species i to species j indicates that the abundance of j can be affected by the abundance of i . This implies that we need first to identify a directionality for the links between species that is consistent with the dynamics of the community (Figure 2). In some ecological networks, establishing the directionality can appear relatively straightforward, for example when links represent biomass transfer or energy flow (Isbell & Loreau 2013). Interspecific effects in pollination networks, however, are not strictly directed since the benefit is mutual between interacting species. Nevertheless, the relative extent to which a given pair of interacting species affect each other can be quantified by the magnitude of the mutual dependence between them (Bascompte *et al.* 2006). The dependence of plant i on pollinator j , d_{ij} , is the proportion of the visits from pollinator j compared to all pollinator visits to plant i . Likewise, the dependence of pollinator j on plant i , d_{ji} , is the proportion of the visits by pollinator j to plant i compared to all visits of pollinator j . Using dependences generates a weighted bipartite network in which all interacting pairs are connected by two directed links (Figure 2b).

Given the respective dependences, the extent to which species i affects species j relative to the extent to which j affects i can be summarised by the interaction asymmetry (Bascompte *et al.* 2006) given by

$$a(i, j) = a(j, i) = \frac{|d_{ij} - d_{ji}|}{\max(d_{ij}, d_{ji})}.$$

Previous research has shown that mutual dependences are often highly asymmetric in natural communities (Bascompte *et al.* 2006); in other words, if a plant species is largely dependent on a pollinator species, then that pollinator tends to depend rather weakly on the plant (or vice versa). We therefore simplified the network so that interacting species are connected by only one directed link when mutual dependences are asymmetric (Figure 2c). This simplification, while maintaining ecological realism, is advantageous for several reasons. First, it is consistent with previous advances in structural controllability; second, it avoids complications related to the introduction of artificial control cycles (Ruths & Ruths 2014); and third it significantly reduces the computational resources necessary for the application of our approach (Supporting Information S6). Moreover, we found that changing to unidirectional interactions based on the direction of asymmetry does not alter the relative n_D of different networks (Table S3).

To find a maximum matching in a network with interaction directions and weights determined by the asymmetry, we adopted a strategy based on an alternative bipartite representation of the directed network with two levels that indicate the outgoing and incoming links to each node (Supporting Information S1). Once we had this alternative representation we used the maximum bipartite matching algorithm implemented in the `max_bipartite_match` function of `igraph` 1.0.1 (Csardi & Nepusz 2006) on each

network.

Statistical analysis

We also wanted to test whether invasion status or other predictors had an impact on the observed values of n_D . We therefore used a set of generalised linear models (with binomial error structure) to investigate the effect of invasion status while also including covariates related to species richness, since one might naively expect to see a negative relationship between richness and manageability (Menge 1995). These covariates included the total number of species, plant richness, pollinators richness, the ratio of plant to pollinator richness, the link density (connectance), and the study site (as a two-level factor).

We next explored whether real networks differ in their architecture from random ones in a concerted way that affects manageability. Previous research indicates a direct link between a network's degree distribution and the number of nodes necessary to fully control it (Liu *et al.* 2011), but the strength and applicability of this relationship have not been tested for in weighted ecological networks. We therefore compared the driver-node density n_D of the empirical networks to networks generated by a null model that maintained each species' strength (its total sum of visits) while allowing their degrees (its number of interactions) to vary. Beyond network structure, the dependence asymmetry plays a fundamental role in determining the direction of control in each two-species interaction and therefore has the potential to influence the network n_D results above. We therefore performed an additional randomisation in which we kept the structure of each network constant but randomised the direction of the interaction asymmetries. That is, we first calculated the observed asymmetries for each community and then shuffled the direction of the link between each pair of species.

Additional details about the statistical models and the randomisations can be found in the Supporting Information S7.

Relative importance

Our second key question was related to how species differ in their ability to drive the population dynamics of the community. To quantify this importance, we computed all maximal cardinality matchings in each network. We then calculated the frequency with which each species i was a driver (f_D) or a superior node (f_S) in the set of matchings that had a matching weight greater or equal to 0.8 times the weight of the maximum matching. We selected this threshold as it provided a high agreement between networks quantified by visitation and pollination efficiency as well as between our weighting/directionality assumptions; however, the choice of this threshold had a negligible impact on any results (Supporting Information S8). Details about the computational procedure to find all maximal cardinality matchings of a network can be found in Supporting Information S1 and Figure S2.

250 We then examined whether any species-level structural properties could predict our metrics of species
 251 importance—the frequency with which a species was a driver or a superior node (f_D and f_S , respectively).
 252 We used a set of generalised linear mixed-effects models (with binomial error structure) with the relative
 253 frequencies as the response variables. As predictors in this model, we included measures of centrality
 254 (degree and eigen-centrality), which have been found to be strong predictors of importance in a coextinction
 255 context (Memmott *et al.* 2004); a measure related to network robustness (contribution to nestedness), as
 256 nestedness has been proposed as one of the key properties that promote stability in mutualistic networks
 257 (Saavedra *et al.* 2011); a measure of strength of association (visitation strength, the sum of visits a
 258 species receives or performs) and a measure of strength of dependence (species strength, the sum of
 259 dependences of all species on the focal species), as their distribution determines the extent of interspecific
 260 effects (Bascompte *et al.* 2006). In addition, we also included guild (plant or pollinator) and whether the
 261 species is invasive or not as categorical fixed effects. Lastly, we allowed for variation between different
 262 communities by including the network identity as a random effect (Bates *et al.* 2015). Supplementary
 263 details about the statistical models can be found in the Supporting Information S7.

Results

Manageability

264 All of the networks had a driver-node density n_D between 0.55 and 0.92 (mean 0.76; Figure 3a). In addition,
 265 we found that, when controlling for potential species richness effects, the n_D of invaded communities
 266 was smaller to those of non-invaded communities (Figure 3b). Nevertheless, of the various covariates we
 267 explored, the ratio of plants to pollinators showed the strongest relationship with n_D (Figure 3c; Table
 268 S5). Specifically as the proportion of pollinators increases and the ratio plant/pollinator approaches unity,
 269 n_D decreases (all our communities had more pollinators than plants). Other covariates—connectance (link
 270 density) and species richness—had negative, but comparatively less important, relationships with n_D .

271 When exploring the effect of network structure itself, we observed that the driver-node density n_D of
 272 empirical networks was, in general, not significantly different to the manageability of network randomi-
 273 sations that maintained the degree of individual species (Figure 4). However, we found that the n_D of
 274 empirical networks was significantly larger than that of randomisations that maintained the network
 275 structure but that differed only in the direction of the asymmetries.

Relative importance

Invasive species were classified as superior nodes *and* driver nodes in every single network they were present; that is, they always had the highest relative f_S and f_D (Figure 5a). The model results suggest that these differences between invasive and native species are not underpinned by any intrinsic property of the invasive species; instead, they are due to species properties that apply to invasive and native species alike (Table 1). Specifically, we found that a species is more likely to be classified as a superior node if it had a large species strength (the sum of the dependences of all other species on the species of interest). To a smaller extent, visitation strength (a species' sum of visits) and degree also had a positive relationship with f_S . In contrast, the relationships between species structural properties and f_D were less clear cut (Table 1). Both invasive species (which generally have a larger degree and high dependence strength) and pollinators (which generally have a smaller degree and low dependence strength) were classified as driver nodes in a large proportion of matchings.

Discussion

Contrary to our initial hypothesis, we found some evidence that invaded communities might be easier to manage than uninvaded ones from a control-theoretic perspective. Our results reveal, however, that this effect is comparatively small, and the structural differences among different networks are more strongly related to potential differences in our ability to alter the state of the community via external interventions. Despite the small effect of invasion status at the network level, we found that invasive mutualists occupy a particularly dominant role in their communities for two reasons. First, as species with a high f_S , changes on their abundance have the potential to propagate broadly through the community and, in turn, affect the abundances of many other species. Second, as species with a high f_D , they are also indispensable when it comes to fully controlling the plant-pollinator network. At a community level, we demonstrate that the manageability of mutualistic networks is strongly governed by the asymmetric nature of mutual dependences—which constitute the foundations of the structure and stability of mutualistic networks (Memmott *et al.* 2004; Vázquez & Aizen 2004; Bascompte *et al.* 2006; Lever *et al.* 2014; Astegiano *et al.* 2015). Moreover, these mutual dependences seem to be constrained by the effects of both the patterns of species richness at each trophic guild and a network's degree distribution (Melián & Bascompte 2002; Blüthgen *et al.* 2007). Indeed, the difference between the driver-node density (n_D) of our empirical networks and that of randomisations depended strongly on the null model's randomisation approach. While the empirical n_D was indistinguishable from that of networks with a random structure that maintained the degree of each species in the community, it was larger than that of randomisations in which the directed network was unchanged but where the observed patterns of dependence were broken.

Invasive species have been previously found to exacerbate the asymmetries in their communities (Aizen *et al.* 2008; Bartomeus *et al.* 2008; Henriksson *et al.* 2016). Although this might cause differences both at the community and the species level, we found that invasive plants are not inherently different to their native counterparts (Stouffer *et al.* 2014; Emer *et al.* 2016). Invasive plants, just like any other mutualist in our data set, tend to be classified as a superior node proportional to the degree to which their interaction partners are collectively more dependent on them than the other way around. Previous studies have found that supergeneralists, like invasive species, play a central role in their networks (Vilà *et al.* 2009; Palacio *et al.* 2016). Our results take this one step further and indicate that dependence strength, rather than generalism or other metrics of centrality, is the factor that best explains the cascading effects a species could trigger on its community.

Because of the ability that our approach has to infer the magnitude of the effects that each species has on others in the community, it is tantalising to use it to select promising candidates for management interventions. To this end, the two indices we have used to characterise a species provide two complementary pieces of information. Our first index f_D —the frequency with which a species is classified as a driver node—provides an indication of the likelihood that a species forms part of the set of species that must be manipulated in order to control *all* species in the community. This driver-node concept has received considerable attention in the structural-control literature and indeed shows substantial potential to provide useful ecological insight. Nevertheless, we anticipate two caveats that hinder its direct utility for management applications. First, unlike some other types of complex systems, fully controlling an ecological community is almost certainly out of reach for all but the simplest, due to either the number of required interventions or the practical difficulties of their implementation (Motter 2015). For instance, our results suggest that full control of the pollination networks would require direct interventions on anywhere from 40–90% of the species. Second, Ruths & Ruths (2014) established that driver nodes arise due to distinct mechanisms and therefore species with markedly different network metrics can act as driver nodes in their community (Supporting Information S2).

Our second index f_S , however, is directly related to the likelihood a species affects the abundance of another species in all of the control strategies considered. Importantly, this is irrespective of whether controlling the entire network is ultimately desired and/or feasible. In fact, because superior nodes are always at the beginning of a matched link, species with a high f_S are more likely to be the subjects of management interventions when controlling the abundances of a target set of species—as opposed to the entire network—is desired (Gao *et al.* 2014). An important advantage here is that the target set of species does not have to be the same set to which interventions are applied. For instance, despite inconsistent outcomes in practice (Suding *et al.* 2004; Rodewald *et al.* 2015; Smith *et al.* 2016), our results suggest that current restoration approaches that focus on direct eradication of invasive species might indeed be

an effective way to modify ecosystem state. Nevertheless, our results also indicate that removals must be exercised with caution. Not only it is hard to predict the direction in which the system will change, but invaded communities also tend to be highly dependent on invaders and therefore acutely vulnerable to their eradication (Traveset *et al.* 2013; Albrecht *et al.* 2014).

Despite the apparent similarities, our approach is different to previous attempts to quantify species importance in a few key ways. Existing metrics usually harness species features, like centrality, position, co-extinction or uniqueness, to infer their effect on other species (Allesina & Bodini 2004; Jordán *et al.* 2006; Jordán 2009; Lai *et al.* 2012). In contrast, our control-based approach tackles that question directly. Although they are relatively simple to calculate, classic species-level network metrics do not necessarily reveal the best set of species to manage (Eklöf *et al.* 2013; McDonald-Madden *et al.* 2016). Our approach, however, is not based on a single structural metric but instead acknowledges the existence of multiple management strategies. By allowing for the fact that some strategies are better than others depending on the context, control theory implicitly highlights that management decisions should not be based on a single technique. As such, ours and other flexible approaches that take a network-wide approach might prove more useful to guide ecosystem management (McDonald-Madden *et al.* 2016).

In this study, we illustrate how a control-theoretic approach can be better adopted in network ecology to evaluate the effect of invasions and other kinds of perturbations. Although our pollination specific results might not be directly translatable to other ecological systems because of the degree constraints imposed by bipartite networks, the approach we propose is applicable wherever species abundances are influenced by their interactions, and exciting open questions lie ahead. How to design the precise “control signals” to reach a desired ecosystem state or conservation outcome? What are the implications of assuming linear species dynamics? How important it is to include several interaction types for our understanding of manageability and species importance? What are the implications for species coexistence? Which are the trade-offs between persistence at the species and the community level? Answering each of these questions might require its evaluation in different ecological systems, an explicit integration of control theory with numerical models of species densities (Cornelius *et al.* 2013; Gibson *et al.* 2016), and experimental tests on simple communities. Nevertheless, the potential rewards are encouraging from both an ecological and conservation perspective, where an integrated approach can shift our focus beyond the identification of ideal targets for intervention to the design of informed interventions that legitimately achieve restoration goals.

Appendix 1: Glossary

Driver node An unmatched node in a maximal cardinality matching or a maximum matching. From the control perspective, driver nodes are those to which external control signals must be applied in order to gain full control of the network.

Matched/unmatched link A link is referred to as *matched* if it is part of a matching, and *unmatched* otherwise.

Matched/unmatched node A node is referred to as *matched* if it is at the end of a matched link, and *unmatched* otherwise.

Matching A set of links in which no two of them share a common starting or ending node.

Matching size The number of matched links in a matching.

Matching weight The sum of the weights of all matched links in a matching.

Maximal cardinality matching A matching with the largest possible matching size. In unweighted/binary networks, all maximal cardinality matchings are also maximum matchings.

Maximum matching A matching with the largest possible matching size *and* largest possible matching weight.

Superior node The node at the start of a matched link. From the control perspective, superior nodes make up the chains that propagate the control signals through the network.

Acknowledgments

The authors thank Dr. Takeuki Uno for the insight provided to find the set of all maximum matching algorithms, and Jason Tylianakis, Bernat Bramon Mora, Guadalupe Peralta, Rogini Runghen, Warwick Allen, Matthew Hutchinson, and Marilia Gaiarsa for feedback and valuable discussions. EFC acknowledges the support from the University of Canterbury Doctoral Scholarship, the University of Canterbury Meadow Mushrooms Postgraduate Scholarship, a New Zealand International Doctoral Research Scholarship, and a travel grant from the European Space Agency. DBS acknowledges the support of a Marsden Fast-Start grant and a Rutherford Discovery Fellowship, administered by the Royal Society of New Zealand.

References

- Aizen, M.A., Morales, C.L. & Morales, J.M. (2008). Invasive mutualists erode native pollination webs. *PLOS Biology*, 6, 0396–0403.
- Albrecht, M., Padrón, B., Bartomeus, I. & Traveset, A. (2014). Consequences of plant invasions on

396 compartmentalization and species' roles in plant-pollinator networks. *Philosophical Transactions of the*
397 *Royal Society B: Biological Sciences*, 281, 20140773.

398 Allesina, S. & Bodini, A. (2004). Who dominates whom in the ecosystem? Energy flow bottlenecks and
399 cascading extinctions. *Journal of Theoretical Biology*, 230, 351–358.

400 Astegiano, J., Massol, F., Vidal, M.M., Cheptou, P.-O. & Guimarães, P.R. (2015). The Robustness of
401 Plant-Pollinator Assemblages: Linking Plant Interaction Patterns and Sensitivity to Pollinator Loss.
402 *PLOS ONE*, 10, e0117243.

403 Ballantyne, G., Baldock, K.C.R. & Willmer, P.G. (2015). Constructing more informative plant-pollinator
404 networks: visitation and pollen deposition networks in a heathland plant community. *Proceedings of the*
405 *Royal Society B*, 282, 20151130.

406 Bartomeus, I., Vilà, M. & Santamaría, L. (2008). Contrasting effects of invasive plants in plant-pollinator
407 networks. *Oecologia*, 155, 761–770.

408 Bascompte, J. & Stouffer, D.B. (2009). The assembly and disassembly of ecological networks. *Philosophical*
409 *Transactions of the Royal Society B: Biological Sciences*, 364, 1781–7.

410 Bascompte, J., Jordano, P. & Olesen, J.M. (2006). Asymmetric Coevolutionary Networks Facilitate
411 Biodiversity Maintenance. *Science*, 312, 431–433.

412 Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models using lme4.
413 *Journal of Statistical Software*, 67, 1–48.

414 Baxter, C.V., Fausch, K.D., Murakami, M. & Chapman, P.L. (2004). Fish invasion restructures stream
415 and forest food webs by interrupting reciprocal prey subsidies. *Ecology*, 85, 2656–2663.

416 Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: A critique
417 and an ecologist's guide. *Basic and Applied Ecology*, 11, 185–195.

418 Blüthgen, N., Menzel, F., Hovestadt, T., Fiala, B. & Blüthgen, N. (2007). Specialization, Constraints,
419 and Conflicting Interests in Mutualistic Networks. *Current Biology*, 17, 341–346.

420 Cornelius, S.P., Kath, W.L. & Motter, A.E. (2013). Realistic control of network dynamics. *Nature*
421 *Communications*, 4, 1942.

422 Csardi, G. & Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal,*
423 *Complex Systems*, 1695, 1–9.

424 Ehrenfeld, J.G. (2010). Ecosystem Consequences of Biological Invasions. *Annual Review of Ecology,*
425 *Evolution, and Systematics*, 41, 59–80.

426 Eklöf, A., Tang, S. & Allesina, S. (2013). Secondary extinctions in food webs: A Bayesian network

approach. *Methods in Ecology and Evolution*, 4, 760–770.

Emer, C., Memmott, J., Vaughan, I.P., Montoya, D. & Tylianakis, J.M. (2016). Species roles in plant-pollinator communities are conserved across native and alien ranges. *Diversity and Distributions*, 22, 841–852.

Gao, J., Liu, Y.-Y., D'Souza, R.M. & Barabási, A.-L. (2014). Target control of complex networks. *Nature Communications*, 5.

Gibson, R.H., Knott, B., Eberlein, T. & Memmott, J. (2011). Sampling method influences the structure of plant-pollinator networks. *Oikos*, 120, 822–831.

Gibson, T.E., Bashan, A., Cao, H.-t., Weiss, S.T. & Liu, Y.-y. (2016). On the Origins and Control of Community Types in the Human Microbiome. *PLOS Computational Biology*, 12, e1004688.

Green, J.L., Hastings, A., Arzberger, P., Ayala, F.J., Cottingham, K.L. & Cuddington, K. *et al.* (2005). Complexity in Ecology and Conservation: Mathematical, Statistical, and Computational Challenges. *BioScience*, 55, 501–510.

Henriksson, A., Wardle, D.A., Trygg, J., Diehl, S. & Englund, G. (2016). Strong invaders are strong defenders - implications for the resistance of invaded communities. *Ecology Letters*, 19, 487–494.

Isbell, F. & Loreau, M. (2013). Human impacts on minimum subsets of species critical for maintaining ecosystem structure. *Basic and Applied Ecology*, 14, 623–629.

Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*, 30, 1883–1893.

Jordán, F. (2009). Keystone species and food webs. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364, 1733–1741.

Jordán, F., Liu, W.-c. & Davis, A.J. (2006). Topological keystone species: measures of positional importance in food webs. *Oikos*, 112, 535–546.

Jørgensen, S.E., Mejer, H. & Nielsen, S.N. (1998). Ecosystem as self-organizing critical systems. *Ecological Modelling*, 111, 261–268.

Kaiser-Bunbury, C.N. & Blüthgen, N. (2015). Integrating network ecology with applied conservation: a synthesis and guide to implementation. *AoB Plants*, 7, plv076.

Kaiser-Bunbury, C.N., Muff, S., Memmott, J., Müller, C.B. & Caflisch, A. (2010). The robustness of pollination networks to the loss of species and interactions: a quantitative approach incorporating pollinator behaviour. *Ecology Letters*, 13, 442–52.

Kalman, R.E. (1963). Mathematical Description of Linear Dynamical Systems. *Journal of the Society for*

457 *Industrial and Applied Mathematics Series A Control*, 1, 152–192.

458 Lai, S.-M., Liu, W.-C. & Jordan, F. (2012). On the centrality and uniqueness of species from the network
459 perspective. *Biology Letters*, 8, 570–573.

460 Lever, J.J., Nes, E.H. van, Scheffer, M. & Bascompte, J. (2014). The sudden collapse of pollinator
461 communities. *Ecology Letters*, 17, 350–359.

462 Levin, S. (1999). *Fragile Dominion*. Perseus, Cambridge, MA.

463 Liu, Y.-Y. & Barabási, A.-L. (2016). Control principles of complex systems. *Reviews of Modern Physics*,
464 88, 035006.

465 Liu, Y.-Y., Slotine, J.-J. & Barabási, A.-L. (2011). Controllability of complex networks. *Nature*, 473,
466 167–173.

467 Lopezaraiza-Mikel, M.E., Hayes, R.B., Whalley, M.R. & Memmott, J. (2007). The impact of an alien
468 plant on a native plant-pollinator network: An experimental approach. *Ecology Letters*, 10, 539–550.

469 McDonald-Madden, E., Sabbadin, R., Game, E.T., Baxter, P.W.J., Chadès, I. & Possingham, H.P. (2016).
470 Using food-web theory to conserve ecosystems. *Nature Communications*, 7, 10245.

471 Melián, C.J. & Bascompte, J. (2002). Complex networks: two ways to be robust? *Ecology Letters*, 5,
472 705–708.

473 Memmott, J., Waser, N.M. & Price, M.V. (2004). Tolerance of pollination networks to species extinctions.
474 *Proceedings of the Royal Society B*, 271, 2605–2611.

475 Menge, B.A. (1995). Indirect effects in marine rocky intertidal interaction webs: patterns and importance.
476 *Ecological Monographs*, 65, 21–74.

477 Montoya, M., Pimm, S.L. & Sole, R.V. (2006). Ecological networks and their fragility. *Nature*, 442,
478 259–264.

479 Motter, A.E. (2015). Networkcontrology. *Chaos*, 25, 097621.

480 Ne'Eman, G., Jürgens, A., Newstrom-Lloyd, L., Potts, S.G. & Dafni, A. (2010). A framework for comparing
481 pollinator performance: Effectiveness and efficiency. *Biological Reviews*, 85, 435–451.

482 Olesen, J.M., Eskildsen, L.I. & Venkatasamy, S. (2002). Invasion of pollination networks on oceanic
483 islands: Importance of invader complexes and endemic super generalists. *Diversity and Distributions*, 8,
484 181–192.

485 Palacio, R.D., Valderrama-Ardila, C. & Kattan, G.H. (2016). Generalist Species Have a Central Role In a

486 Highly Diverse Plant-Frugivore Network. *Biotropica*, 48, 349–355.

487 Pascual, M. & Dunne, J.A. (Eds.). (2005). *Ecological Networks: Linking Structure to Dynamics in Food*
488 *Webs*. Oxford University Press, New York.

489 Rodewald, A.D., Rohr, R.P., Fortuna, M.A. & Bascompte, J. (2015). Does removal of invasives restore
490 ecological networks? An experimental approach. *Biological Invasions*, 17, 2139.

491 Ruths, J. & Ruths, D. (2014). Control profiles of complex networks. *Science*, 343, 1373–6.

492 Saavedra, S., Stouffer, D.B., Uzzi, B. & Bascompte, J. (2011). Strong contributors to network persistence
493 are the most vulnerable to extinction. *Nature*, 478, 233–235.

494 Scotti, M., Podani, J. & Jordán, F. (2007). Weighting, scale dependence and indirect effects in ecological
495 networks: A comparative study. *Ecological Complexity*, 4, 148–159.

496 Smith, A.M., Reinhardt Adams, C., Wiese, C. & Wilson, S.B. (2016). Re-vegetation with native species
497 does not control the invasive *Ruellia simplex* in a floodplain forest in Florida, USA. *Applied Vegetation*
498 *Science*, 19, 20–30.

499 Solé, R.V. & Montoya, J.M. (2001). Complexity and fragility in ecological networks. *Proceedings of the*
500 *Royal Society B*, 268, 2039–45.

501 Stouffer, D.B., Cirtwill, A.R. & Bascompte, J. (2014). How exotic plants integrate into pollination
502 networks. *Journal of Ecology*, 102, 1442–1450.

503 Stouffer, D.B., Sales-Pardo, M., Sirer, M.I. & Bascompte, J. (2012). Evolutionary Conservation of Species’
504 Roles in Food Webs. *Science*, 335, 1489–1492.

505 Suding, K.N., Gross, K.L. & Houseman, G.R. (2004). Alternative states and positive feedbacks in
506 restoration ecology. *Trends in Ecology and Evolution*, 19, 46–53.

507 Traveset, A., Heleno, R., Chamorro, S., Vargas, P., McMullen, C.K. & Nogales, M. *et al.* (2013). Invaders
508 of pollination networks in the Galápagos Islands : emergence of novel communities. *Proceedings of the*
509 *Royal Society B*, 280, 20123040.

510 Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and species
511 interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351–1363.

512 Tylianakis, J.M., Laliberté, E., Nielsen, A. & Bascompte, J. (2010). Conservation of species interaction
513 networks. *Biological Conservation*, 143, 2270–2279.

514 Tylianakis, J.M., Tscharnkte, T. & Lewis, O.T. (2007). Habitat modification alters the structure of
515 tropical host-parasitoid food webs. *Nature*, 445, 202–205.

516 Vázquez, D.P. & Aizen, M.A. (2004). Asymmetric specialization: a pervasive feature of plant-pollinator

517 interactions. *Ecology*, 85, 1251–1257.

518 Vázquez, D.P., Melián, C.J., Williams, N.M., Blüthgen, N., Krasnov, B.R. & Poulin, R. (2007). Species
519 abundance and asymmetric interaction strength in ecological networks. *Oikos*, 116, 1120–1127.

520 Vázquez, D.P., Morris, W.F. & Jordano, P. (2005). Interaction frequency as a surrogate for the total
521 effect of animal mutualists on plants. *Ecology Letters*, 8, 1088–1094.

522 Vilà, M., Bartomeus, I., Dietzsch, A.C., Petanidou, T., Steffan-Dewenter, I. & Stout, J.C. *et al.* (2009).
523 Invasive plant integration into native plant-pollinator networks across Europe. *Proceedings of the Royal*
524 *Society B*, 276, 3887–3893.

525 West, D. (2001). *Introduction to graph theory*. Prentice Hall, Upper Saddle River.

526 Woodford, D.J., Richardson, D.M., Macisaac, H.J., Mandrak, N.E., Van Wilgen, B.W. & Wilson, J.R.U.
527 *et al.* (2016). Confronting the wicked problem of managing biological invasions. *NeoBiota*, 86, 63–86.

Table legends

- 1 **Factors explaining species importance.** Factor estimates correspond to the average over all models that accounted for 95% of the AIC evidence. Confidence intervals correspond to $\alpha = 0.05$.

Figure legends

- 1 **Matchings of a simple network.** (a) We start with a network in which the direction of the links indicates the potential direction of control; for example a link from a_I to p_I indicates that the state of p_I is influenced by the state of a_I . The numbers indicate the weight of each link. (b & c) This network has two maximal cardinality matchings; that is, two configurations in which it would be possible to exert full control of the network via external input signals to a minimal set of nodes. In both cases, the three matched links (purple arrows) represent the control paths through the network and provide an indication of the matched nodes (purple), which are controlled by superior nodes within the network (circular nodes). Unmatched nodes (orange) are called driver nodes because full network control requires external signals to be applied to them. Out of the two maximal cardinality matchings only one (c) has maximum weight and therefore is also a maximum matching. Further examples can be found in the Supporting Information S1.
- 2 **Different ways to depict quantitative mutualistic networks.** (a) Pollination networks are frequently described by the observed number of visits between each plant and animal species. (b) Based on that visitation data, the mutual dependences between interacting species are calculated directly based on the relative visitation frequencies. (c) The relative differences of these dependences—the interaction asymmetry—then provide a means to estimate the dominant direction of the interspecific effects.
- 3 **Driver-node density.** (a) Histogram of the driver-node density (n_D) for the twenty networks. (b) Invaded communities have lower n_D than uninvaded communities even when controlling for factors related to species richness. The boxes cover the 25th–75th percentiles, the middle lines mark the median, and the maximum length of the whiskers is 1.5 times the interquartile range. (c) Out of the richness metrics, the ratio of plants to pollinators showed a strong, negative relationship with n_D . In both plots, partial residuals correspond to the partial working residuals of the invasion status in our generalised linear mixed model.

- 4 **The driver-node density n_D of empirical networks compared to network randomisations.** For each randomisation approach, we show the standardised rank of the empirical value compared to the set of randomisations. A scaled mean rank—akin to a p-value—less than 0.025 or greater than 0.975 (the areas shaded in light grey) suggests a significant difference between the empirical network and its randomisations. The empirical n_D is much larger than that of network randomisations in which the direction of asymmetries has been randomised. In contrast, the manageability of networks in which the species degrees were randomly shuffled were not significantly different. All boxes are as in Figure 3a.
- 5 **Relationships between f_S and f_D and species structural properties.** (a) In all networks where they were present, invasive species were classified as superior (f_S) and driver (f_D) nodes in all possible control configurations. (b) Species strength (the sum of the dependences of other species on the species of interest) is the single most important factor explaining f_S . Visitation strength and degree also had an important albeit comparatively smaller effect (dashed lines correspond to \pm one standard deviation of these factors). Invasive species are depicted as circles.

Table 1

	imp.	est.	C.I.
f_s			
(Intercept)	1.00	2.69	2.5
species strength	1.00	34.26	15
visitation strength	1.00	1.37	1.1
degree	0.90	4.12	5.5
contribution to nestedness	0.56	0.44	1.3
guild (pollinator)	0.48	0.72	2.6
eigen-centrality	0.25	0.00	0.19
invasive sp.	0.24	-6.23	3.2E+06
f_d			
(Intercept)	1.00	-0.19	0.83
guild (pollinator)	1.00	4.05	0.99
contribution to nestedness	1.00	1.41	0.62
degree	1.00	-5.31	2.5
species strength	1.00	4.65	2.6
visitation strength	1.00	3.07	2.7
eigen-centrality	0.71	0.72	1.5
invasive sp.	0.08	10.95	4.5E+06




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5