

# Reworked analyses and figure drafts for paper1 after JEB reviews

## 1 Distribution of mutations

This is a redraw of the theoretical figure (fig 1) using R code:


### 1.1 Distribution of mutations relative to control using means only.

Here are various figures emphasizing variation among mutant lines compared to controls. In this first figure, the distribution of line means is plotted with the distribution of all control plants as well as 116 resampled means of control replicates equal in size to the average number of reps per SALK line


```
[1] "pdf"
```

This figure compares just the resampled means to the SALK line means

```
[1] "pdf"
```



Here are the number of line means less than control




```
##           Df Sum Sq Mean Sq F value Pr(>F)
## GeneFamilyName 1 8841 8841 10.38 0.00146 **
## Residuals    227 193427 852
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 72 observations deleted due to missingness
##
## Welch Two Sample t-test
##
```


```
## data:  fitdev by GeneFamilyName
## t = 3.263, df = 224.06, p-value = 0.001275
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##    4.930097 19.964495
## sample estimates:
##                mean in group Control
##                -1.935837e-15
## mean in group MADS-box Transcription Factor Family
##                -1.244730e+01
```

Here is the distribution of fitness effects among mads box genes


```
## Warning in (function (x, y, box.ratio = 1, box.width = box.ratio/(1 + box.ratio),
:  NAs introduced by coercion
```



```
##           Df Sum Sq Mean Sq F value    Pr(>F)
## SALK_Line   14  26332   1880.9     2.288 0.00616 **
## Residuals  214 175937    822.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 72 observations deleted due to missingness
```



## 1.2 Fitness components



## 2 Analysis of fitness

Taking all SALK lines and pooling them and comparing to the control:


You can see from the resampled line distributions that salk lines and controls have similar mean fitnesses, with definite differences in variance among groups

```
##
##  Welch Two Sample t-test
##
## data:  fitness by treat
## t = -0.63525, df = 155.85, p-value = 0.5262
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  -8.141375  4.179152
## sample estimates:
## mean in group control    mean in group treat
##           39.39545           41.37657
##
##  Kruskal-Wallis rank sum test
##
## data:  fitmerg$treat and fitmerg$fitness
## Kruskal-Wallis chi-squared = 1106.5, df = 763, p-value = 4.391e-15
```


```
## [1] "permutation typeI prob:"  
## [1] 0.717
```

### 3 Effects of tdna insert number

Here is a plot that relates our total measure of fitness in the lines used in the original pilot study to the ratio of tdna to endogenous genes







```
Call:
lm(formula = fitdev ~ log(area.ratio), data = fruit.by.line)
```

Residuals:

Min	1Q	Median	3Q	Max
-35.920	-19.680	-3.513	14.541	64.439

Coefficients:

Estimate	Std. Error	t value	Pr(> t )

```
(Intercept)      -4.0472      2.8887  -1.401    0.164
log(area.ratio)   0.8829      2.6995   0.327    0.744

Residual standard error: 23.52 on 112 degrees of freedom
(7 observations deleted due to missingness)
Multiple R-squared:  0.0009542, Adjusted R-squared:  -0.007966
F-statistic: 0.107 on 1 and 112 DF,  p-value: 0.7442
```

Clearly no pattern there and a regression confirms.

Just to make sure, I also lumped the ratios into categories and looked for a pattern again:

When focusing on medians, it looks a little bit like there might be some variation across categories

```
summary(aov(fitdev~as.factor(ratiocat),fruit.by.line))

              Df Sum Sq Mean Sq F value Pr(>F)
as.factor(ratiocat)    2    187    93.3   0.168  0.846
Residuals              111  61805   556.8
7 observations deleted due to missingness
```

Alas, no pattern there either. I might suggest that, at the least, we worry about copy number less than other factors when choosing lines for UnPAK projects

## 4 Fitness as a function of gene family size

In the original pilot study we chose genes from different families with differing sizes. Again, there does not seem to be a significant relationship between gene family size and reproductive output in the non-regulatory genes, but there does seem to be a slight pattern in the regulatory genes.

This figure is based on the number of genes in the Gene Family data on tair

This is the same figure but with the means of each line plotted instead of all the reps for each line.

```
Call:
lm(formula = fitdev ~ FamilySize * Regulatory, data = fruit.by.line)

Residuals:
    Min       1Q   Median       3Q      Max
-37.50 -17.46  -1.68   15.80   60.80

Coefficients:
```

```

              Estimate Std. Error t value Pr(>|t|)
(Intercept)    0.031853  10.582719   0.003   0.998
FamilySize     -0.004502   0.038440  -0.117   0.907
Regulatoryyes  -16.152593  12.906961  -1.251   0.213
FamilySize:Regulatoryyes  0.188779   0.112813   1.673   0.097 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 23.2 on 112 degrees of freedom
(5 observations deleted due to missingness)
Multiple R-squared:  0.02922, Adjusted R-squared:  0.003212
F-statistic: 1.124 on 3 and 112 DF,  p-value: 0.3427


Call:
lm(formula = fitdev ~ FamilySize, data = fruit.by.line, subset = Regulatory ==
    "yes")

Residuals:
    Min       1Q   Median       3Q      Max
-34.700 -16.930  -1.639   15.799   60.804

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept) -16.1207     7.3637  -2.189   0.0313 *
FamilySize    0.1843     0.1057   1.743   0.0848 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 23.12 on 86 degrees of freedom
(5 observations deleted due to missingness)
Multiple R-squared:  0.03414, Adjusted R-squared:  0.0229
F-statistic: 3.039 on 1 and 86 DF,  p-value: 0.08484

```



And some analyses on the means of each line's fitness deviation from the ancestor: First OLS ancova. Then regressions for non-regulatory and then regulatory genes

```
summary(aov(fitdev~FamilySize*Regulatory,fruit.by.line))
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
FamilySize	1	306	306.3	0.569	0.452
Regulatory	1	1	0.7	0.001	0.972
FamilySize:Regulatory	1	1507	1507.0	2.800	0.097
Residuals	112	60277	538.2		

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1  
5 observations deleted due to missingness

```
summary(lm(fitdev~FamilySize,subset=Regulatory=="no",fruit.by.line))
```

Call:

```
lm(formula = fitdev ~ FamilySize, data = fruit.by.line, subset = Regulatory ==  
"no")
```

Residuals:

Min	1Q	Median	3Q	Max
-37.504	-18.043	-1.843	16.262	47.859

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.031853	10.701052	0.003	0.998
FamilySize	-0.004502	0.038869	-0.116	0.909

Residual standard error: 23.46 on 26 degrees of freedom

(5 observations deleted due to missingness)

Multiple R-squared: 0.0005156, Adjusted R-squared: -0.03793

F-statistic: 0.01341 on 1 and 26 DF, p-value: 0.9087

```
summary(lm(fitdev~FamilySize,subset=Regulatory=="yes",fruit.by.line))
```

Call:

```
lm(formula = fitdev ~ FamilySize, data = fruit.by.line, subset = Regulatory ==  
    "yes")
```

Residuals:

Min	1Q	Median	3Q	Max
-34.700	-16.930	-1.639	15.799	60.804

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	-16.1207	7.3637	-2.189	0.0313 *
FamilySize	0.1843	0.1057	1.743	0.0848 .

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 23.12 on 86 degrees of freedom

(5 observations deleted due to missingness)

Multiple R-squared: 0.03414, Adjusted R-squared: 0.0229

F-statistic: 3.039 on 1 and 86 DF, p-value: 0.08484

```
summary(lm(log(fitdev+40)~FamilySize,subset=Regulatory=="yes",fruit.by.line))
```

Call:

```
lm(formula = log(fitdev + 40) ~ FamilySize, data = fruit.by.line,  
    subset = Regulatory == "yes")
```

Residuals:

Min	1Q	Median	3Q	Max
-3.7990	-0.5099	0.2162	0.6402	1.2325

```

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  3.02358    0.27102  11.156   <2e-16 ***
FamilySize   0.00439    0.00389   1.128    0.262
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.8509 on 86 degrees of freedom
(5 observations deleted due to missingness)
Multiple R-squared:  0.01459, Adjusted R-squared:  0.00313
F-statistic: 1.273 on 1 and 86 DF, p-value: 0.2623

```

Here is a little more sophisticated analysis of the family size effect

```

library(nlme)

##
## Attaching package: 'nlme'
##
## The following object is masked from 'package:lme4':
##
##      lmList

fm1 <- lme(fixed=fitdev~FamilySize,random=~1|SALK_Line,method="ML",subset=Regulatory=="y")
#plot(fm1)
vfix <- varFixed(~FamilySize)
fm2 <- lme(fixed=fitdev~FamilySize,random=~1|SALK_Line,method="ML",subset=Regulatory=="y")
#plot(resid(fm2,type="pearson")~fitted(fm2))
fm3 <- lme(fixed=fitdev~1,random=~1|SALK_Line,method="ML",subset=Regulatory=="yes",na.action=na.omit)

anova(fm3,fm1,fm2) #looks like fm1 is the best model

##      Model df      AIC      BIC    logLik    Test  L.Ratio p-value
## fm3      1  3 7933.316 7947.407 -3963.658
## fm1      2  4 7931.925 7950.713 -3961.962 1 vs 2 3.390619 0.0656
## fm2      3  4 7985.516 8004.304 -3988.758

anova(fm1)

##      numDF denDF  F-value p-value
## (Intercept)      1    722 1.105524 0.2934
## FamilySize      1     86 3.460323 0.0663

```

```
anova(fm2)

##              numDF denDF  F-value p-value
## (Intercept)      1    722 1.276404  0.2589
## FamilySize       1     86 3.534842  0.0635

anova(fm3)

##              numDF denDF  F-value p-value
## (Intercept)      1    722 1.100184  0.2946
```

Ok, here is the analysis of family size with line as random intercept. Terrible heteroscedasticity, repaired using a fixed variance structure. No signal of family size in the final model.

Now, the joint categories approach: Two tests. The first assumes that the rows and columns are independent, but the expected values come from the marginal totals. The second assumes that the number of fitnesses in each of the four categories is equal.

```
Pearson's Chi-squared test with Yates' continuity correction

data:  tbl
X-squared = 2.3999, df = 1, p-value = 0.1213
[1] 6.025056e-07
```

Here is a test of the change in variance through time:

```
brks=c(0,seq(10,150,10),166)
family.size.cat <- cut(fitmerg$FamilySize,breaks=brks)
sds <- with(fitmerg,tapply(fitdev,family.size.cat,sd))
brkmid <- (brks+(c(brks[-1],166)-brks)/2)[-length(sds)]
bartlett.test(fitmerg$fitdev,family.size.cat)

##
## Bartlett test of homogeneity of variances
##
## data:  fitmerg$fitdev and family.size.cat
## Bartlett's K-squared = 38.058, df = 6, p-value = 1.095e-06

plot(sds~brkmid)
summary(lm(sds~brkmid))

##
```

```
## Call:
## lm(formula = sds ~ brkmid)
##
## Residuals:
##      (0,10]      (20,30]      (40,50]      (50,60]      (60,70]      (90,100]      (130,140]
##      -7.426      -5.119       2.664      11.247      -2.590      14.254      -13.031
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   38.7467     7.4525   5.199  0.00347 **
## brkmid        -0.1638     0.1023  -1.602  0.17017
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 10.91 on 5 degrees of freedom
##      (9 observations deleted due to missingness)
## Multiple R-squared:  0.339, Adjusted R-squared:  0.2069
## F-statistic: 2.565 on 1 and 5 DF,  p-value: 0.1702
```

So not much change in variance, though not a lot of power either.

```
Welch Two Sample t-test

data:  fitdev by Regulatory
t = 1.2283, df = 398.63, p-value = 0.2201
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -1.900829  8.231144
sample estimates:
mean in group no mean in group yes
      4.215966      1.050808
```

## 5 Multiple environment experiment

### 5.1 Figures for the 1st multi-environment experiment

This figure is fitness deviation ignoring germination (we don't have per-sowed seed estimates (in other words, replicated) of germination for the first experiment, just average germination for that line for that germination effort. We do have those data for the second three treatments/time points




```
[1] "SALK_017933C"
[1] "SALK_033462C"
[1] "SALK_038957C"
[1] "SALK_042704C"
[1] "SALK_050488C"
[1] "SALK_054680C"
[1] "SALK_059835C"
[1] "SALK_063722C"
[1] "SALK_094332C"
[1] "SALK_126600C"
[1] "SALK_134535C"
[1] "SALK_150522C"
[1] "CS60000"
```

I'm going to try and address courtney's question about the gene family size of high and low lines

```
## [1] "SALK_Line"          "fitdev"          "gene.family.size"
## [4] "lohi"
```


### Compare family sizes for the high and low lines chosen for exp2



Now here is the figure with straight mean fruit number per line:

```
[1] "SALK_017933C"  
[1] 1  
[1] "SALK_033462C"  
[1] 1  
[1] "SALK_038957C"  
[1] 1  
[1] "SALK_042704C"  
[1] 1
```

```
[1] "SALK_050488C"  
[1] 1  
[1] "SALK_054680C"  
[1] 1  
[1] "SALK_059835C"  
[1] 1  
[1] "SALK_063722C"  
[1] 1  
[1] "SALK_094332C"  
[1] 1  
[1] "SALK_126600C"  
[1] 1  
[1] "SALK_134535C"  
[1] 1  
[1] "SALK_150522C"  
[1] 1  
[1] "CS60000"  
[1] 2
```



### 5.1.1 Copy number

In the following figure line width is proportional to copy number category

## 5.2 Various tests of GxE

```
#MixedEffects
fit1 <- lmer(fitdevng~1+(1|SALK_Line),subset=SALK_Line!="CS60000",data=intresults)
```

```

fit2 <- lmer(fitdevng~treattype+(1|SALK_Line),subset=SALK_Line!="CS60000",data=intresult
fit3 <- lmer(fitdevng~treattype+treattype:SALK_Line+(1|SALK_Line),subset=SALK_Line!="CS6

anova(fit1,fit2,fit3)

refitting model(s) with ML (instead of REML)

Data: intresults
Subset: SALK_Line != "CS60000"
Models:
fit1: fitdevng ~ 1 + (1 | SALK_Line)
fit2: fitdevng ~ treattype + (1 | SALK_Line)
fit3: fitdevng ~ treattype + treattype:SALK_Line + (1 | SALK_Line)
      Df    AIC    BIC logLik deviance Chisq Chi Df Pr(>Chisq)
fit1  3 4643.4 4655.4 -2318.7  4637.4
fit2  6 4589.9 4613.9 -2288.9  4577.9 59.548      3 7.343e-13 ***
fit3 50 4578.4 4778.1 -2239.2  4478.4 99.466     44 3.555e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#fit using OLS
fitaov1 <- aov(fitdevng~treattype*SALK_Line,subset=SALK_Line!="CS60000",data=intresults)
summary(fitaov1)

              Df Sum Sq Mean Sq F value    Pr(>F)
treattype      3  350733   116911   24.814 1.37e-14 ***
SALK_Line     11  191431    17403    3.694 5.42e-05 ***
treattype:SALK_Line 33  329706     9991    2.121 0.000476 ***
Residuals    353 1663151     4711
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
63 observations deleted due to missingness

#fit using ML
fitaov1.glm <-glm(fitdevng~treattype*SALK_Line,subset=SALK_Line!="CS60000",data=intresul
anova(fitaov1.glm,test="Chisq")

Analysis of Deviance Table

Model: gaussian, link: identity

Response: fitdevng

```

Terms added sequentially (first to last)

		Df	Deviance	Resid. Df	Resid. Dev	Pr(>Chi)
NULL				400	2535021	
treattype	3	350733		397	2184288	4.771e-16 ***
SALK_Line	11	191431		386	1992857	2.789e-05 ***
treattype:SALK_Line	33	329706		353	1663151	0.0001816 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

#

*#now look in each environment and ask if there are line differences*

#

```
for (trt in unique(intresults$treattype))
```

```
{
```

```
  fit.line <- glm(fitdevng~1+as.factor(SALK_Line),subset=SALK_Line!="CS60000",data=int
```

```
  fit.intercept <- glm(fitdevng~1,subset=SALK_Line!="CS60000",data=intresults[intresul
```

```
  cat(rep("-",30));cat("\n")
```

```
  print(paste("Effect of including line in environment: ",trt))
```

```
  print(anova(fit.intercept,fit.line,test="Chisq"))
```

```
  cat(rep("-",30));cat("\n")
```

```
}
```

-----

[1] "Effect of including line in environment: nutrient"

Analysis of Deviance Table

Model 1: fitdevng ~ 1

Model 2: fitdevng ~ 1 + as.factor(SALK\_Line)

	Resid. Df	Resid. Dev	Df	Deviance	Pr(>Chi)
1	98	1417594			
2	87	1125616	11	291978	0.02033 *

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

-----

-----

[1] "Effect of including line in environment: control"

Analysis of Deviance Table

Model 1: fitdevng ~ 1

Model 2: fitdevng ~ 1 + as.factor(SALK\_Line)

	Resid. Df	Resid. Dev	Df	Deviance	Pr(>Chi)
--	-----------	------------	----	----------	----------

```

1      101      532531
2       90      439049 11      93482  0.05823 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
- - - - -
- - - - -

[1] "Effect of including line in environment:  highwater"
Analysis of Deviance Table

Model 1: fitdevng ~ 1
Model 2: fitdevng ~ 1 + as.factor(SALK_Line)
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1         95      14671
2         84      11170 11    3500.2 0.005812 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
- - - - -
- - - - -

[1] "Effect of including line in environment:  FIRST.EXP"
Analysis of Deviance Table

Model 1: fitdevng ~ 1
Model 2: fitdevng ~ 1 + as.factor(SALK_Line)
  Resid. Df Resid. Dev Df Deviance  Pr(>Chi)
1        103      219492
2         92      87315 11    132177 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
- - - - -
- - - - -

```

Here is an analysis with Prior information and ecotypes included

```

results$genotype.cls = rep("control",dim(results)[1])
results$genotype.cls[results$fitdev<0] = "low"
results$genotype.cls[results$fitdev>0] = "high"
results$genotype.cls[results$SALK_Line=="CS60000"] = "control"
results$genotype.cls[grepl("ECO",results$SALK_Line)] = "ecotype"

fit1 <- lm(fitness~treattype*genotype.cls,subset=treattype!="FIRST.EXP",data=results)
Anova(fit1,contrasts = list(treattype=contr.sum,genotype.cls=contr.sum),type=3)

Anova Table (Type III tests)

```

Response: fitness

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	110881	1	49.774	7.596e-12 ***
treatype	44786	2	10.052	5.497e-05 ***
genotype.cls	490361	3	73.373	< 2.2e-16 ***
treatype:genotype.cls	330825	6	24.751	< 2.2e-16 ***
Residuals	895536	402		

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

```
fit2 <- lm(fitness~treatype*genotype.cls,subset=(treatype!="FIRST.EXP")&(genotype.cls!=  
Anova(fit2,contrasts = list(treatype=contr.sum,genotype.cls=contr.sum),type=3)
```

Anova Table (Type III tests)

Response: fitness

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	110881	1	54.756	9.043e-13 ***
treatype	44786	2	11.058	2.157e-05 ***
genotype.cls	490177	2	121.031	< 2.2e-16 ***
treatype:genotype.cls	328440	4	40.548	< 2.2e-16 ***
Residuals	759380	375		

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

This analysis just looks for GxE and main effects in the ecotypes. Not much signal  
Ecotype lines only

```
fitaov.genotype.cls.nomut <- aov(fitness~treatype*SALK_Line,subset=(treatype!="FIRST.E  
summary(fitaov.genotype.cls.nomut)
```

##	Df	Sum Sq	Mean Sq	F value	Pr(>F)
## treatype	2	20455	10228	1.714	0.234
## SALK_Line	9	23496	2611	0.438	0.883
## treatype:SALK_Line	9	58971	6552	1.098	0.446
## Residuals	9	53689	5965		

```
fitall <- glm(fitdev~treatype*SALK_Line,subset=treatype!="FIRST.EXP",data=intresults)  
(anova(fitall,test="F"))
```

## Analysis of Deviance Table

##



```
## Model: gaussian, link: identity
##
## Response: fitdev
##
## Terms added sequentially (first to last)
##
##
##              Df Deviance Resid. Df Resid. Dev      F      Pr(>F)
## NULL                      383      3103317
## treattype           2      647029      381      2456288 59.5646 < 2.2e-16 ***
## SALK_Line          12      234750      369      2221538  3.6018 4.360e-05 ***
## treattype:SALK_Line 24      347732      345      1873806  2.6676 5.317e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#add in a classifier for early experiment performance
firstmns <- with(intresults[intresults$treattype=="FIRST.EXP",c("fitdev", "SALK_Line")],
                aggregate(cbind(first.fitdev=fitdev),by=list(SALK_Line=SALK_Line),mean,
intresults <- merge(intresults,firstmns,all.x=T)

fitlo <- glm(fitdev~treattype*SALK_Line,subset=((treattype!="FIRST.EXP")&(first.fitdev<
(anova(fitlo,test="F"))

## Analysis of Deviance Table
##
## Model: gaussian, link: identity
##
## Response: fitdev
##
## Terms added sequentially (first to last)
##
##
##              Df Deviance Resid. Df Resid. Dev      F      Pr(>F)
## NULL                      236      1805730
## treattype           2      278944      234      1526786 25.0251 1.731e-10 ***
## SALK_Line           7      146974      227      1379812  3.7673 0.0007137 ***
## treattype:SALK_Line 14      192704      213      1187108  2.4697 0.0029519 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

fithi <- glm(fitdev~treattype*SALK_Line,subset=((treattype!="FIRST.EXP")&(first.fitdev>
```

```
## Analysis of Deviance Table
##
## Model: gaussian, link: identity
##
## Response: fitdev
##
## Terms added sequentially (first to last)
##
##
##              Df Deviance Resid. Df Resid. Dev      F      Pr(>F)
## NULL                                146      1232611
## treattype          2    428976          144      803635 41.2297 1.228e-14 ***
## SALK_Line          4     18145          140      785490  0.8720  0.48267
## treattype:SALK_Line 8     98791          132      686698  2.3738  0.02021 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

intresults$lohi <- factor(ifelse(intresults$first.fitdev<=0,"low","high"))
fitlowhi <- glm(fitdev~treattype*lohi,subset=(treattype!="FIRST.EXP"),data=intresults)
(anova(fitlowhi,test="F"))

## Analysis of Deviance Table
##
## Model: gaussian, link: identity
##
## Response: fitdev
##
## Terms added sequentially (first to last)
##
##
##              Df Deviance Resid. Df Resid. Dev      F      Pr(>F)
## NULL                                383      3103317
## treattype          2    647029          381      2456288 52.4749 < 2.2e-16 ***
## lohi                1     68562          380      2387726 11.1209 0.0009381 ***
## treattype:lohi      2     57305          378      2330421  4.6475 0.0101397 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Anova(lm(fitdev~treattype*lohi,subset=(treattype!="FIRST.EXP"),data=intresults),contrast

## Anova Table (Type III tests)
##
## Response: fitdev
```

```
##              Sum Sq  Df F value    Pr(>F)
## (Intercept)    23242   1  3.7699  0.05293 .
## treattype     428976   2 34.7905 1.353e-14 ***
## lohi          15911   1  2.5808  0.10900
## treattype:lohi  57305   2  4.6475  0.01014 *
## Residuals     2330421 378
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

fiteco <- glm(fitdev~treattype*SALK_Line,data=intresults.ecotypes)
(anova(fiteco,test="F"))

## Analysis of Deviance Table
##
## Model: gaussian, link: identity
##
## Response: fitdev
##
## Terms added sequentially (first to last)
##
##              Df Deviance Resid. Df Resid. Dev      F Pr(>F)
## NULL                                29      154737
## treattype           2      18581           27      136156 1.5574 0.2625
## SALK_Line           9      23496           18      112660 0.4376 0.8829
## treattype:SALK_Line  9      58971           9       53689 1.0984 0.4456
```

## 5.3 Tables for the MS

### 5.3.1 Gene families

```
famtable <- unique(fitmerg[!is.na(fitmerg$Regulatory),c("GeneFamilyName","FamilySize","R
linesfromfams <- with(unique(fitmerg[fitmerg$GeneFamilyName!="Control",c("SALK_Line","Ge
famtable <- merge(famtable,linesfromfams,all.x=T)
famtable <- famtable[order(famtable$Regulatory,-famtable$FamilySize),]
famtable$Function <- ifelse(famtable$Regulatory=="yes","Regulatory","Metabolic")
famtable <- famtable[,~which(names(famtable)=="Regulatory")]

require(xtable)

## Loading required package: xtable
##
```

```
## Attaching package: 'xtable'
##
## The following objects are masked from 'package:Hmisc':
##
## label, label<-
print(file="redundancy-tables-figs/tbl1-genefams.html",xtable(famtable),type="html",
      include.rownames=F)
```

### 5.3.2 SALK Line list

```
salktbl <- unique(fitmerg[,c("SALK_Line","Gene","Gene_Family")])
salktbl <- salktbl[complete.cases(salktbl),]
salktbl <- salktbl[order(salktbl$Gene_Family,salktbl$SALK_Line),]
print(file="redundancy-tables-figs/tbls1-salk-lines.html",xtable(salktbl),type="html",
      include.rownames=F)
```

## 5.4 Test for effect of line on fitness for first exp.

```
summary(aov(log(fitness+1)~SALK_Line,data=fitmerg[grepl("SALK.[0-9]+C",fitmerg$SALK_Line)

##              Df Sum Sq Mean Sq F value Pr(>F)
## SALK_Line    115  604.2    5.254    2.991 <2e-16 ***
## Residuals    928 1630.4    1.757
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

## 6 Naturally occurring variants

We took the data from Cao et al 2010 and determined how many of the lines in this experiment also showed some sort of natural variation in gene function

```
suppressMessages(require(dplyr))
snp <- unique(read.csv(paste0(csvdir,"/phen-snp.csv"))[,1:2])
names(snp)[2] <- "snp.strains"
snp <- snp %>% group_by(Accession) %>% summarise(snp.strains.mn=sum(snp.strains))
sv <- unique(read.csv(paste0(csvdir,"/phen-sv.csv"))[,1:2])
```

```
names(sv)[2] <- "sv.strains"
sv <- sv %>% group_by(Accession) %>% summarise(sv.strains.mn=sum(sv.strains))
write.table(file="cao-digested.csv", sep=",", row.names=F, unique(merge(snp, sv)))
```

There are definitely lines that are knocked out in nature. The first table is the frequency of lines with no natural variants (false) versus variants for SNPs that should drastically alter gene function. The second is for the distribution of lines with large structural variants

```
with(unique(snp), table(snp.strains.mn>0))
```


```
##
## FALSE  TRUE
##    99    16
```

```
with(unique(sv), table(sv.strains.mn>0))
```


```
##
## FALSE  TRUE
##   106     9
```

The following figure illustrates the distribution of the naturally occurring variants in our

**Distribution of natural accessions with large effect SNPs**




**Distribution of natural accessions with large structural variants**



collection of lines.

## 6.1 Some additional comparisons we have not emphasized

### 6.1.1 Germination rate as a function of gene-family



And here's another test of the same hypothesis using permutation approach (may be less affected by unequal var).

```
[1] "typeI prob:"
[1] 0.017
```

## 6.2 Survival to harvest as a function of gene family

Here are analyses that examine the effect of gene family upon survival:

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
GeneFamilyName	12	0.1829	0.01524	0.758	0.692
Residuals	105	2.1122	0.02012		