



[Course](#) > [Unit 9: Integer Optimization](#) > [Sports Scheduling: An Introduction to Integer Optimization](#) > Quick Question

Quick Question

🔖 Bookmark this page

Quick Question

2/2 points (graded)

Suppose we had two more teams in our tournament (for a total of 6 teams). Each division would have 3 teams. So each team plays TWO teams twice (the teams in their division), and each team plays three teams once (the teams in the other division). This means that the tournament will last for 7 weeks. How many decision variables would we have?

✔ Answer: 105

Explanation

We would have 105 decision variables because we have 7 weeks, and 15 different pairs of teams.


How many constraints would we have? Don't include the constraints that force the variables to be binary when counting the constraints here. (HINT: We would have 6 division constraints, since each pair in each division needs to play twice.)

✔ Answer: 57

Explanation


We would have 6 division constraints, 9 non-division constraints (each of the three teams in one division has to play each of the three teams in the other division), and 42 constraints to make sure each team only plays one team each week (6 teams times 7 weeks).

You have used 2 of 4 attempts

 Your answers have been saved but not graded. Click 'Submit' to grade them.

© All Rights Reserved



 English ▼

© 2012–2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

POWERED BY
OPENedX®

