



[Course](#) > [Unit 4: Trees](#) > [Assignment 4](#) > Understanding Why People Vote

## Understanding Why People Vote

🔖 Bookmark this page

### understanding why people vote

In August 2006 three researchers (Alan Gerber and Donald Green of Yale University, and Christopher Larimer of the University of Northern Iowa) carried out a large scale field experiment in Michigan, USA to test the hypothesis that one of the reasons people vote is social, or extrinsic, pressure. To quote the first paragraph of their [2008 research paper](#):

Among the most striking features of a democratic political system is the participation of millions of voters in elections. Why do large numbers of people vote, despite the fact that ... "the casting of a single vote is of no significance where there is a multitude of electors"? One hypothesis is adherence to social norms. Voting is widely regarded as a citizen duty, and citizens worry that others will think less of them if they fail to participate in elections. Voters' sense of civic duty has long been a leading explanation of vote turnout...

In this homework problem we will use both logistic regression and classification trees to analyze the data they collected.

### The data

The researchers grouped about 344,000 voters into different groups randomly - about 191,000 voters were a "control" group, and the rest were categorized into one of four "treatment" groups. These five groups correspond to five binary variables in the dataset.

1. "Civic Duty" (variable **civicduty**) group members were sent a letter that simply said "DO YOUR CIVIC DUTY - VOTE!"
2. "Hawthorne Effect" (variable **hawthorne**) group members were sent a letter that had the "Civic Duty" message plus the additional message "YOU ARE BEING STUDIED" and they were informed that their voting behavior would be examined by means of public records.

3. "Self" (variable **self**) group members received the "Civic Duty" message as well as the recent voting record of everyone in that household and a message stating that another message would be sent after the election with updated records.
4. "Neighbors" (variable **neighbors**) group members were given the same message as that for the "Self" group, except the message not only had the household voting records but also that of neighbors - maximizing social pressure.
5. "Control" (variable **control**) group members were not sent anything, and represented the typical voting situation.

Additional variables include **sex** (0 for male, 1 for female), **job** (year of birth), and the dependent variable **voting** (1 if they voted, 0 otherwise).

---

## Problem 1.1 - Exploration and Logistic Regression

1/1 point (graded)

We will first get familiar with the data. Load the CSV file gerber.csv into R. What proportion of people in this dataset voted in this election?

0.3159

✓ Answer: 0.316

### Explanation

Load the dataset into R by using the read.csv command:

```
gerber = read.csv("gerber.csv")
```

Then we can compute the percentage of people who voted by using the table function:

```
table(gerber$voting)
```

The output tells us that 235,388 people did not vote, and 108,696 people did vote. This means that  $108696/(108696+235388) = 0.316$  of all people voted in the election.

Submit

You have used 1 of 3 attempts

---

**i** Answers are displayed within the problem

---

## Problem 1.2 - Exploration and Logistic Regression

1/1 point (graded)

Which of the four "treatment groups" had the largest percentage of people who actually voted (voting = 1)?

☐ Civic Duty☐ Hawthorne Effect☐ Self☒ Neighbors ✓

### Explanation

There are several ways to get this answer. One is to use the `tapply` function, and compute the mean value of "voting", sorted by whether or not the people were in each group:

```
tapply(gerber$voting, gerber$civicduty, mean)
```

```
tapply(gerber$voting, gerber$hawthorne, mean)
```

```
tapply(gerber$voting, gerber$self, mean)
```

```
tapply(gerber$voting, gerber$neighbors, mean)
```

The variable with the largest value in the "1" column has the largest fraction of people voting in their group - this is the Neighbors group.

You have used 1 of 2 attempts

 Answers are displayed within the problem

## Problem 1.3 - Exploration and Logistic Regression

1/1 point (graded)

Build a **logistic regression** model for *voting* using the four treatment group variables as the independent variables (civicduty, hawthorne, self, and neighbors). Use all the data to build the model (DO NOT split the data into a training set and testing set). Which of the following coefficients are significant in the logistic regression model? Select all that apply.

☒ Civic Duty ✓☒ Hawthorne Effect ✓☒ Self ✓

☒ Neighbors ✓

### Explanation

You can build the logistic regression model with the following command:

```
LogModel = glm(voting ~ civicduty + hawthorne + self + neighbors, data=gerber, family="binomial")
```

If you look at the output of `summary(LogModel)`, you can see that all of the variables are significant.

You have used 1 of 3 attempts

 Answers are displayed within the problem

## Problem 1.4 - Exploration and Logistic Regression

2/2 points (graded)

Using a threshold of **0.3**, what is the accuracy of the logistic regression model? (When making predictions, you don't need to use the `newdata` argument since we didn't split our data.)

✓ Answer: 0.542

### Explanation

First compute predictions:

```
predictLog = predict(LogModel, type="response")
```

Then, use the table function to make a confusion matrix:

```
table(gerber$voting, predictLog > 0.3)
```

We can compute the accuracy of the sum of the true positives and true negatives, divided by the sum of all numbers in the table:

```
(134513+51966)/(134513+100875+56730+51966) = 0.542
```

You have used 1 of 5 attempts

 Answers are displayed within the problem

## Problem 1.5 - Exploration and Logistic Regression

1/1 point (graded)

Using a threshold of **0.5**, what is the accuracy of the logistic regression model?

0.6841004

✓ Answer: 0.684

### Explanation

First compute predictions:

```
predictLog = predict(LogModel, type="response")
```

Then, use the table function to make a confusion matrix:

```
table(gerber$voting, predictLog > 0.5)
```

We can compute the accuracy of the sum of the true positives and true negatives, divided by the sum of all numbers in the table:

```
(235388+0)/(235388+108696) = 0.684
```

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Problem 1.6 - Exploration and Logistic Regression

1/1 point (graded)

Compare your previous two answers to the percentage of people who did not vote (the baseline accuracy) and compute the AUC of the model. What is happening here?

- ☒ Even though all of the variables are significant, this is a weak predictive model. ✓
- ☐ The model's accuracy doesn't improve over the baseline, but the AUC is high, so this is a strong predictive model.

### Explanation

You can compute the AUC with the following commands (if your model's predictions are called "predictLog"):

```
library(ROCR)
```

```
ROCRpred = prediction(predictLog, gerber$voting)
```

```
as.numeric(performance(ROCRpred, "auc")@y.values)
```

Even though all of our variables are significant, our model does not improve over the baseline model of just predicting that someone will not vote, and the AUC is low. So while the treatment groups do make a difference, this is a weak predictive model.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Problem 2.1 - Trees

1/1 point (graded)

We will now try out trees. Build a CART tree for *voting* using all data and the same four treatment variables we used before. Don't set the option `method="class"` - we are actually going to create a regression tree here. We are interested in building a tree to explore the fraction of people who vote, or the probability of voting. We'd like CART to split our groups if they have different probabilities of voting. If we used `method='class'`, CART would only split if one of the groups had a probability of voting above 50% and the other had a probability of voting less than 50% (since the predicted outcomes would be different). However, with regression trees, CART will split even if both groups have probability less than 50%.

Leave all the parameters at their default values. You can use the following command in R to build the tree:

```
CARTmodel = rpart(voting ~ civicduty + hawthorne + self + neighbors, data=gerber)
```

Plot the tree. What happens, and if relevant, why?

- ☐ Only the "Neighbors" variable is used in the tree - it is the only one with a big enough effect.
- ☐ All variables are used - they all make a difference.
- ☒ No variables are used (the tree is only a root node) - none of the variables make a big enough effect to be split on. ✓

### Explanation

If you plot the tree, with `prp(CARTmodel)`, you should just see one leaf! There are no splits in the tree, because none of the variables make a big enough effect to be split on.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Problem 2.2 - Trees

1/1 point (graded)

Now build the tree using the command:

```
CARTmodel2 = rpart(voting ~ civicduty + hawthorne + self + neighbors, data=gerber,
cp=0.0)
```

to force the complete tree to be built. Then plot the tree. What do you observe about the order of the splits?

☐ Civic duty is the first split, neighbor is the last.

☒ Neighbor is the first split, civic duty is the last. ✓

### Explanation

You can plot the tree with `prp(CARTmodel2)`.

We saw in Problem 1 that the highest fraction of voters was in the Neighbors group, followed by the Self group, followed by the Hawthorne group, and lastly the Civic Duty group. And we see here that the tree detects this trend.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Problem 2.3 - Trees

1/1 point (graded)

Using only the CART tree plot, determine what fraction (a number between 0 and 1) of "Civic Duty" people voted:

.31

✓ Answer: 0.31

### Explanation

You can find this answer by reading the tree - the people in the civic duty group correspond to the bottom right split, which has value 0.31 in the leaf.

Submit

You have used 2 of 3 attempts

**i** Answers are displayed within the problem

## Problem 2.4 - Trees

0/2 points (graded)

Make a new tree that includes the "sex" variable, again with  $cp = 0.0$ . Notice that sex appears as a split that is of secondary importance to the treatment group.

In the control group, which gender is more likely to vote?

☐ Men (0) ✓

☒ Women (1) ✗

In the "Civic Duty" group, which gender is more likely to vote?

☐ Men (0) ✓

☒ Women (1) ✗

### Explanation

You can generate the new tree using the command:

```
CARTmodel3 = rpart(voting ~ civicduty + hawthorne + self + neighbors + sex, data=gerber, cp=0.0)
```

Then, if you plot the tree with `prp(CARTmodel3)`, you can see that there is a split on the "sex" variable after every treatment variable split. For the control group, which corresponds to the bottom left, sex = 0 (male) corresponds to a higher voting percentage.

For the civic duty group, which corresponds to the bottom right, sex = 0 (male) corresponds to a higher voting percentage.

Submit

You have used 1 of 1 attempt



 Answers are displayed within the problem

## Problem 3.1 - Interaction Terms

2/2 points (graded)

We know trees can handle "nonlinear" relationships, e.g. "in the 'Civic Duty' group **and** female", but as we will see in the next few questions, it is possible to do the same for logistic regression. First, let's explore what trees can tell us some more.

Let's just focus on the "Control" treatment group. Create a regression tree using **just the "control" variable**, then create another tree with the "control" and "sex" variables, both with  $cp=0.0$ .

In the "control" only tree, what is the absolute value of the difference in the predicted probability of voting between being in the control group versus being in a different group? You can use the absolute value function to get answer, i.e.  $\text{abs}(\text{Control Prediction} - \text{Non-Control Prediction})$ . Add the argument "digits = 6" to the prp command to get a more accurate estimate.

✓ Answer: 0.043362

### Explanation

You can build the two trees with the following two commands:

```
CARTcontrol = rpart(voting ~ control, data=gerber, cp=0.0)
```

```
CARTsex = rpart(voting ~ control + sex, data=gerber, cp=0.0)
```

Then, plot the "control" tree with the following command:

```
prp(CARTcontrol, digits=6)
```

The split says that if control = 1, predict 0.296638, and if control = 0, predict 0.34. The absolute difference between these is 0.043362.

Submit

You have used 1 of 5 attempts

 Answers are displayed within the problem

## Problem 3.2 - Interaction Terms

1/1 point (graded)

Now, using the second tree (with control and sex), determine who is affected **more** by NOT being in the control group (being in any of the four treatment groups):

- ☐ Men, by a margin of more than 0.001
- ☐ Women, by a margin of more than 0.001
- ☒ They are affected about the same (change in probability within 0.001 of each other).  
✓

### Explanation

You can plot the second tree using the command:

```
prp(CARTsex, digits=6)
```

The first split says that if control = 1, go left. Then, if sex = 1 (female) predict 0.290456, and if sex = 0 (male) predict 0.302795. On the other side of the tree, where control = 0, if sex = 1 (female) predict 0.334176, and if sex = 0 (male) predict 0.345818. So for women, not being in the control group increases the fraction voting by 0.04372. For men, not being in the control group increases the fraction voting by 0.04302. So men and women are affected about the same.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Problem 3.3 - Interaction Terms

1/1 point (graded)

Going back to logistic regression now, create a model using "sex" and "control". Interpret the coefficient for "sex":

- ☒ Coefficient is negative, reflecting that women are less likely to vote ✓
- ☐ Coefficient is negative, reflecting that women are more likely to vote
- ☐ Coefficient is positive, reflecting that women are less likely to vote

- ☐ Coefficient is positive, reflecting that women are more likely to vote

### Explanation

You can create the logistic regression model by using the following command:

```
LogModelSex = glm(voting ~ control + sex, data=gerber, family="binomial")
```

If you look at the summary of the model, you can see that the coefficient for the "sex" variable is -0.055791. This means that women are less likely to vote, since women have a larger value in the sex variable, and a negative coefficient means that larger values are predictive of 0.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Problem 3.4 - Interaction Terms

1/1 point (graded)

The regression tree calculated the percentage voting exactly for every one of the four possibilities (Man, Not Control), (Man, Control), (Woman, Not Control), (Woman, Control). Logistic regression has attempted to do the same, although it wasn't able to do as well because it can't consider exactly the joint possibility of being a women and in the control group.

We can quantify this precisely. Create the following dataframe (this contains all of the possible values of sex and control), and evaluate your logistic regression using the predict function (where "LogModelSex" is the name of your logistic regression model that uses both control and sex):

```
Possibilities = data.frame(sex=c(0,0,1,1),control=c(0,1,0,1))
predict(LogModelSex, newdata=Possibilities, type="response")
```

The four values in the results correspond to the four possibilities in the order they are stated above ( (Man, Not Control), (Man, Control), (Woman, Not Control), (Woman, Control) ). What is the **absolute difference** between the tree and the logistic regression for the **(Woman, Control)** case? Give an answer with five numbers after the decimal point.

0.00035

✓ Answer: 0.00035

### Explanation

The CART tree predicts 0.290456 for the (Woman, Control) case, and the logistic regression model predicts 0.2908065. So the absolute difference, to five decimal places, is 0.00035.

Submit

You have used 2 of 3 attempts

**i** Answers are displayed within the problem

## Problem 3.5 - Interaction Terms

1/1 point (graded)

So the difference is not too big for this dataset, but it is there. We're going to add a new term to our logistic regression now, that is the **combination of the "sex" and "control" variables** - so if this new variable is 1, that means the person is a woman AND in the control group. We can do that with the following command:

```
LogModel2 = glm(voting ~ sex + control + sex:control, data=gerber, family="binomial")
```

How do you interpret the coefficient for the new variable **in isolation**? That is, how does it relate to the dependent variable?

- ☐ If a person is a woman or in the control group, the chance that she voted goes up.
- ☐ If a person is a woman and in the control group, the chance that she voted goes up.
- ☐ If a person is a woman or in the control group, the chance that she voted goes down.
- ☒ If a person is a woman and in the control group, the chance that she voted goes down. ✓

### Explanation

This coefficient is negative, so that means that a value of 1 in this variable decreases the chance of voting. This variable will have variable 1 if the person is a woman and in the control group.

Submit

You have used 1 of 2 attempts

**i** Answers are displayed within the problem

## Problem 3.6 - Interaction Terms

1/1 point (graded)

Run the same code as before to calculate the average for each group:

```
predict(LogModel2, newdata=Possibilities, type="response")
```

Now what is the difference between the logistic regression model and the CART model for the (Woman, Control) case? Again, give your answer with five numbers after the decimal point.

✓ Answer: 0

### Explanation

The logistic regression model now predicts 0.2904558 for the (Woman, Control) case, so there is now a very small difference (practically zero) between CART and logistic regression.

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Problem 3.7 - Interaction Terms

1/1 point (graded)

This example has shown that trees can capture nonlinear relationships that logistic regression can not, but that we can get around this sometimes by using variables that are the combination of two variables. Should we always include all possible interaction terms of the independent variables when building a logistic regression model?

☐ Yes

☒ No ✓

### Explanation

We should not use all possible interaction terms in a logistic regression model due to overfitting. Even in this simple problem, we have four treatment groups and two values for sex. If we have an interaction term for every treatment variable with sex, we will double the number of variables. In smaller data sets, this could quickly lead to overfitting.

[Submit](#)

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

Please remember not to ask for or post complete answers to homework questions in this discussion forum.

## Discussion

[Hide Discussion](#)

**Topic:** Unit 4 / Unit 4, Homework: Understanding Why People Vote

[Add a Post](#)

### ◀ All Posts

#### Question 2.4

question posted 5 days ago by **icephoenix1802**

In problem 2.4, we are asked: "In the control group, which gender is more likely to vote?" However, the model we build does not include 'control' variable. Can someone clarify this for me?

This post is visible to everyone.

[Add a Response](#)

4 responses

**huynh057**

5 days ago

I just added the control variable to the rpart equation. It might be an error, it might not.

**rahensle**

4 days ago



Hi,

I think this was an error in the problems. Neither the model in 2.2 or in the explanation in 2.4 includes the control group. However, the question and context of the explanation in 2.4 imply that the control group should have been included. In addition, when you graph this without the control group, there are no control group splits. Thus, the civic duty = yes split to female & male and the civic duty = no split to female and male could be in either control = yes or control = no since this would be unknown.

Staff, can you check if this was an error please? This can be confusing to others. Were we supposed to include the control group in the previous models or were we supposed to add both sex and control for just 2.4?

Thank you,

Rebecca

I remembered answering this in the thread you made, but you seem to have deleted it.



It's not an error. Each treatment variable is split at 0.5. Less than 0.5 means not belonging to the corresponding group and more than 0.5 means belong to that group.

By that reasoning, you trace the far left path, where every leaf answers **yes** to "**less than 0.5**". Each participant can only belong to 1 group. Hence the leaf at the bottom left corner is the control group.

posted 4 days ago by **lethutrang101**

I don't get how you can infer a group that is not in the model (2.2 and the model in 2.4)...that doesn't seem to fit with basic scientific and data validity thinking. When I ran the model with control in it, my graph matches the explanation where they discuss what is at the bottom left and bottom right. Run the model with control vs. not control and graph prp using type 4 which labels the branches. It's easier to see the trace. From what I've found so far, the model should have included control...and I'm not the only one given the original question and the response.



Unfortunately, it's really hard to discuss this in as much detail as I'd like without violating the discussion rules. That's also why I haven't posted pictures of my trees either which would be a lot clearer. It also would have been a lot easier if their explanation actually included the graph.

Consequently, I'd really prefer to see what the staff says. They're the only ones who can say for sure if they made a mistake or not.

Thank you,

Rebecca

posted 3 days ago by **rahensle**

Add a comment

**msi888**

a day ago



I guess that, if You can generate the new using the command: `CARTmodelX = rpart(voting ~ ....+ ....+ sex, data=gerber, cp=0.0)`, i.e. **without** "control" when you plot the tree you can see that the control group corresponds to the bottom left. It's what remains after all other possibilities have been calculated. The group members who were not sent anything m

Add a comment

**sebastien martin**

a day ago



Hello icephoenix1802,



The control group is by definition people who are not civic duty, hawthorn, self, or neighbors. So the a tree that does not use the control variable can still be used to have information about the control group:

Just look for the leaf that corresponds to people that are not civic duty, hawthorn, self, or neighbors.

Best,

Sebastien (TA)

Add a comment

**strgeon**

32 minutes ago



I had the same question about the control group but when I saw the answer, I realized the control group would come if all others were false. However, in my model, the opposite gender had a higher percentage, I don't understand why.

Add a comment

Preview

Submit

Showing all responses

Add a response:

Preview

Submit

© All Rights Reserved



English ▼

© 2012–2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

POWERED BY  
OPENedX®

