



[Course](#) > [Unit 6: Clustering](#) > [Assignment 6](#) > Predicting Stock Returns with Cluster-Then-Predict

Predicting Stock Returns with Cluster-Then-Predict

🔖 Bookmark this page

Predicting Stock Returns with Cluster-Then-Predict

In the second lecture sequence this week, we heard about cluster-then-predict, a methodology in which you first cluster observations and then build cluster-specific prediction models. In the lecture sequence, we saw how this methodology helped improve the prediction of heart attack risk. In this assignment, we'll use cluster-then-predict to predict future stock prices using historical stock data.

When selecting which stocks to invest in, investors seek to obtain good future returns. In this problem, we will first use clustering to identify clusters of stocks that have similar returns over time. Then, we'll use logistic regression to predict whether or not the stocks will have positive future returns.

For this problem, we'll use [StocksCluster.csv](#), which contains monthly stock returns from the NASDAQ stock exchange. The NASDAQ is the second-largest stock exchange in the world, and it lists many technology companies. The stock price data used in this problem was obtained from [infochimps](#), a website providing access to many datasets.

Each observation in the dataset is the monthly returns of a particular company in a particular year. The years included are 2000-2009. The companies are limited to tickers that were listed on the exchange for the entire period 2000-2009, and whose stock price never fell below \$1. So, for example, one observation is for Yahoo in 2000, and another observation is for Yahoo in 2001. Our goal will be to predict whether or not the stock return in December will be positive, using the stock returns for the first 11 months of the year.

This dataset contains the following variables:

- **ReturnJan** = the return for the company's stock during January (in the year of the observation).

- **ReturnFeb** = the return for the company's stock during February (in the year of the observation).
- **ReturnMar** = the return for the company's stock during March (in the year of the observation).
- **ReturnApr** = the return for the company's stock during April (in the year of the observation).
- **ReturnMay** = the return for the company's stock during May (in the year of the observation).
- **ReturnJune** = the return for the company's stock during June (in the year of the observation).
- **ReturnJuly** = the return for the company's stock during July (in the year of the observation).
- **ReturnAug** = the return for the company's stock during August (in the year of the observation).
- **ReturnSep** = the return for the company's stock during September (in the year of the observation).
- **ReturnOct** = the return for the company's stock during October (in the year of the observation).
- **ReturnNov** = the return for the company's stock during November (in the year of the observation).
- **PositiveDec** = whether or not the company's stock had a positive return in December (in the year of the observation). This variable takes value 1 if the return was positive, and value 0 if the return was not positive.

For the first 11 variables, the value stored is a proportional change in stock value during that month. For instance, a value of 0.05 means the stock increased in value 5% during the month, while a value of -0.02 means the stock decreased in value 2% during the month.

Problem 1.1 - Exploring the Dataset

1/1 point (graded)

Load StocksCluster.csv into a data frame called "stocks". How many observations are in the dataset?

✓ Answer: 11580

Explanation

You can load the dataset with the `read.csv` function:

```
stocks = read.csv("StocksCluster.csv")
```

and see how many observations are included with either the `str` or `nrow` function:

```
str(stocks)
```

```
nrow(stocks)
```

Both tell us that there are 11580 observations in this dataset.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 1.2 - Exploring the Dataset

1/1 point (graded)

What proportion of the observations have positive returns in December?

0.546114

✓ Answer: 0.546114

Explanation

You can compute the proportion of observations with positive returns by using the `table` function:

```
table(stocks$PositiveDec)
```

It tells us that 6324 observations have `PositiveDec = 1`, so $6324/11580 = 0.546$ of the observations have positive returns in December.

Alternatively, you could use the `mean` function to compute the proportion:

```
mean(stocks$PositiveDec)
```

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 1.3 - Exploring the Dataset

1/1 point (graded)

What is the maximum correlation between any two return variables in the dataset? You should look at the pairwise correlations between `ReturnJan`, `ReturnFeb`, `ReturnMar`, `ReturnApr`, `ReturnMay`, `ReturnJune`, `ReturnJuly`, `ReturnAug`, `ReturnSep`, `ReturnOct`, and `ReturnNov`.

✔ Answer: 0.19167279

Explanation

From `cor(stocks)`, we see the largest correlation coefficient is 0.19167279, between ReturnOct and ReturnNov.

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 1.4 - Exploring the Dataset

2/2 points (graded)

Which month (from January through November) has the largest mean return across all observations in the dataset?

✔ Answer: April

Which month (from January through November) has the smallest mean return across all observations in the dataset?

✔ Answer: September

Explanation

These can be determined using the summary function:

```
summary(stocks)
```

If you look at the mean value for each variable, you can see that April has the largest mean value (0.026308), and September has the smallest mean value (-0.014721).

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 2.1 - Initial Logistic Regression Model

2/2 points (graded)

Run the following commands to split the data into a training set and testing set, putting 70% of the data in the training set and 30% of the data in the testing set:

```
set.seed(144)
```

```
spl = sample.split(stocks$PositiveDec, SplitRatio = 0.7)
```

```
stocksTrain = subset(stocks, spl == TRUE)
```

```
stocksTest = subset(stocks, spl == FALSE)
```

Then, use the stocksTrain data frame to train a logistic regression model (name it StocksModel) to predict PositiveDec using all the other variables as independent variables. Don't forget to add the argument family=binomial to your glm command.

What is the overall accuracy on the training set, using a threshold of 0.5?

✓ Answer: 0.5711818

Explanation

We can train the model with:

```
StocksModel = glm(PositiveDec ~ ., data=stocksTrain, family=binomial)
```

Then, we can compute our predictions on the training set with:

```
PredictTrain = predict(StocksModel, type="response")
```

And construct a classification matrix with the table function:

```
table(stocksTrain$PositiveDec, PredictTrain > 0.5)
```

The overall accuracy of the model is $(990 + 3640)/(990 + 2689 + 787 + 3640) = 0.571$.

You have used 1 of 5 attempts

i Answers are displayed within the problem

Problem 2.2 - Initial Logistic Regression Model

1/1 point (graded)

Now obtain test set predictions from StocksModel. What is the overall accuracy of the model on the test, again using a threshold of 0.5?

✓ Answer: 0.5670697

Explanation

You can compute predictions on the test set using the predict function:

```
PredictTest = predict(StocksModel, newdata=stocksTest, type="response")
```

Then, you can compute the classification matrix on the test set with the table function:

```
table(stocksTest$PositiveDec, PredictTest > 0.5)
```

The overall accuracy of the model on the test set is $(417 + 1553)/(417 + 1160 + 344 + 1553) = 0.567$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 2.3 - Initial Logistic Regression Model

1/1 point (graded)

What is the accuracy on the test set of a baseline model that always predicts the most common outcome (PositiveDec = 1)?

0.546056419

✓ Answer: 0.5460564

Explanation

This can be computed by making a table of the outcome variable in the test set:

```
table(stocksTest$PositiveDec)
```

The baseline model would get all of the PositiveDec = 1 cases correct, and all of the PositiveDec = 0 cases wrong, for an accuracy of $1897/(1577 + 1897) = 0.5460564$.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 3.1 - Clustering Stocks

1/1 point (graded)

Now, let's cluster the stocks. The first step in this process is to remove the dependent variable using the following commands:

```
limitedTrain = stocksTrain
```

```
limitedTrain$PositiveDec = NULL
```

```
limitedTest = stocksTest
```

```
limitedTest$PositiveDec = NULL
```

Why do we need to remove the dependent variable in the clustering phase of the cluster-then-predict methodology?

- ☐ Leaving in the dependent variable might lead to unbalanced clusters
- ☐ Removing the dependent variable decreases the computational effort needed to cluster
- ☒ Needing to know the dependent variable value to assign an observation to a cluster defeats the purpose of the methodology ✓

Explanation

In cluster-then-predict, our final goal is to predict the dependent variable, which is unknown to us at the time of prediction. Therefore, if we need to know the outcome value to perform the clustering, the methodology is no longer useful for prediction of an unknown outcome value.

This is an important point that is sometimes mistakenly overlooked. If you use the outcome value to cluster, you might conclude your method strongly outperforms a non-clustering alternative. However, this is because it is using the outcome to determine the clusters, which is not valid.

Submit

You have used 1 of 1 attempt

i Answers are displayed within the problem

Problem 3.2 - Clustering Stocks

2/2 points (graded)

In the market segmentation assignment in this week's homework, you were introduced to the `preProcess` command from the `caret` package, which normalizes variables by subtracting by the mean and dividing by the standard deviation.

In cases where we have a training and testing set, we'll want to normalize by the mean and standard deviation of the variables in the training set. We can do this by passing just the training set to the `preProcess` function:

```
library(caret)
```

```
preproc = preProcess(limitedTrain)
```

```
normTrain = predict(preproc, limitedTrain)
```

```
normTest = predict(preproc, limitedTest)
```

What is the mean of the ReturnJan variable in normTrain?

✓ Answer: 2.100586e-17

What is the mean of the ReturnJan variable in normTest?

✓ Answer: -0.0004185886

Explanation

After running the provided normalization commands, we can read the means with `mean(normTrain$ReturnJan)` and `mean(normTest$ReturnJan)`.

You have used 1 of 3 attempts

 Answers are displayed within the problem

Problem 3.3 - Clustering Stocks

1/1 point (graded)

Why is the mean ReturnJan variable much closer to 0 in normTrain than in normTest?

- ☐ Small rounding errors exist in the normalization procedure
- ☒ The distribution of the ReturnJan variable is different in the training and testing set
✓
- ☐ The distribution of the dependent variable is different in the training and testing set

Explanation

From `mean(stocksTrain$ReturnJan)` and `mean(stocksTest$ReturnJan)`, we see that the average return in January is slightly higher in the training set than in the testing set. Since `normTest` was constructed by subtracting by the mean `ReturnJan` value from the training set, this explains why the mean value of `ReturnJan` is slightly negative in `normTest`.

You have used 1 of 1 attempt

 Answers are displayed within the problem

Problem 3.4 - Clustering Stocks

1/1 point (graded)

Set the random seed to 144 (it is important to do this again, even though we did it earlier). Run k-means clustering with 3 clusters on `normTrain`, storing the result in an object called `km`.

Which cluster has the largest number of observations?

☐ Cluster 1☒ Cluster 2 ✓☐ Cluster 3

Explanation

We can set the seed and run the k-means algorithm with:

```
set.seed(144)
```

```
km = kmeans(normTrain, centers = 3)
```

From `table(km$cluster)`, we can see that cluster 2 has the largest number of observations. Alternatively, you can see the number of observations in each cluster by typing `km$size` in your console.

You have used 1 of 1 attempt

 Answers are displayed within the problem

Problem 3.5 - Clustering Stocks

1/1 point (graded)

Recall from the recitation that we can use the flexclust package to obtain training set and testing set cluster assignments for our observations (note that the call to `as.kcca` may take a while to complete):

```
library(flexclust)
```

```
km.kcca = as.kcca(km, normTrain)
```

```
clusterTrain = predict(km.kcca)
```

```
clusterTest = predict(km.kcca, newdata=normTest)
```

How many test-set observations were assigned to Cluster 2?

✓ Answer: 2080

Explanation

After running the provided commands, we can obtain the breakdown of the testing set clusters with `table(clusterTest)`.

Submit

You have used 1 of 3 attempts

📘 Answers are displayed within the problem

Problem 4.1 - Cluster-Specific Predictions

1/1 point (graded)

Using the subset function, build data frames `stocksTrain1`, `stocksTrain2`, and `stocksTrain3`, containing the elements in the `stocksTrain` data frame assigned to clusters 1, 2, and 3, respectively (be careful to take subsets of `stocksTrain`, not of `normTrain`). Similarly build `stocksTest1`, `stocksTest2`, and `stocksTest3` from the `stocksTest` data frame.

Which training set data frame has the highest average value of the dependent variable?

☒ stocksTrain1 ✓

☐ stocksTrain2☐ stocksTrain3

Explanation

We can obtain the necessary subsets with:

```
stocksTrain1 = subset(stocksTrain, clusterTrain == 1)
```

```
stocksTrain2 = subset(stocksTrain, clusterTrain == 2)
```

```
stocksTrain3 = subset(stocksTrain, clusterTrain == 3)
```


```
stocksTest1 = subset(stocksTest, clusterTest == 1)
```

```
stocksTest2 = subset(stocksTest, clusterTest == 2)
```

```
stocksTest3 = subset(stocksTest, clusterTest == 3)
```

From `mean(stocksTrain1$PositiveDec)`, `mean(stocksTrain2$PositiveDec)`, and `mean(stocksTrain3$PositiveDec)`, we see that `stocksTrain1` has the observations with the highest average value of the dependent variable.

You have used 1 of 1 attempt

 Answers are displayed within the problem

Problem 4.2 - Cluster-Specific Predictions

2/2 points (graded)

Build logistic regression models `StocksModel1`, `StocksModel2`, and `StocksModel3`, which predict `PositiveDec` using all the other variables as independent variables. `StocksModel1` should be trained on `stocksTrain1`, `StocksModel2` should be trained on `stocksTrain2`, and `StocksModel3` should be trained on `stocksTrain3`.

Which variables have a positive sign for the coefficient in at least one of `StocksModel1`, `StocksModel2`, and `StocksModel3` and a negative sign for the coefficient in at least one of `StocksModel1`, `StocksModel2`, and `StocksModel3`? Select all that apply.

☒ ReturnJan ✓☒ ReturnFeb ✓☒ ReturnMar ✓

☐ ReturnApr☐ ReturnMay☒ ReturnJune ✓☐ ReturnJuly☒ ReturnAug ✓☐ ReturnSep☒ ReturnOct ✓☐ ReturnNov

Explanation

We can build the models with:

```
StocksModel1 = glm(PositiveDec ~ ., data=stocksTrain1, family=binomial)
```

```
StocksModel2 = glm(PositiveDec ~ ., data=stocksTrain2, family=binomial)
```

```
StocksModel3 = glm(PositiveDec ~ ., data=stocksTrain3, family=binomial)
```

From `summary(StocksModel1)`, `summary(StocksModel2)`, and `summary(StocksModel3)`, ReturnJan, ReturnFeb, ReturnMar, ReturnJune, ReturnAug, and ReturnOct differ in sign between the models.

You have used 1 of 3 attempts

i Answers are displayed within the problem

Problem 4.3 - Cluster-Specific Predictions

6/6 points (graded)

Using `StocksModel1`, make test-set predictions called `PredictTest1` on the data frame `stocksTest1`. Using `StocksModel2`, make test-set predictions called `PredictTest2` on the data frame `stocksTest2`. Using `StocksModel3`, make test-set predictions called `PredictTest3` on the data frame `stocksTest3`.

What is the overall accuracy of StocksModel1 on the test set stocksTest1, using a threshold of 0.5?

✓ Answer: 0.6194145

What is the overall accuracy of StocksModel2 on the test set stocksTest2, using a threshold of 0.5?

✓ Answer: 0.5504808

What is the overall accuracy of StocksModel3 on the test set stocksTest3, using a threshold of 0.5?

✓ Answer: 0.6458333

Explanation

The predictions can be obtained with:

```
PredictTest1 = predict(StocksModel1, newdata = stocksTest1, type="response")
```

```
PredictTest2 = predict(StocksModel2, newdata = stocksTest2, type="response")
```

```
PredictTest3 = predict(StocksModel3, newdata = stocksTest3, type="response")
```

And the classification matrices can be computed with:

```
table(stocksTest1$PositiveDec, PredictTest1 > 0.5)
```

```
table(stocksTest2$PositiveDec, PredictTest2 > 0.5)
```

```
table(stocksTest3$PositiveDec, PredictTest3 > 0.5)
```

The overall accuracy of StocksModel1 is $(30 + 774)/(30 + 471 + 23 + 774) = 0.6194145$, the overall accuracy of StocksModel2 is $(388 + 757)/(388 + 626 + 309 + 757) = 0.5504808$, and the overall accuracy of StocksModel3 is $(49 + 13)/(49 + 13 + 21 + 13) = 0.6458333$.

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

Problem 4.4 - Cluster-Specific Predictions

1/1 point (graded)

To compute the overall test-set accuracy of the cluster-then-predict approach, we can combine all the test-set predictions into a single vector and all the true outcomes into a single vector:

```
AllPredictions = c(PredictTest1, PredictTest2, PredictTest3)
```

```
AllOutcomes = c(stocksTest1$PositiveDec, stocksTest2$PositiveDec,  
stocksTest3$PositiveDec)
```

What is the overall test-set accuracy of the cluster-then-predict approach, again using a threshold of 0.5?

✓ Answer: 0.5788716

Explanation

After combining the predictions and outcomes with the provided code, we can compute the overall test-set accuracy by creating a classification matrix:

```
table(AllOutcomes, AllPredictions > 0.5)
```

Which tells us that the overall accuracy is $(467 + 1544)/(467 + 1110 + 353 + 1544) = 0.5788716$.

We see a modest improvement over the original logistic regression model. Since predicting stock returns is a notoriously hard problem, this is a good increase in accuracy. By investing in stocks for which we are more confident that they will have positive returns (by selecting the ones with higher predicted probabilities), this cluster-then-predict model can give us an edge over the original logistic regression model.

You have used 1 of 3 attempts


i Answers are displayed within the problem

Please remember not to ask for or post complete answers to homework questions in this discussion forum.

Discussion

Topic: Unit 6 / Unit 6, Homework: Predicting Stock Returns with Cluster-Then-Predict

© All Rights Reserved



© 2012–2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

