

Easy Level Activity Packet

Label Reading, Time Matching, and Basic Safety

Use this standalone packet when the session goal is basic medication label reading, matching medication directions to daily time slots, filling a simple one-day organizer, and identifying common medication safety mistakes.

Activity 1

Match Medication to Time of Day

Activity 2

Fill a One-Day Pill Organizer

Activity 3

Identify Medication Safety Mistakes

MEDICATION BOTTLE BANK USED IN EASY LEVEL

M1 Metformin 500 mg	Take 1 tablet twice daily with meals.
M2 Lisinopril 10 mg	Take 1 tablet every morning.
M3 Atorvastatin 20 mg	Take 1 tablet at bedtime.
M4 Aspirin 81 mg	Take 1 tablet every morning.
M5 Amoxicillin 500 mg	Take 1 capsule every 12 hours until finished.
M6 Gabapentin 300 mg	Take 1 capsule three times daily.
M7 Clopidogrel 75 mg	Take 1 tablet once daily.
M8 Melatonin 5 mg	Take 1 tablet at bedtime as needed.

Clinical simulation resource only

Do not use this packet to make real medication changes. Real medication questions should go to a prescriber or pharmacist.

Activity 1: Match Medication to Time of Day

EASY LEVEL

The patient reads four medication labels and matches each medication to the correct time slot.

SAY "We are going to match each medication to the time of day it should be taken. Read the label, then place it in the right box."

M1 Metformin 500 mg
Take 1 tablet twice daily with meals.

M3 Atorvastatin 20 mg
Take 1 tablet at bedtime.

M4 Aspirin 81 mg
Take 1 tablet every morning.

M8 Melatonin 5 mg
Take 1 tablet at bedtime as needed.

SETUP

Cards
None

Layout
4 time boxes: Morning, Lunch, Dinner, Bedtime

PATIENT WORK AREA

Morning

Lunch

Dinner

Bedtime

The patient fills a four-compartment pill organizer using three unique medication bottles.

SAY "Each section is for a different time of day. Put the right medications in the right sections."

M2 Lisinopril 10 mg
Take 1 tablet every morning.

M6 Gabapentin 300 mg
Take 1 capsule three times daily.

M8 Melatonin 5 mg
Take 1 tablet at bedtime as needed.

SETUP

Cards
PRN card optional

Layout
One-day organizer with Morning, Afternoon, Evening, Bedtime

PATIENT WORK AREA

Morning

Afternoon

Evening

Bedtime

Activity 3: Identify Medication Safety Mistakes

EASY LEVEL

The patient hears short medication behavior statements and decides whether each one is safe or unsafe.

SAY "I am going to read things people do with medications. After each one, tell me safe or unsafe."

M2 Lisinopril 10 mg

Take 1 tablet every morning.

M5 Amoxicillin 500 mg

Take 1 capsule every 12 hours until finished.

M6 Gabapentin 300 mg

Take 1 capsule three times daily.

M7 Clopidogrel 75 mg

Take 1 tablet once daily.

SETUP

Cards

Safety statements

Layout

Safe / unsafe response sheet

PATIENT WORK AREA

Statement	Safe	Unsafe
1. Take Lisinopril exactly as prescribed.	☐	☐
2. Skip Lisinopril if you feel fine today.	☐	☐
3. Call the pharmacy if Amoxicillin directions are unclear.	☐	☐
4. Use someone else's Gabapentin because it helped them.	☐	☐
5. Take double the Gabapentin dose to catch up.	☐	☐
6. Store medications away from children and heat.	☐	☐
7. Stop Amoxicillin early because you feel better.	☐	☐
8. Keep a written medication list for doctor visits.	☐	☐

ACTIVITY 1: MATCH MEDICATION TO TIME OF DAY

The patient reads four medication labels and matches each medication to the correct time slot.

SAY "We are going to match each medication to the time of day it should be taken. Read the label, then place it in the right box."

WHAT YOU NEED

- This printed packet
- Four bottles from the 8-bottle medication bank
- A quiet room with no distractions
- 5 to 10 minutes

STEP-BY-STEP THERAPIST GUIDE

- Orient the patient to Morning, Lunch, Dinner, and Bedtime boxes.
- Work through medications one at a time and ask the patient to read each label aloud.
- Let the patient self-check by reading each slot back before you review.
- Confirm correct answers and double-check errors by returning to the label.

ANSWER KEY

Morning	Metformin 500 mg + Aspirin 81 mg
Lunch	Empty - no medication scheduled
Dinner	Metformin 500 mg
Bedtime	Atorvastatin 20 mg + Melatonin 5 mg as needed

WATCH FOR

- Atorvastatin placed in morning instead of bedtime
- Only one morning medication placed
- Selection by pill color instead of label reading
- Metformin missed as a twice-daily medication

IF THEY STRUGGLE

- Read the last line on this bottle. What time does it say?
- Count how many labels say morning. Let us count together.
- Ignore the colors. Just read the words on the label.

Great work. You read every label before placing anything. That is exactly the habit that keeps you safe at home.

ACTIVITY 2: FILL A ONE-DAY PILL ORGANIZER

The patient fills a four-compartment pill organizer using three unique medication bottles.

SAY "Each section is for a different time of day. Put the right medications in the right sections."

WHAT YOU NEED

- This printed packet or a real 4-slot pill organizer
- Three unique bottles from the medication bank
- Pen and paper if completing on paper
- About 10 minutes

STEP-BY-STEP THERAPIST GUIDE

- Show the pill organizer and read the four compartments aloud.
- Introduce medications one at a time and ask the patient to read the name and directions.
- Pause after each placement and ask the patient to re-check the label.
- Ask why the empty slot is okay.
- Use teach-back: have the patient explain the completed organizer.

ANSWER KEY

Morning	Lisinopril 10 mg + Gabapentin 300 mg
Afternoon	Gabapentin 300 mg
Evening	Gabapentin 300 mg
Bedtime	Melatonin 5 mg as needed

WATCH FOR

- Melatonin placed in Evening instead of Bedtime
- Gabapentin not placed three times across the day
- Patient chooses by color before reading
- Patient rushes without reading each label

IF THEY STRUGGLE

- Does the label say evening or bedtime? Point to the exact word.
- Does any label say afternoon?
- Pretend all pills are the same color. Now where does the label say it goes?

You filled a real pill organizer correctly by reading every label. This is exactly what you will do at home each week.

ACTIVITY 3: IDENTIFY MEDICATION SAFETY MISTAKES

The patient hears short medication behavior statements and decides whether each one is safe or unsafe.

SAY "I am going to read things people do with medications. After each one, tell me safe or unsafe."

WHAT YOU NEED

- This printed packet
- Read each statement aloud
- Patient responds verbally or by pointing
- About 10 to 15 minutes

STEP-BY-STEP THERAPIST GUIDE

- Explain that the patient only needs to answer safe or unsafe.
- Read each statement slowly and pause before prompting.
- Ask the patient to explain each answer before correcting.
- Return to missed items at the end for discussion.

ANSWER KEY

Safe	Take Lisinopril exactly as prescribed; call pharmacy if Amoxicillin directions are unclear; store medications safely; keep a written medication list.
Unsafe	Skip Lisinopril if you feel fine; use someone else's Gabapentin; double-dose Gabapentin; stop Amoxicillin early.

WATCH FOR

- Skipping blood pressure medication viewed as safe
- Stopping antibiotics early when feeling better
- Fast answers without reasoning
- Double-dosing treated as not serious

IF THEY STRUGGLE

- What if blood pressure is high but there is no headache?
- What does your body experience when it suddenly gets twice the medication?
- What happens to bacteria that are still there when you stop early?

You reviewed medication safety rules that protect you every day at home. Well done.