

# TITLE

AIRS Science Team Meeting

---

L. Larrabee Strow<sup>1,2</sup>, Sergio DeSouza-Machado<sup>1,2</sup>, Steven Leroy<sup>3</sup>, Howard Motteler<sup>2</sup>, Chris Hepplewhite<sup>2</sup>, and Steven Buczkowski<sup>2</sup>

October 3, 2018

<sup>1</sup>UMBC Physics Dept.

<sup>2</sup>UMBC JCET

<sup>3</sup>AER

## SAMPLE: Motivation

- Produce Level 1b CHIRP radiances for retrievals
- Produce Level 3 climate-level gridded CHIRP radiance products
- Goals
  - Minimize sensitivity to a-priori estimates, etc.
  - Remove artificial sampling biases
  - Perform as much analysis in radiance space for error traceability
- Geophysical Products
  - Level 3 T/Q anomalies and trends (and surface T?)


This approach is in principle very simple and quick. Allows frequency re-processing.

What's Hard:

- Dealing with clouds
- AIRS radiometric stability estimates (ie. how good?)


# SAMPLE 2x Figs

Some Small Title



AIRS, CrIS, IASI are *all* very stable  
CLARREO has removed us from this figure!

Another Small Title




These are  $2\text{-}\sigma$  B(T) statistical uncertainties  
due to inter-annual variability.


Some channels, some latitudes not gaussian  
(strat sudden warmings, QBO, etc.)

# SAMPLE 4x Figs


Some Small Title



Another Small Title



Some Small Title



Another Small Title

