

Climate Hyperspectral InfraRed Product (CHIRP) Combining AIRS, CrIS, and IASI

AIRS Science Team Meeting

L. Larrabee Strow^{1,2}, Sergio DeSouza-Machado^{1,2}, Howard Motteler²,
Chris Hepplewhite², and Steven Buczowski²

Sept 25, 2019

¹UMBC Physics Dept.

²UMBC JCET

A Climate Hyperspectral InfraRed Product (CHIRP)

Motivation

- Provide climate-level radiance time series spanning AIRS + CrIS, IASI
- User friendly by using a single spectral instrument line shape (ILS)
- A high-stability 25+ year long record of climate forcings and response
- Allows Level 2 retrievals to use a common Forward Model (RTA) and channel selection

Implementation

- Convert AIRS (and IASI) to a slightly modified CrIS ILS
- Use satellite overlaps to adjust AIRS radiometry to SNPP CrIS (0.2-0.3K)
- Three Levels of CHIRP Products:
 - CHIRP L1c (all scenes converted for Level 2 retrievals)
 - CHIRP L1 Gridded: 16 day, ~1 deg gridded radiance averages
 - Likely most used product for climate research
 - CHIRP L3 (geophysical) Anomalies
 - $T(z)$, $H_2O(z)$, T_{surf} , etc retrievals derived from CHIRP L1 Gridded
 - Quick to produce for frequent reprocessing
 - Will use very minimal a-priori information