

Physics 30 – Lesson 20

Magnetic Forces, Charged Particles.


Possible 82/74

- 1) A permanent magnet involves the alignment of magnetic domains, while an electromagnet depends on current around a solenoid.

/2

- 2)


/2



The Left Hand Rule (karate grip) is where fingers point in the direction of the magnetic field, thumb points in direction of the particle's initial motion, and the palm indicates the direction of the force. The **force** is directed **out of the page**.

- 3)

/3




Only the right hand aligns with the magnetic field direction, velocity and force on the particle. Therefore it is a **positive** charge.

#2

The particle is not affected by the magnetic field. Therefore it has **no charge**.

#3



Only the left hand aligns with the magnetic field direction, velocity and force on the particle. Therefore it is a **negative** charge.

- 4)

$$F_m = F_c$$

$$qvB_{\perp} = \frac{mv^2}{r}$$

$$qB_{\perp} = \frac{mv}{r}$$

/3

$$r = \frac{m}{qB_{\perp}} v \quad \text{r is directly proportional to v (higher v} \rightarrow \text{larger r)}$$


\therefore #1 - fastest

#2 - slowest

#3 - middle

5)

/6



The magnetic force is a centripetal force, therefore it changes the direction while speed remains constant.

6)

$$F_m = ?$$

$$q = 4.0 \times 10^{-5} \text{ C}$$

/3

$$v = 0.15 \text{ m/s}$$

$$\theta = 30^\circ$$

$$B = 0.025 \text{ T}$$

$$F_m = qvB \sin \theta$$

$$F_m = 4.0 \times 10^{-5} \text{ C} (0.15 \text{ m/s}) (0.025 \text{ T}) \sin 30^\circ$$

$$F_m = 7.5 \times 10^{-8} \text{ N}$$

7)

Since $\theta = 0$ and $\sin 0 = 0$, no force is applied \therefore the electron is unaffected

/2

8)

$$q = +3.20 \times 10^{-19} \text{ C}$$

$$v = 3.00 \times 10^8 \text{ m/s} (0.010)$$

/3

$$v = 3.00 \times 10^6 \text{ m/s}$$

$$B = 0.0030 \text{ T}$$

$$F_m = ?$$

$$F_m = qvB_\perp$$

$$F_m = 3.20 \times 10^{-19} \text{ C} (3.0 \times 10^6 \text{ m/s}) (0.0030 \text{ T})$$

$$F_m = 2.9 \times 10^{-15} \text{ N}$$

9)

$$v = 1.2 \times 10^2 \text{ m/s}$$

$$B = 0.0020 \text{ T}$$

$$q = 1.60 \times 10^{-19} \text{ C}$$

$$F = ?$$

$$F_m = qvB_\perp$$

$$F_m = (1.60 \times 10^{-19} \text{ C}) (1.2 \times 10^2 \text{ m/s}) (0.0020 \text{ T})$$

$$F_m = 3.84 \times 10^{-20} \text{ N}$$

/9

for electrons

$$F_m = 3.84 \times 10^{-20} \text{ N out of page}$$

$$\vec{a} = \frac{\vec{F}}{m}$$

$$\vec{a} = \frac{3.84 \times 10^{-20} \text{ N}}{9.11 \times 10^{-31} \text{ kg}}$$

$$\vec{a} = 4.2 \times 10^{10} \text{ m/s}^2 \text{ out of page}$$

for protons

$$F_m = 3.84 \times 10^{-20} \text{ N in to page}$$

$$\vec{a} = \frac{\vec{F}}{m}$$

$$\vec{a} = \frac{3.84 \times 10^{-20}}{1.67 \times 10^{-27} \text{ kg}}$$

$$\vec{a} = 2.3 \times 10^7 \text{ m/s}^2 \text{ in to page}$$

10)

$$m = 24 \times 1.67 \times 10^{-27} \text{ kg}$$

$$m = 4.008 \times 10^{-26} \text{ kg}$$

/5

$$F_c = F_m$$

$$\frac{mv^2}{r} = qvB_{\perp}$$

$$r = \frac{mv}{qB_{\perp}}$$

$$r = \frac{(4.008 \times 10^{-26} \text{ kg})(60000 \text{ m/s})}{(3.20 \times 10^{-19} \text{ C})(0.0800 \text{ T})}$$

$$\boxed{r = 9.39 \times 10^{-2} \text{ m}}$$

11)

Direction

For the right hand: fingers
point upward, thumb
points south – palm points
west.

/4

$$F_m = qvB_{\perp}$$

$$F_m = (3.20 \times 10^{-19} \text{ C})(7.4 \times 10^4 \text{ m/s})(5.50 \times 10^{-3} \text{ T})$$

$$\boxed{F_m = 1.30 \times 10^{-16} \text{ N west}}$$

12)

$$F_c = F_m$$

$$\frac{mv^2}{r} = qvB_{\perp}$$

$$r = \frac{mv}{qB_{\perp}}$$

/6

electron

$$r = \frac{(9.11 \times 10^{-31} \text{ kg})(9.0 \times 10^6 \text{ m/s})}{(1.60 \times 10^{-19} \text{ C})(1.2 \times 10^{-7} \text{ T})}$$

$$\boxed{r = 4.3 \times 10^2 \text{ m}}$$

proton

$$r = \frac{(1.67 \times 10^{-27} \text{ kg})(9.0 \times 10^6 \text{ m/s})}{(1.60 \times 10^{-19} \text{ C})(1.2 \times 10^{-7} \text{ T})}$$

$$\boxed{r = 7.8 \times 10^5 \text{ m}}$$

13)

$$F_c = F_m$$

$$\frac{mv^2}{r} = qvB_{\perp}$$

/4

$$q = \frac{mv}{rB}$$

$$q = \frac{(6.6 \times 10^{-27} \text{ kg})(4.4 \times 10^5 \text{ m/s})}{(0.012 \text{ m})(0.75 \text{ T})}$$

$$\boxed{q = 3.2 \times 10^{-19} \text{ C}}$$

14)

Accelerator

$$\Delta E = qV$$

/6

$$\frac{1}{2}mv^2 = qV$$

$$v = \sqrt{\frac{2qV}{m}}$$

$$v = \sqrt{\frac{2(1.60 \times 10^{-19} \text{ C})(2000 \text{ V})}{2 \times 1.67 \times 10^{-27} \text{ kg}}}$$

$$v = 4.38 \times 10^5 \text{ m/s}$$

Mass spectrometer

$$F_m = F_c$$



$$qvB_{\perp} = \frac{mv^2}{r}$$

$$r = \frac{mv}{qB_{\perp}}$$

$$r = \frac{(2 \times 1.67 \times 10^{-27} \text{ kg})(4.38 \times 10^5 \text{ m/s})}{(1.60 \times 10^{-19} \text{ C})(0.600 \text{ T})}$$

$$\boxed{r = 0.0152 \text{ m}}$$

15)

Bonus
/8

The particle follows a corkscrew path with a radius (r) and a distance (d) between adjacent rotations in the magnetic field. The velocity is broken into vertical and horizontal components. The vertical component is used to calculate the radius (r) and period (T) of rotation and the horizontal component is used to find (d).

find r

$$F_m = F_c$$

$$qvB_{\perp} = \frac{mv^2}{r}$$

$$r = \frac{mv}{qB}$$

$$r = \frac{9.11 \times 10^{-31} \text{ kg}(3.5 \times 10^6 \text{ m/s})}{1.60 \times 10^{-19} \text{ C}(0.0373 \text{ T})}$$

$$r = 5.3376 \times 10^{-4} \text{ m}$$

find period of revolution

$$T = \frac{2\pi r}{v}$$

$$T = \frac{2\pi(5.3376 \times 10^{-4} \text{ m})}{3.5 \times 10^6 \text{ m/s}}$$

$$T = 9.3 \times 10^{-10} \text{ s}$$

find d


$$d = v_H T$$

$$d = 9.37 \times 10^5 \text{ m/s}(9.3 \times 10^{-10} \text{ s})$$

$$\boxed{d = 8.73 \times 10^{-4} \text{ m}}$$

16)

/5



undeflected means
 $F_m = F_e$

$$F_E = F_m$$

$$q|\vec{E}| = qvB_{\perp}$$

$$v = \frac{|\vec{E}|}{B_{\perp}}$$

$$v = \frac{V}{dB}$$

$$v = \frac{2000V}{(0.080m)(0.0028T)}$$

$$v = 8.9 \times 10^6 \text{ m/s}$$

17)

/11

$$F_E = F_m$$

$$q|\vec{E}| = qvB_{\perp}$$

$$v = \frac{|\vec{E}|}{B_{\perp}}$$

$$v = \frac{V}{dB_{\perp}}$$

$$v = \frac{56.3 \times 10^3 V}{(0.00400m)(0.845T)}$$

$$v = 1.68 \times 10^7 \text{ m/s}$$

$$F_c = F_m$$

$$\frac{mv^2}{r} = qvB$$

$$m = \frac{qBr}{v}$$

$$m = \frac{2(1.60 \times 10^{-19} C)(0.845T)(4.15m)}{(1.68 \times 10^7 \text{ m/s})}$$

$$m = 6.68 \times 10^{-26} \text{ kg}$$

$$\text{atomic mass} = \frac{\text{mass of nucleus}}{m_{p^+ \text{ or } n}}$$

$$\text{atomic mass} = \frac{6.68 \times 10^{-26} \text{ kg}}{1.67 \times 10^{-27} \text{ kg}}$$

$$\text{atomic mass} = 40$$

look up the element with an atomic mass
 of 40 on the periodic table – ion is Ca^{2+}