

Topology of the $O(3)$ non-linear sigma model under the gradient flow

Stuart Thomas^{α,*} and Christopher Monahan^α

^α*Department of Physics, William & Mary, Williamsburg, Virginia 23187, USA*

E-mail: snthomas01@email.wm.edu, cjmonahan@wm.edu

The $O(3)$ non-linear sigma model (NLSM) is a prototypical field theory for QCD and ferromagnetism, featuring topological qualities. Though the topological susceptibility χ_t should vanish in physical theories, lattice simulations of the NLSM find that χ_t diverges in the continuum limit. We study the effect of the gradient flow on this quantity using a Markov Chain Monte Carlo method, finding that a logarithmic divergence persists. This result supports a previous study and indicates that either the definition of topological charge is problematic or the NLSM has no well-defined continuum limit. We also introduce a θ -term and analyze the topological charge as a function of θ under the gradient flow.

*The 38th International Symposium on Lattice Field Theory, LATTICE2021 26th-30th July, 2021
Zoom/Gather@Massachusetts Institute of Technology*

*Speaker




Figure 1

1. The Non-Linear Sigma Model

We study the $O(3)$ non-linear sigma model (NLSM) in 1+1 dimensions, defined by the Euclidean action

$$S_E = \frac{\beta}{2} \int d^2x \left[(\partial_t e)^2 + (\partial_x e)^2 \right]$$

where, e is 3-component real vector constrained by $|e| = 1$ and β is the inverse coupling constant. In solid-state systems, this model describes Heisenberg ferromagnets [1] and in nuclear physics, it acts as a prototype for quantum chromodynamics (QCD), the gauge theory that describes the strong nuclear force. In general, the NLSM shares key features with non-Abelian gauge theories such as QCD, including a mass gap and asymptotic freedom [2]. Therefore, the NLSM is a useful model for exploring the effect of these properties in a simpler system.

Following [3], we define topological charge density $q(x^*)$ for each plaquette x^* such that

$$Q = \sum_{x^*} q(x^*) \quad (1)$$

where

$$q(x^*) = \frac{1}{4\pi} \left[A(e(x_1), e(x_2), e(x_3)) + A(e(x_1), e(x_3), e(x_4)) \right]. \quad (2)$$

The topology of the NLSM is visualized in Fig. 1

$$S[e] \rightarrow S[e] - i\theta Q[e].$$

A nonzero θ implies nonzero $\langle Q \rangle$. Furthermore,

$$\chi_t \propto \left. \frac{d \text{Im} \langle Q \rangle}{d\theta} \right|_{\theta=0} \quad (3)$$

2. Computational Methods

Our study of the gradient flow in the NLSM is based on a computational system that simulates quantum fields numerically. We begin by implementing a numerical Monte Carlo method to simulate the lattice in two dimensions and verify our program with the well-studied ϕ^4 scalar field theory. We then generalize our model to a vector field to simulate the NLSM and implement a

numerical solution to the gradient flow. We use the data produced by this program to study the topological charge and susceptibility under the gradient flow.

We first implement the ϕ^4 model in Python. Afterwards, we transition to the NLSM, using C++ for increased efficiency. To compile the C++ simulation, we use the gcc compiler with the highest level of optimization.

3. Results




Figure 2: $\chi_t \xi_2^2$ as a function of L . ξ_2 is the second moment of the correlation function and t_0 is a scale-independent unit of flow time. We fit the data with both a logarithmic and power fit. Simulation run with 10,000 measurements every 50 sweeps, 1,000 sweep thermalization. In the $\tau = 0$ case, we have compared our result with the curve fits found in [4].




Figure 3: Nontrivial $\text{Im}\langle Q \rangle$ as a function of θ . Simulation run with 10,000 measurements, every 50 sweeps, 1,000 sweep thermalization. Note the different scaling of the y-axis.

References

- [1] C. Callan, D. Friedan, E. Martinec and M. Perry, *Strings in background fields*, .

- [2] A. Polyakov, *Interaction of goldstone particles in two dimensions. Applications to ferromagnets and massive Yang-Mills fields*, .
- [3] B. Berg and M. Lüscher, *Definition and statistical distributions of a topological number in the lattice $O(3)$ σ -model*, .
- [4] W. Bietenholz, P. de Forcrand, U. Gerber, H. Mejía-Díaz and I.O. Sandoval, *Topological Susceptibility of the 2d $O(3)$ Model under Gradient Flow*, [1808.08129](#).