

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 2 of 11

- 6 1. Find the open interval of convergence for the series

$$\sum_{n=3}^{\infty} \frac{(n+1)!}{n^n} (x+1)^{4n+3}$$

$$\sum_{n=3}^{\infty} \frac{(n+1)!}{n^n} (x+1)^{4n+3} = (x+1)^3 \sum_{n=3}^{\infty} \frac{(n+1)!}{n^n} (x+1)^{4n}$$

Let us set  $y = (x+1)^4$ , and consider the series

$$\sum_{n=3}^{\infty} \frac{(n+1)!}{n^n} y^n$$

Its radius of convergence is

$$R_y = \lim_{n \rightarrow \infty} \left| \frac{\frac{(n+1)!}{n^n}}{\frac{(n+2)!}{(n+1)^{n+1}}} \right| = \lim_{n \rightarrow \infty} \left[ \frac{(n+1)!}{n^n} \cdot \frac{(n+1)^{n+1}}{(n+2)(n+1)!} \right]$$

$$= \lim_{n \rightarrow \infty} \left[ \left( \frac{n+1}{n+2} \right) \left( \frac{n+1}{n} \right)^n \right] = e$$

Hence,  $R_x = e^{1/4}$ , and the open interval of convergence is

$$|x+1| < e^{1/4}$$

$$-e^{1/4} < x+1 < e^{1/4}$$

$$-1 - e^{1/4} < x < -1 + e^{1/4}$$



THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 4 of 11

10 3. Find the sum of the series

$$\sum_{n=1}^{\infty} (n+1)2^n x^{n-1}.$$

Answer:  $\sum_{n=1}^{\infty} (n+1)2^n x^{n-1} = \sum_{n=0}^{\infty} (n+2)2^{n+1} x^n$

The radius of convergence of the series is

$$R = \lim_{n \rightarrow \infty} \left| \frac{(n+2)2^{n+1}}{(n+3)2^{n+2}} \right| = \frac{1}{2}.$$

If we set  $S(x) = \sum_{n=1}^{\infty} (n+1)2^n x^{n-1}$ , then

$$xS(x) = \sum_{n=1}^{\infty} (n+1)2^n x^n.$$

We may integrate the series term-by-term,

$$\begin{aligned} \int xS(x)dx &= \sum_{n=1}^{\infty} 2^n \frac{x^{n+1}}{n+1} + C \\ &= \frac{2x^2}{1-2x} + C. \end{aligned}$$

Differentiation now gives

$$xS(x) = \frac{(1-2x)(4x) - 2x^2(-2)}{(1-2x)^2} = \frac{4x - 4x^2}{(1-2x)^2}.$$

Division by  $x$  now gives

$$S(x) = \frac{4-4x}{(1-2x)^2}, \text{ provided } x \neq 0.$$

Since the sum of the series when  $x=0$  is

$$\sum_{n=1}^{\infty} (n+1)2^n (0)^{n-1} = 4,$$

and the above formula also gives  $S(0)=4$ , we may

write that  $\sum_{n=1}^{\infty} (n+1)2^n x^{n-1} = \frac{4(1-x)}{(1-2x)^2}, \quad -\frac{1}{2} < x < \frac{1}{2}.$

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 5 of 11

12 4. Solve the initial-value problem

$$y'' = yy', \quad y(0) = 1, \quad y'(0) = 1/2.$$

If we set  $v = \frac{dy}{dx}$  and  $\frac{d^2y}{dx^2} = v \frac{dv}{dy}$ ,

$$v \frac{dv}{dy} = vy - 1$$

$$dv = y dy - \frac{1}{v}, \text{ provided } v \neq 0.$$

A 1-parameter family of solutions of this equation is defined implicitly by

$$v = \frac{y^2}{2} + C.$$

Since  $v = y'$  is equal to  $\frac{1}{2}$  when  $y = 1$ ,

$$\frac{1}{2} = \frac{1}{2} + C \Rightarrow C = 0.$$

$$\therefore v = \frac{y^2}{2}$$

$$\frac{dy}{dx} = \frac{y^2}{2}$$

$$\frac{dy}{y^2} = \frac{dx}{2} \text{ provided } y \neq 0.$$

A 1-parameter family of solutions of this equation is defined implicitly by

$$-\frac{1}{y} = \frac{x}{2} + D$$

Since  $y(0) = 1$ ,  $-1 = D$

$$\therefore -\frac{1}{y} = \frac{x}{2} - 1 = \frac{x-2}{2}$$

$$\therefore y = \frac{2}{2-x}$$

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 6 of 11

---

- 6 5. Find the form of a particular solution of the differential equation

$$D(D^2 - 1)(D^2 + 4)y = 3x^2e^{-x} + 10x + 5\cos x$$

as predicted by the method of undetermined coefficients. Do NOT solve for the coefficients.

The auxiliary equation is

$$m(m^2 - 1)(m^2 + 4) = 0$$

Roots are  $m = 0, \pm 1, \pm 2i$

$$y_h(x) = C_1 + C_2 e^x + C_3 e^{-x} + C_4 \cos 2x + C_5 \sin 2x$$

$$y_p(x) = Ax^3 e^{-x} + Bx^2 e^{-x} + Cx e^{-x} + Dx^2 + Ex + F \cos x + G \sin x$$

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 7 of 11

- 12 6. A 100 gram mass is suspended from a spring with constant  $25/2$  newtons per metre. It is set into vertical motion by pulling it 5 centimetres below its equilibrium position and giving it velocity 2 metres per second upward. During its subsequent motion, damping is equal to 3 times velocity.

- (a) Find the position of the mass as a function of time.  
(b) Is the motion underdamped, overdamped, or critically damped?  
(c) Determine whether the mass ever passes through its equilibrium position. If it does, find the time(s) when this occurs.

(a) The initial value problem for displacements is:  

$$\frac{1}{10} \frac{d^2x}{dt^2} + 3 \frac{dx}{dt} + \frac{25}{2}x = 0, \quad x(0) = -\frac{1}{20}, \quad x'(0) = 2.$$

The auxiliary equation is:  

$$0 = \frac{1}{10}m^2 + 3m + \frac{25}{2} = \frac{1}{10}(m^2 + 30m + 125) = \frac{1}{10}(m+5)(m+25)$$

Roots are  $m = -5, -25$ . Thus,

The initial conditions give  

$$x(t) = Ae^{-5t} + Be^{-25t}$$

$$-\frac{1}{20} = A + B, \quad 2 = -5A - 25B$$

Solutions are  $A = \frac{3}{80}$  and  $B = -\frac{7}{80}$ .

Thus,  $x(t) = \frac{3}{80}e^{-5t} - \frac{7}{80}e^{-25t}$  m.

(b) Motion is overdamped.

(c) The mass passes through equilibrium if

$$0 = \frac{3}{80}e^{-5t} - \frac{7}{80}e^{-25t}$$

$$3e^{-5t} = 7e^{-25t}$$

$$e^{20t} = \frac{7}{3}$$

$$20t = \ln\left(\frac{7}{3}\right)$$

$$t = \frac{1}{20} \ln\left(\frac{7}{3}\right) \text{ s.}$$

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 8 of 11

- 8 7. Find the Laplace transform of the function


$$f(t) = e^{4t} \sin 3t h(t-2).$$

$$\begin{aligned} F(s) &= e^{-2s} \mathcal{L} \{ e^{4(t+2)} \sin 3(t+2) \} \\ &= e^{-2s} e^8 \mathcal{L} \{ e^{4t} (\sin 3t \cos 6 + \cos 3t \sin 6) \} \\ &= e^{8-2s} \left[ \frac{\cos 6(3)}{(s-4)^2+9} + \frac{(s-4) \sin 6}{(s-4)^2+9} \right] \end{aligned}$$

or

$$\begin{aligned} F(s) &= \mathcal{L} \{ \sin 3t h(t-2) \}_{s-4} \\ &= e^{-2(s-4)} \mathcal{L} \{ \sin 3(t+2) \}_{s-4} \\ &= e^{8-2s} \mathcal{L} \{ \sin 3t \cos 6 + \cos 3t \sin 6 \}_{s-4} \\ &= e^{8-2s} \left[ \frac{3 \cos 6}{(s-4)^2+9} + \frac{(s-4) \sin 6}{(s-4)^2+9} \right] \end{aligned}$$

- 10 8. Find the Laplace transform of the function in the figure below. You need not simplify your result.



$$\begin{aligned} F(s) &= \frac{1}{1-e^{-2s}} \int_0^5 f(t) e^{-st} dt \\ &= \frac{1}{1-e^{-2s}} \mathcal{L} \{ 2t^2 [h(t)-h(t-1)] + 2[h(t-1)-h(t-2)] \} \\ &= \frac{1}{1-e^{-2s}} \left[ \frac{4}{s^3} - 2e^{-s} \mathcal{L} \{ (t+1)^2 \} + \frac{2}{s} e^{-s} - \frac{2}{s} e^{-2s} \right] \\ &= \frac{1}{1-e^{-2s}} \left[ \frac{4}{s^3} - 2e^{-s} \mathcal{L} \{ t^2 + 2t + 1 \} + \frac{2}{s} e^{-s} - \frac{2}{s} e^{-2s} \right] \\ &= \frac{1}{1-e^{-2s}} \left[ \frac{4}{s^3} - 2e^{-s} \left( \frac{2}{s^3} + \frac{2}{s^2} + \frac{1}{s} \right) + \frac{2}{s} e^{-s} - \frac{2}{s} e^{-2s} \right] \end{aligned}$$

THE UNIVERSITY OF MANITOBA

DATE: December 16, 2013 (Afternoon)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 9 of 11

- 10 9. Find the inverse Laplace transform for the function

$$\frac{(3s^2 + s - 6)e^{-2s}}{s^3 + 3s^2}$$

Since  $\frac{3s^2 + s - 6}{s^3 + 3s^2} = \frac{3s^2 + s - 6}{s^2(s+3)} = \frac{1}{s} - \frac{2}{s^2} + \frac{2}{s+3}$

$$\mathcal{L}^{-1} \left\{ \frac{3s^2 + s - 6}{s^3 + 3s^2} \right\} = 1 - 2t + 2e^{-3t}$$

Thus,  $\mathcal{L}^{-1} \left\{ \frac{(3s^2 + s - 6)e^{-2s}}{s^3 + 3s^2} \right\} = [1 - 2(t-2) + 2e^{-3(t-2)}] h(t-2)$

$$= (5 - 2t + 2e^{-3t}) h(t-2)$$



DATE: April 11, 2011 (Morning)

FINAL EXAMINATION

DEPARTMENT & COURSE NO: MATH2132

TIME: 3 hours

EXAMINATION: Engineering Mathematical Analysis 2 EXAMINER: D. Trim

PAGE NO: 10 of 11

12 10. Solve the initial-value problem

$$\frac{d^2y}{dt^2} + 4\frac{dy}{dt} + 6y = 4\delta(t-3), \quad y(0) = 2, \quad y'(0) = 1.$$

When we take Laplace transforms,

$$[s^2Y - 2s - 1] + 4[sY - 2] + 6Y = 4e^{-3s}$$

$$Y(s) = \frac{2s + 9 + 4e^{-3s}}{s^2 + 4s + 6}$$

$$= \frac{2s + 9}{s^2 + 4s + 6} + \frac{4e^{-3s}}{s^2 + 4s + 6}$$

$$= \frac{2(s+2) + 5}{(s+2)^2 + 2} + \frac{4e^{-3(s+2)+6}}{(s+2)^2 + 2}$$

$$y(t) = e^{-2t} \mathcal{L}^{-1} \left\{ \frac{2s+5}{s^2+2} \right\} + 4e^{-2t} \mathcal{L}^{-1} \left\{ \frac{e^{-3s}}{s^2+2} \right\}$$

$$= e^{-2t} \left[ 2 \cos \sqrt{2}t + \frac{5}{\sqrt{2}} \sin \sqrt{2}t \right]$$

$$+ \frac{4}{\sqrt{2}} e^{-2t} \sin \sqrt{2}(t-3) h(t-3)$$