

Math 2130 - Engineering Mathematical Analysis 1

Tutorial § 11.6, 11.10, 11.11

11.6.5 Determine the area of the triangle that passes through the points: $(3, 0, 2)$, $(2, 1, 3)$ and $(1, 3, 7)$.
Hint: See tutorial question 11.5.1.

11.10.1 Find parametric equations of the ellipse: $\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} = 1$, traversed (i.e., travelled or gone over) once, counter-clockwise; a and b are positive constants. Note: (x_0, y_0) is the centre of the ellipse and a and b are the semi-axes.
Hint: Use $\cos^2 t + \sin^2 t = 1$. Note: Unlike the case of a circle (when $a = b$), if $a \neq b$, then this “ t ” will *not* be the angle between the line joining a point on the ellipse and the horizontal line through the centre. Can you prove this?

In questions 11.10.2-11.10.4 below, find parametric equations of the given curve and also name the surfaces which intersect in the curve and describe the shape of the curve:

11.10.2 Find parametric equations of the curve: $x + 2y + 3z = 6$, $y - 2z = 3$, directed so that z increases along the curve.

11.10.3 Find parametric equations of the curve: $z = x^2 + y^2$, $z = \sqrt{6 - x^2 - y^2}$, directed so that y decreases when x is positive.

11.10.4 Find parametric equations of the curve: $z = 4x^2 + y^2$, $z = 5 + 8x$, so that the curve is traversed once, counter-clockwise, when viewed from above.

11.11.1 For each of 11.10.1-11.10.4 above, determine a *unit* tangent vector in the direction of the curve.

11.11.2 For the curve in 11.10.2 above, calculate the length of that portion where: $-14 \leq x \leq 7$.

11.11.3 For 11.10.3 above, calculate the length of the curve.

11.11.4 For 11.10.4 above, set up an integral for the length of the curve, and *simplify*, but do *not evaluate* this integral.

11.11.5 Calculate the length of the curve: $y = x^{(2/3)}$, $z = 2x^{(2/3)}$, for the portion where: $0 \leq x \leq 1$.
Use the parametrization where $x = t^3$.

Answers: 11.6.5: $\text{Area} = \frac{\sqrt{14}}{2}$.

11.10.1: $x = x_0 + a \cos t$, $y = y_0 + b \sin t$; $0 \leq t \leq 2\pi$. Consider the line: $t = \frac{\pi}{4}$.

11.10.2: $x = -7t$, $y = 2t + 3$, $z = t$; all real t . Two planes; the curve is a line.

11.10.3: Two standard forms are:

$$x = \sqrt{2} \cos(t), \quad y = -\sqrt{2} \sin(t), \quad z = 2; \quad 0 \leq t \leq 2\pi.$$

$$\text{or, } x = \sqrt{2} \sin(t), \quad y = \sqrt{2} \cos(t), \quad z = 2; \quad 0 \leq t \leq 2\pi.$$

A circular paraboloid and a sphere; the curve is a circle.

11.10.4: $x = 1 + \frac{3}{2} \cos t$, $y = 3 \sin t$, $z = 13 + 12 \cos t$; $0 \leq t \leq 2\pi$.

An elliptic paraboloid and a plane; the curve is an ellipse.

11.11.1: (1) $\hat{T} = \frac{(-a \sin t, b \cos t)}{\sqrt{a^2 \sin^2 t + b^2 \cos^2 t}}$; (2) $\hat{T} = \frac{(-7, 2, 1)}{\sqrt{54}} = \frac{(-7, 2, 1)}{3\sqrt{6}}$;

(3) $\hat{T} = (-\sin(t), -\cos(t), 0)$, or, $\hat{T} = (\cos(t), -\sin(t), 0)$ (depending on your parametrization);

(4) $\hat{T} = \frac{(-\sin t, 2 \cos t, -8 \sin t)}{\sqrt{4 \cos^2 t + 65 \sin^2 t}} = \frac{(-\sin t, 2 \cos t, -8 \sin t)}{\sqrt{4 + 61 \sin^2 t}}$.

11.11.2: (Re 11.10.2): $L = 9\sqrt{6}$.

11.11.3: (Re 11.10.3): $L = (2\sqrt{2})\pi$.

11.11.4: (Re 11.10.4): $L = \int_0^{2\pi} \frac{3}{2} \sqrt{4 \cos^2 t + 65 \sin^2 t} \, dt = \int_0^{2\pi} \frac{3}{2} \sqrt{4 + 61 \sin^2 t} \, dt$.

Note: This is called an *elliptic integral* and *cannot* be calculated exactly.

11.11.5: $L = \int_0^1 \sqrt{9t^4 + 20t^2} \, dt = \int_0^1 t \sqrt{9t^2 + 20} \, dt = \frac{1}{27} \left[29^{3/2} - 20^{3/2} \right] = \frac{1}{27} \left[29^{3/2} - 40\sqrt{5} \right]$.