## THE UNIVERSITY OF MANITOBA

Name

**Student Number** 

COMP1012: Computer Programming for Scientists and Engineers Midterm In Class Exam (A01—Andres)

2015 October 29, 8:30 am Time: 50 minutes

### Instructions:

- 1. Answer all questions on this paper. For multiple choice questions, circle the letter of the **best** or most complete choice. For short answer questions, write your answer in the space provided.
- 2. Extra work space is available on page 3.
- 3. You will find a Python Guide along with your midterm; ask if you don't have one. You may *not* use your own copy. No other aids (such as calculators or cell phones) are permitted.
- 4. You have 50 minutes to complete the exam.

Marks for Part 1	Part 2A	Part 2B	Part 3	Total
/ 4	/ 4	/ 4	/ 4	/16

#### Part 1: Predict the output [4 x 1 mark]

In each row of the table below, mentally execute the code on the left and enter the expected output in the box on the right. Each table row is separate. Use the space below for scrap work.

	Code Fragment	Expected output		
Α.	What is printed by print(2 / 2 + 3 / 1) ?			
В.	<pre>What is printed by print(tuple(range(1,-3,-2))) ?</pre>			
С.	What is printed by print(1 // 4 and 1 / 4) ?			
D.	What is printed by print(2 + 1 != 1 < 2) ?			

Work space:

Name

**Student Number** 

COMP1012: Computer Programming for Scientists and Engineers Midterm In Class Exam (A01—Andres)

2015 October 29, 8:30 am Time: 50 minutes

# Part 2: Write a program [Total 8 marks]

2. [8 marks] A friend has described to you a strange function that has this infinite series:

$$f(x) = \frac{x}{1!} - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

She speculates that  $f(x) \equiv \sin(x)$ . Write a complete program to print out the table of values shown to the right, to compare the two functions. In evaluating the series, sum all terms that are at least  $10^{-16}$  in absolute value and only those terms. In comparing the function values, use a comparison tolerance of  $10^{-13}$ .

# put imports here

TESTING SIN SERIES sin(x)f(x)Equal x[rad] 0.1 0.0998334166468 0.0998334166468 True 0.2 0.1986693307951 0.1986693307951 True 0.3 0.2955202066613 0.2955202066613 True 0.4 0.3894183423087 0.3894183423087 True 0.5 0.4794255386042 0.4794255386042 True 0.5646424733950 0.6 0.5646424733950 True 0.7 0.6442176872377 0.6442176872377 0.7173560908995 0.7173560908995 True 0.7833269096275 0.7833269096275 True

# global constants here
EPS1 =

EPS1 = EPS2 =

def main() : # put your code for the table here

For marker use only

Item Mark

A

B

C

D

E

Sum

 $\operatorname{def}$  fnc(xx) : # your code to return the series sum

THE UNIVERSITY OF MANITOBA

Name

**Student Number** 

COMP1012: Computer Programming for Scientists and Engineers Midterm In Class Exam (A01—Andres)

2015 October 29, 8:30 am

Time: 50 minutes

main()

Page intentionally left blank (apart from this line); test continues on next page

Name

**Student Number** 

COMP1012: Computer Programming for Scientists and Engineers Midterm In Class Exam (A01—Andres)

2015 October 29, 8:30 am Time: 50 minutes

# Part 3: Circle the letter of the best answer, or provide the required answer [4 x 1 mark]

Given the following lines have just been executed, which of the options below creates a list of all numbers from seq1 that contain the digit '3', and only those numbers?

$$seq1 = [-3, 4, -4, -5, 4, 0, 1, -4, 3, -3]$$
  
 $seq2 = []$ 

- a) for num in seq1: seq2.append((num % 3 == 0) \* num)
- for num in seq1: seq2 = [num] \* (num % 3 == 0)b)
- seq2 += [num] \* (num % 3 == 0)c) for num in seq1:
- seq2 += [num] \* ('3' in str(num)) d) for num in seq1:
- for num in seq1: seq2 = [num] \* ('3' in str(num))e)
- B. Which of the following does NOT print "any\_banana" given this definition?

```
string = "any banana"
```

- string.replace(' ', '\_'); print(string)
  print(string[:3] + '\_' + string[4:]) a)
- b)
- print(string.replace(" ", "\_")) c)
- d) print('\_'.join(string.split()))
- strList = list(string); strList[3] = '\_'; print(''.join(strList)) e)
- C. Which of the following statements about tuples, lists and strings is false, assuming seq is a tuple, a list or a string that is large enough to make the expressions valid?
  - seq[-2] is the second last item if seq is any of these data types. a)
  - b) bool(seq[3:3]) is False for any of these data types.
  - 2 \* seq is twice the length of seq for any of these data types. c)
  - d) seg[2] is the second item if seg is any these data types.
  - seg += seg doubles the size of seg for any of these data types.
- D. Using good coding practices and the same rules as QuizMaster, write a Python expression to evaluate this math expression, assuming math has already been imported:

$$\ln \left( \left\lceil \sin \left( \tan \left( \log_{10} \left( 5 \right) - \frac{|b|}{-5} \right) \right) \right\rceil \right)$$

Put expression here