Studybeesofficial
EDUCATIONAL PLATFORM
www.studybeesofficial.com

SSC-I NATIONAL BOOK FOUNDATION FBISE

MATHEMATICS TEST - CHAPTER 3: SETS AND RELATIONS

Name:		Roll No.		Section:

Total Time: 2:30 Hour **Total Marks: 150**
Time Allowed: 150 Minutes

PAPER OBJECTIVE
SECTION-A (MARKS 150)

NOTE: Section-A is compulsory. All parts of this section are to be answered on these pages and handed over to the center superintendent. Deleting / overwriting is not allowed. Do not use lead pencil.

Q.1 Fill the relevant bubble for each part. All parts carry one mark.

Questions	A	B	C	D	A B C D
1. The set of all letters in the word "set" is	{s, e, t, s}	{s, e, t}	{s, t}	{e, t}	○●○○
2. The set of natural numbers N is	{0, 1, 2, ...}	{1, 2, 3, ...}	{..., −1, 0, 1, ...}	{2, 3, 5, ...}	○●○○
3. The set of rational numbers Q is written as	{p/q | p, q ∈ Z, q ≠ 0}	{p/q | p, q ∈ Z, q = 0}	{x | x ≠ p/q}	{x | x ∈ N}	●○○○
4. The set of squares of natural numbers {1², 2², 3², ...} is	{1, 4, 9, 16, ...}	{1, 2, 3, 4, ...}	{2, 4, 6, 8, ...}	{1, 3, 5, 7, ...}	●○○○
5. Writing a set by listing all its elements is called	Set builder form	Tabular (roster) form	Rule method	Descriptive form	○●○○
6. Sets of numbers N, W, Z, E, O, P are first written in	Descriptive then tabular form	Set builder form only	Venn form only	Graph form	●○○○
7. The set of natural numbers less than 1 is	{ } (empty)	{0}	{1}	{0, 1}	●○○○
8. Which of the following is a well-defined set?	Lighter stones	Stones weighing less than 1 kg	Beautiful flowers	Tall students	○●○○
9. The set of integers is denoted by	N	W	Q	Z	○○○●
10. Another name for set builder form is	Rule method	Roster method	Tabular method	Listing method	●○○○
11. The set Q′ of irrational numbers is written as	{p/q | q ≠ 0}	{x | x ≠ p/q, p, q ∈ Z, q ≠ 0}	{x | x ∈ Z}	{x | x ∈ Q}	○●○○
12. Sets Q, Q′ and R are first written in descriptive form and then in	Tabular form	Set builder form	Table form	Arrow form	○●○○
13. The set of even numbers E is	{±1, ±3, ±5, ...}	{2, 3, 5, ...}	{1, 2, 3, ...}	{0, ±2, ±4, ...}	○○○●
14. The set of odd numbers O is	{0, ±2, ±4, ...}	{2, 3, 5, ...}	{1, 2, 3, ...}	{±1, ±3, ±5, ...}	○○○●

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio

Questions	A	B	C	D	A B C D
15. Writing a set by its defining rule using a variable is called	Tabular form	Roster form only	Set builder form	Venn form	○○●○
16. The set of real numbers R can be written as	{x | x ∈ Q and x ∈ Q′}	{x | x ∈ Z}	{x | x ∈ Q or x ∈ Q′}	{x | x ∈ N or x ∈ W}	○○●○
17. The set of whole numbers W is	{1, 2, 3, ...}	{..., −2, −1, 0, ...}	{0}	{0, 1, 2, 3, ...}	○○○●
18. In the pattern of mangoes in boxes (2, 4, 6, 8 ...), box 6 contains	10 mangoes	14 mangoes	11 mangoes	12 mangoes	○○○●
19. The set of prime numbers P begins as	{1, 2, 3, 5, ...}	{2, 3, 5, 7, 11, ...}	{1, 3, 5, 7, ...}	{2, 4, 6, 8, ...}	○●○○
20. The set of prime numbers which are also even is	{2}	{ }	{2, 4}	{1, 2}	●○○○
21. The set of real numbers cannot be written in tabular form because	It is empty	It is finite	Its elements cannot be listed	It has one element	○○●○
22. The symbol "|" in set builder notation is read as	Divides	Or	Not	Such that	○○○●
23. A set is a well-defined collection of	Repeated objects	Only numbers	Distinct objects	Only letters	○○●○
24. Tabular form of {x | x is a multiple of 3, x < 20, x ∈ N} is	{3, 9, 15}	{2, 3, 6, 9, 12, 15, 18}	{3, 6, 9, 12, 15, 18}	{1, 2, 3, 4, 5, 6, 7}	○○●○
25. The set {y | y ∈ W ∧ $y^3 - 1 = 7$} in tabular form is	{1}	{2, −2}	{−2}	{2}	○○○●
26. Set builder form of {1, 2, 3, 6} is	{x | x is a multiple of 6}	{x | x ∈ N ∧ x < 6}	{x | x is a prime < 6}	{x | x is a factor of 6}	○○○●
27. Set builder form of "set of prime numbers less than 17" is	{x | x ∈ N ∧ x < 17}	{x | x ∈ P ∧ x > 17}	{x | x ∈ P ∧ x < 17}	{x | x ∈ Z ∧ x < 17}	○○●○
28. Tabular form of {3x | x ∈ N ∧ x < 5} is	{3, 9}	{3, 6, 9, 12}	{2, 3, 6, 9, 12}	{3, 6, 9}	○●○○
29. Tabular form of {x | x ∈ N ∧ x is even and x < 10} is	{2, 4, 6}	{2, 4, 6, 8, 9}	{3, 5, 7, 9}	{2, 4, 6, 8}	○○○●
30. Tabular form of {x | x ∈ W ∧ x ≤ 3} is	{0, 1, 2, 3}	{0, 1, 2, 3, 4}	{1, 2, 3, 4}	{0, 1, 2}	●○○○
31. The complement of A, written A^c or A′, is equal to	A − U	A ∪ U	U − A	A ∩ U	○○●○

Questions	A	B	C	D	A B C D
32. (A ∪ B) ∪ C = A ∪ (B ∪ C) is the	Associative property of union	Distributive property	Commutative property	Complement law	●○○○
33. For disjoint sets A and B, n(A ∪ B) equals	n(A) − n(B)	n(A) · n(B)	n(A) + n(B) − 1	n(A) + n(B)	○○○●
34. For any sets A and B, A ∩ B is always a	Subset of A	Superset of A	Empty set	Universal set	●○○○
35. Two sets are disjoint if	A ∪ B = ∅	A ⊂ B	A = B	A ∩ B = ∅	○○○●
36. The intersection of sets A and B contains elements that are	In A only	In B only	In neither	Common to both A and B	○○○●
37. The difference A − B consists of elements that are	In A but not in B	In B but not in A	In both A and B	In neither A nor B	●○○○
38. In general, A − B and B − A are	Always equal	Always empty	Always disjoint from A ∪ B	Not equal	○○○●
39. For disjoint sets A and B, n(A − B) equals	n(B)	0	n(A)	n(A) − n(B)	○○●○
40. A pictorial representation of sets is called a	Bar graph	Venn diagram	Pie chart	Number line	○●○○
41. For any sets A and B, A is always a	Superset of A ∪ B	Subset of A ∪ B	Proper subset of A ∩ B	Disjoint from A ∪ B	○●○○
42. The union of A and B is denoted by	A ∪ B	A ∩ B	A − B	A^c	●○○○
43. For three sets, brackets are used in (A ∪ B) ∩ C because	They look nicer	The two expressions differ	Union has priority	They are always equal	○●○○
44. For any two sets, n(A ∪ B) also equals	n(A − B) + n(B − A)	n(A − B) + n(A ∩ B) + n(B − A)	n(A) + n(B)	n(A ∩ B) − n(A − B)	○●○○
45. For any two sets, n(A − B) equals	n(A) + n(A ∩ B)	n(A) − n(B)	n(B) − n(A ∩ B)	n(A) − n(A ∩ B)	○○○●
46. A ∩ B in set builder form is	{x | x ∈ A ∨ x ∈ B}	{x | x ∈ A ∧ x ∉ B}	{x | x ∈ A ∧ x ∈ B}	{x | x ∉ A}	○○●○
47. For overlapping sets, n(A ∪ B) equals	n(A) + n(B) + n(A ∩ B)	n(A) + n(B) − n(A ∩ B)	n(A) · n(B)	n(A) − n(B)	○●○○
48. A − B in set builder form is	{x | x ∈ B ∧ x ∉ A}	{x | x ∈ A ∨ x ∈ B}	{x | x ∈ A ∧ x ∉ B}	{x | x ∈ A ∧ x ∈ B}	○○●○

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio

Questions	A	B	C	D	A B C D
49. For any two sets, n(A − B) also equals	n(A ∪ B) − n(B)	n(A ∪ B) + n(B)	n(A ∩ B) − n(B)	n(A) + n(B)	●○○○
50. For any set A, A ∪ { } equals	∅	U	{∅}	A	○○○●
51. The union of sets A and B contains elements that are	In A or in B or in both	In both A and B only	In A but not in B	Not in A and not in B	●○○○
52. The complement of A in set builder form is	{x | x ∈ A ∧ x ∉ U}	{x | x ∈ U ∧ x ∉ A}	{x | x ∈ A}	{x | x ∈ U ∧ x ∈ A}	○●○○
53. For any set A, A ∪ A equals	∅	A	U	2A	○●○○
54. For any set A, A ∩ { } equals	A	U	∅	{∅}	○○●○
55. For any set A, A ∩ A equals	A	∅	U	2A	●○○○
56. A ∪ B in set builder form is	{x | x ∈ A ∧ x ∈ B}	{x | x ∈ A ∧ x ∉ B}	{x | x ∈ A ∨ x ∈ B}	{x | x ∉ A ∧ x ∉ B}	○○●○
57. The elements of A are never present in	A^c	A ∪ B	U	A ∪ A	●○○○
58. In a Venn diagram, the universal set is represented by	A circle	A triangle	A point	The interior of a rectangle	○○○●
59. Two sets are overlapping if they have a common element and	Neither is a subset of the other	One is a subset of the other	They are equal	They are empty	●○○○
60. A is a subset of B if	Every element of A is in B	Every element of B is in A	A and B are disjoint	A has more elements than B	●○○○
61. If A = {4, 8, 9}, B = {4, 5, 9}, C = {2, 3, 4, 5, 8, 9}, find (A ∩ B) ∪ C	{5, 8}	{4, 5, 8, 9}	{1, 2, 3, 4, 5, 8, 9}	{2, 3, 4, 5, 8, 9}	○○○●
62. If A = {2, 3, 5, 7, 8}, B = {2, 3, 4, 6, 8, 9}, find A − B	{1, 4, 6, 9, 10}	{2, 3, 4, 5, 6, 7, 8, 9}	{5, 7}	{4, 6, 9}	○○●○
63. If n(A) = 48, n(B) = 40 and n(A ∪ B) = 84, then n(A ∩ B) =	4	8	88	5	●○○○
64. If A = {2, 3, 5, 6, 7, 8}, B = {2, 3, 4, 5, 6}, C = {1, 3, 4, 6, 7, 9}, find A ∪ (B ∩ C)	{3, 4, 6, 7}	{2, 3, 4, 5, 6, 7, 8}	{1, 2, 3, 4, 5, 6, 7, 8}	{4, 7, 8}	○●○○
65. Of 120 students, 70 like English, 60 like Maths, 25 both. How many like English only?	45	46	35	70	●○○○

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio

Questions	A	B	C	D	A B C D
66. If A = {4, 5, 6, 7}, B = {1, 5, 7, 8, 9}, C = {3, 5, 6, 8}, find (A ∩ B) − C	∅	{4, 6}	{7}	{1, 4, 6, 8, 9}	○○●○
67. If n(A) = 30, n(B) = 49 and n(A ∩ B) = 18, then n(B − A) =	19	49	31	61	○○●○
68. If n(A) = 19, n(B) = 12 and n(A ∪ B) = 28, then n(A ∩ B) =	9	4	2	3	○○○●
69. If n(A) = 52, n(B) = 43 and n(A ∩ B) = 10, then n(A − B) =	42	85	33	62	●○○○
70. If n(A) = 54, n(B) = 26 and n(A ∩ B) = 20, then n(A ∪ B) =	60	80	100	79	●○○○
71. Out of 50 people, 30 like tea and 35 like coffee; each likes at least one. Both =	17	5	15	13	○○●○
72. For disjoint sets with n(A) = 11 and n(B) = 15, n(A ∪ B) =	15	26	4	165	○●○○
73. Of 100 students, 55 like English, 45 like Maths and 20 like both. How many like neither?	40	0	80	20	○○○●
74. Of 70 students, 40 like English, 35 like Maths and 18 like both. How many like neither?	13	31	12	57	●○○○
75. Of 50 students, 8 learn both languages, 26 learn English. All learn at least one. Chinese learners =	26	24	32	18	○○●○
76. If A = {4, 6, 7}, B = {3, 4, 7, 8}, C = {2, 3, 4, 6, 7, 8}, find (A ∪ B) ∩ C	{3, 4, 6, 7, 8}	{4, 6, 7}	{4, 7}	{3, 6, 8}	●○○○
77. If A = {1, 2, 3, 6, 9}, B = {2, 8, 9}, C = {2, 3, 4, 5, 6, 8}, find A ∩ (B ∪ C)	{8}	{2, 3, 6, 8}	{2, 3, 6, 9}	{1, 3, 6, 8}	○○●○
78. If A = {1, 3, 4, 5, 8, 9}, B = {1, 2, 3, 4, 7}, C = {1, 4, 5, 7}, find (A ∩ B) − C	{2, 5, 7, 8, 9}	{1, 4, 5, 7}	{3}	{1, 3, 4}	○○●○
79. If n(A) = 52, n(B) = 17 and n(A ∩ B) = 11, then n(A − B) =	63	41	58	52	○●○○
80. If n(A) = 31, n(B) = 19 and n(A ∩ B) = 15, then n(A ∪ B) =	50	65	49	35	○○○●
81. If n(A) = 43, n(B) = 35 and n(A ∩ B) = 26, then n(A − B) =	43	17	69	52	○●○○
82. If n(A) = 47, n(B) = 12 and n(A ∩ B) = 4, then n(A ∪ B) =	58	54	55	59	○○●○

Questions	A	B	C	D	A B C D
83. In a girl college, 350 play badminton, 280 play table tennis, 150 play both. Total students =	630	480	280	780	○●○○
84. If U = {1, ..., 10} and A = {1, 2, 4, 9}, find A′	{3, 5, 6, 7, 8, 10}	{1, 3, 5, 6, 7, 8, 10}	{3, 5, 6, 7, 8}	{1, 3, 4, 7, 8, 10}	●○○○
85. If U = {1, ..., 10} and A = {1, 2, 6, 9}, find A′	{1, 5, 6, 7, 10}	{3, 4, 5, 7, 8, 10}	{1, 2, 6, 9}	{3, 4, 5, 7, 8}	○●○○
86. If n(A) = 33, n(B) = 18 and n(A ∪ B) = 42, then n(A ∩ B) =	9	15	8	10	●○○○
87. If U = {1, ..., 10} and A = {1, 2, 7}, find A′	{3, 4, 5, 6, 8, 9, 10}	{1, 2, 3, 6, 7}	{2, 7}	{3, 4, 5, 6, 8, 9}	●○○○
88. If A = {1, 3, 5}, B = {5, 7, 8}, find B − A	{1, 3, 5, 7, 8}	{7}	{1, 7, 8}	{7, 8}	○○○●
89. Of 90 students, 50 like English, 40 like Maths and 15 like both. How many like neither?	30	75	0	15	○○○●
90. If n(A) = 49, n(B) = 20 and n(A ∩ B) = 11, then n(A − B) =	38	58	29	60	●○○○
91. Out of 70 people, 48 like tea and 40 like coffee; each likes at least one. Both =	8	20	18	16	○○●○
92. Of 80 students, 40 like English, 34 like Maths, 9 both. How many like English only?	49	32	65	31	○○○●
93. If A = {1, 3, 4, 6}, B = {1, 3, 8}, C = {5, 6, 8}, find A − (B ∪ C)	{1, 3, 6}	{4, 6}	{1, 3, 4, 6, 8}	{4}	○○○●
94. Of 60 students, 35 like English, 30 like Maths, 12 both. How many like English only?	24	23	35	47	○●○○
95. If n(A) = 55, n(B) = 11 and n(A ∩ B) = 8, then n(A ∪ B) =	66	57	58	65	○○●○
96. If n(A) = 43, n(B) = 32 and n(A ∪ B) = 73, then n(A ∩ B) =	1	3	2	75	○○●○
97. If n(A) = 48, n(B) = 19 and n(A ∩ B) = 3, then n(A ∪ B) =	64	70	66	67	●○○○
98. Of 200 students, 120 like English, 90 like Maths and 40 like both. How many like neither?	70	160	30	170	○○●○
99. Of 940 children, 400 are at primary, 240 elementary, 175 secondary. Out of school =	135	175	765	125	○○○●

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio

Questions	A	B	C	D	A B C D
100. If n(A) = 30, n(B) = 46 and n(A ∪ B) = 53, then n(A ∩ B) =	24	76	23	16	○○●○
101. If n(A) = 45, n(B) = 35 and n(A ∪ B) = 60, find n(A ∩ B)	19	15	20	80	○○●○
102. If n(A) = 50, n(B) = 12 and n(A ∪ B) = 54, then n(A ∩ B) =	62	4	7	8	○○○●
103. If U = {1, ..., 10} and A = {1, 4, 6, 8}, find A′	{2, 3, 5, 7, 9}	{2, 3, 5, 7, 9, 10}	{1, 2, 3, 5, 7, 9, 10}	{1, 2, 3, 4, 8, 9, 10}	○●○○
104. The number of subsets of A × B is	$n(A \times B)^2$	$2 \times n(A \times B)$	n(A) + n(B)	2^n(A×B)	○○○●
105. If R = {(a, b)}, the inverse relation R^{-1} is	{(a, a)}	{(−a, −b)}	{(a, −b)}	{(b, a)}	○○○●
106. In the ordered pair (a, b), the element a is called the	Second element	First element	Range	Domain only	○●○○
107. n(A × B) and n(B × A) are always	Equal	Different	Zero	Negative	●○○○
108. A × B = B × A if and only if	A = B	A ⊂ B	A ∩ B = ∅	n(A) ≠ n(B)	●○○○
109. If A ⊂ B, then A − B =	A	∅	B	B − A	○●○○
110. A relation can be represented by	Only tables	Tables, ordered pairs and graphs	Only graphs	Only equations	○●○○
111. If n(A) = p, then n(A × A) =	p^2	2p	p	2^p	●○○○
112. For R = {(a, b) | a, b ∈ Z ∧ a + b = 0}	Dom R ⊂ Range R	Dom R = Range R	Range R ⊂ Dom R	Dom R ∩ Range R = ∅	○●○○
113. The range of R^{-1} is equal to the	Domain of R	Range of R	Domain of R^{-1}	Empty set	●○○○
114. Two ordered pairs (a, b) and (c, d) are equal if and only if	a = d and b = c	a + b = c + d	a = c and b = d	a = b and c = d	○○●○
115. If A ∪ B = A and A ∩ B = B, then	A ⊂ B	A ≠ B	A ∩ B = ∅	A ⊇ B	○○○●
116. The domain of R^{-1} is equal to the	Domain of R	Range of R^{-1}	Empty set	Range of R	○○○●

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio

Questions	A	B	C	D	A B C D
117. In general A × B is	Equal to B × A	Equal to A ∩ B	Not equal to B × A	Equal to A ∪ B	○○●○
118. If A − B = B − A = { }, then	A ⊂ B only	A ∩ B = ∅	A ≠ B	A = B	○○○●
119. The set builder form of A − B is	{x | x ∈ A ∧ x ∈ B}	{x | x ∈ B}	{x | x ∉ A ∧ x ∈ B}	{x | x ∈ A ∧ x ∉ B}	○○○●
120. A pair in which the order of numbers matters is called an	Unordered pair	Empty set	Ordered pair	Venn pair	○○●○
121. The empty relation R = { } is called a	Void relation	Universal relation	Identity relation	Inverse relation	●○○○
122. A binary relation is a set of	Ordered pairs	Single elements	Numbers only	Sets of sets	●○○○
123. Any subset of the Cartesian product A × B is a	Function only	Universal set	Binary relation	Venn diagram	○○●○
124. The number of elements in A × B is	n(A) + n(B)	n(A) − n(B)	n(A) ÷ n(B)	n(A) × n(B)	○○○●
125. The ordered pairs (2, 5) and (5, 2) are	Equal	Not equal	Same point	Both on the y-axis	○●○○
126. The set of all first elements of the ordered pairs in a relation is its	Range	Inverse	Domain	Image set only	○○●○
127. A binary relation from A to B is a	Subset of A ∩ B	Subset of A × B	Superset of A × B	Element of A	○●○○
128. If n(B) = t, the number of binary relations in B × B is	t^2	2^(t²)	2t	2^t	○●○○
129. The set of common elements of A and A^c is a	Universal set	Singleton set	Infinite set	Null set	○○○●
130. The set of all second elements of the ordered pairs in a relation is its	Domain	Range	Inverse	Origin	○●○○
131. If R = {(5, 1), (5, 2), (6, 3)}, then Range R =	{5, 6}	{1, 2}	{1, 2, 3}	{1, 2, 3, 4}	○○●○
132. If R = {(2, 0), (4, 2), (6, 4), (8, 6), (10, 8)}, then Dom R =	{0, 2, 4, 6, 8, 10}	{2, 4, 6, 8}	{2, 4, 6, 8, 10}	{0, 2, 4, 6, 8}	○○●○
133. If R = {(0, 1), (3, 4), (3, 5)}, then Range R =	{1, 4}	{1, 4, 5, 6}	{0, 3}	{1, 4, 5}	○○○●

Questions	A	B	C	D	A B C D
134. If n(A) = 3 and n(B) = 2, the number of binary relations in A × B is	128	12	32	64	○○○●
135. If A = {1, 2, 3} and B = {5}, A × B =	{(2, 5), (3, 5)}	{(1, 5), (2, 5), (3, 5)}	{(1, 5), (2, 5), (3, 5), (4, 6)}	{(5, 1), (5, 2), (5, 3)}	○●○○
136. If R = {(0, 1), (3, 4), (3, 5)}, then Dom R =	{0, 1, 3, 4, 5}	{0}	{0, 3}	{0, 3, 4}	○○●○
137. If R = {(5, 1), (6, 2), (7, 3)}, then R^{-1} =	{(1, 1), (2, 2), (3, 3)}	{(5, 5), (6, 6), (7, 7)}	{(1, 5), (2, 6), (3, 7)}	{(5, 1), (6, 2), (7, 3)}	○○●○
138. The number of subsets of a set with 4 elements is	16	8	15	32	●○○○
139. If (x + y, x − 3y) = (8, −4), then (x, y) =	(5, 2)	(6, 3)	(3, 5)	(5, 3)	○○○●
140. If n(A) = 4 and n(B) = 4, then n(A × B) =	256	16	15	8	○●○○
141. If R = {(5, 1), (5, 2), (6, 3)}, then Dom R =	{1, 2, 3}	{5, 6}	{5, 6, 7}	{1, 2, 3, 5, 6}	○●○○
142. If n(A) = 3 and n(B) = 3, the number of binary relations in A × B is	512	1024	64	256	●○○○
143. If n(A) = 2 and n(B) = 5, then n(A × B) =	32	7	10	11	○○●○
144. If n(A) = 1 and n(B) = 3, the number of binary relations in A × B is	4	8	16	3	○●○○
145. If n(A) = 5 and n(B) = 5, then n(A × B) =	25	3125	26	10	●○○○
146. If A = {1, 3}, B = {0, 2, 4}, the number of binary relations in A × B is	6	64	32	12	○●○○
147. If A = {1, 2} and B = {3, 5}, A × B =	{(3, 1), (3, 2), (5, 1), (5, 2)}	{(2, 3), (2, 5)}	{(1, 3), (1, 5), (2, 3), (2, 5)}	{(1, 1), (2, 2)}	○○●○
148. If n(C) = 5, the number of binary relations in C × C is	25	2^25	2^10	32^2	○●○○
149. If R = {(1, 2), (2, 3), (3, 4)}, then Dom R =	{2, 3, 4}	{1, 2, 3}	{1, 2, 3, 4}	{1, 2}	○●○○
150. If (x − 3y, 5x + 1) = (4, 6), then x =	1	5	−1	2	●○○○

Follow Studybeesofficial on Instagram, YouTube Channel & WhatsApp Channel — Link in Bio