


Amines

Question1

Consider the following reaction sequence




The percentage of nitrogen in product ' T ' formed is ____ %. (Nearest integer)


(Given molar mass in gmol^{-1} H : 1, C : 12, N : 14, O : 16)

JEE Main 2026 (Online) 21st January Morning Shift

Answer: 20

Solution:





$$\text{Mol. wt} = 6 \times 12 + (6 \times 1) + (2 \times 14) + (2 \times 16) = 138$$

$$\% \text{ N} = \frac{28}{138} \times 100 = 20.29\%$$

Question2


The mass of benzanilide obtained from the benzoylation reaction of 5.8 g of aniline, if yield of product is 82%, is ____ g (nearest integer).

(Given molar mass in gmol^{-1} H : 1, C : 12, N : 14, O : 16)

JEE Main 2026 (Online) 22nd January Evening Shift

Answer: 10

Solution:



Question3

Consider the following sequence of reactions.



The percentage of nitrogen in the yellow product (X) formed is ____ %.


(Nearest Integer)

(Given Molar mass in gmol^{-1} H : 1, C : 12, N : 14)

JEE Main 2026 (Online) 6th April Morning Shift

Answer: 21

Solution:



Step 1: Hydrolysis – The compound first undergoes acid-catalysed hydrolysis in the presence of HCl. This breaks the bond between the carbon and nitrogen atoms, forming the corresponding amine intermediate.

Step 2: Role of aniline – Aniline is used both as a solvent and as a reactant. It helps in dissolving the intermediate and also participates in the next reaction.

Step 3: Rearrangement on heating – When the mixture is heated gently, a rearrangement occurs, producing a new compound that leads to the formation of the yellow product.

Step 4: Isolation of the product – After the reaction finishes, the mixture is cooled and acetic acid is added. Acetic acid neutralises extra acid and helps in the precipitation of the final yellow product.

Step 5: Calculation of nitrogen percentage

The molecular mass of the product is given as 197, and the total mass of nitrogen atoms present is 42.

Hence, the percentage of nitrogen is:

$$= \frac{42}{197} \times 100 = 21.30\% \approx 21\%$$


Question4

A hydrocarbon 'P' (C_4H_8) on reaction with HCl gives an optically active compound 'Q' (C_4H_9Cl) which on reaction with one mole of ammonia gives compound 'R' ($C_4H_{11}N$). 'R' on diazotization followed by hydrolysis gives 'S'. Identify P, Q, R and S.


JEE Main 2026 (Online) 21st January Morning Shift

Options:

A.




B.



C.




D.



Answer: B

Solution:



Question5

An organic compound (P) on treatment with aqueous ammonia under hot condition forms compound (Q) which on heating with Br₂ and KOH forms compound (R) having molecular formula C₆H₇ N. Names of P, Q and R respectively are.

JEE Main 2026 (Online) 21st January Morning Shift

Options:

A.

Toluic acid, methylbenzamide, 2-methylaniline

B.

Benzoic acid, benzamide, aniline

C.

Benzoic acid, 4-methylbenzamide, 4-methylaniline

D.

Phenylethanoic acid, phenylethanamide, benzamine


Answer: B

Solution:


$$\text{C}_6\text{H}_7\text{N} \Rightarrow \text{DU} = 4$$

DU (Degree of Unsaturation) equal to 4 tells us there is one benzene ring in the molecule.

With the formula $\text{C}_6\text{H}_7\text{N}$ and one benzene ring, the compound must be aniline, which is a benzene ring attached to an NH_2 group.



Question6



Consider the above sequence of reactions. The number of bromine atom(s) in the final product (P) will be :

JEE Main 2026 (Online) 21st January Evening Shift

Options:

A.

B.

5

C.


1

D.

3

Answer: B

Solution:



Number of Br atom in major product (P) = 5

Question7


'A' is a neutral organic compound (M. F : $\text{C}_8\text{H}_9\text{ON}$). On treatment with aqueous $\text{Br}_2 / \text{HO}^{(-)}$, 'A' forms a compound 'B' which is soluble in dilute

acid. ' B ' on treatment with aqueous NaNO_2/HCl ($0 - 5^\circ\text{C}$) produces a compound ' C ' which on treatment with CuCN/NaCN produces ' D '. Hydrolysis of ' D ' produces ' E ' which is also obtainable from the hydrolysis of ' A '. ' E ' on treatment with acidified KMnO_4 produces ' F '. ' F ' contains two different types of hydrogen atoms. The structure of ' A ' is


JEE Main 2026 (Online) 22nd January Morning Shift

Options:


A.




B.



C.





D.




Answer: A

Solution:



Question8

Consider the following sequence of reactions.



Assuming that the reaction proceeds to completion, then 137 mg of 4-nitrotoluene will produce

_____ mg of *B*.

(Given molar mass in gmol^{-1} H : 1, C : 12, N : 14, O : 16, Br : 80)

JEE Main 2026 (Online) 23rd January Morning Shift

Options:

A.

208

B.

228

C.


146

D.

301

Answer: B

Solution:



$$\text{Mole} = \frac{137 \times 10^{-3}}{137} = 0.001 \text{ mole}$$

$$\text{Mole of product} = 0.001 \text{ mole}$$

$$\begin{aligned} \text{Mass of product} &= 0.001 \times 228 \text{ gm} \\ &= 0.228 \text{ gm} = 228 \text{ mg} \end{aligned}$$

Question9

A student has been given a compound " x " of molecular formula- $\text{C}_6\text{H}_7\text{N}$. ' x ' is sparingly soluble in water. However, on addition of dilute mineral acid, ' x ' becomes soluble in water. ' x ' when treated with CHCl_3 and $\text{KOH}(\text{alc})$, ' y ' is produced. ' y ' has a specific unpleasant smell. On treatment with benzenesulphonyl chloride, ' x ' gives a compound ' z ' which is soluble in alkali. The number of different " H " atoms present in ' z ' is:-

Options:

A.

8

B.

4

C.



5

D.


7

Answer: D

Solution:




Number of different H atom in



Question10

Given below are two statements :



In the light of the above statements, choose the correct answer from the options given below

JEE Main 2026 (Online) 23rd January Evening Shift

Options:

A.

Both Statement I and Statement II are true

B.

Statement I is false but Statement II is true

C.

Statement I is true but Statement II is false


D.

Both Statement I and Statement II are false


Answer: A

Solution:

Statement-I




Statement-II



Both statement (I) and (II) are true.

Question11

The correct stability order of the following diazonium salts is



JEE Main 2026 (Online) 24th January Morning Shift

Options:

A.

$C > D > B > A$

B.

$A > C > D > B$

C.

$A > B > C > D$


D.

$C > A > D > B$

Answer: B

Solution:

Correct order of stability



EDG increases stability

EWG decreases stability

Question12

A student has planned to prepare acetanilide from aniline using acetic anhydride.

The student has started from 9.3 g of aniline. However, the student has managed to obtain 11 g of dry acetanilide.

The % yield of this reaction is :-

JEE Main 2026 (Online) 24th January Evening Shift

Options:

A.

59.5%

B.

72.5

C.

97.5


D.

81.5%

Answer: D

Solution:

Reaction (1:1):



- Molar mass of aniline $\text{C}_6\text{H}_7\text{N} \approx 93 \text{ g mol}^{-1}$

$$n(\text{aniline}) = \frac{9.3}{93} = 0.10 \text{ mol}$$

- Molar mass of acetanilide $\text{C}_8\text{H}_9\text{NO} \approx 135 \text{ g mol}^{-1}$

Theoretical mass:

$$m_{\text{theor}} = 0.10 \times 135 = 13.5 \text{ g}$$

Percent yield:

$$\% \text{ yield} = \frac{11}{13.5} \times 100 \approx 81.5\%$$

Answer: Option D (81.5%)

Question 13

Given below are two statements :

Statement I : The dipole moment of R-CN is greater than R-NC and R-NC can



Statement II : R-CN hydrolyses under acidic medium to produce a compound which on treatment with SOCl_2 , followed by the addition of NH_3 gives another compound (x). This compound (x) on treatment with NaOCl/NaOH gives a product, that on treatment with $\text{CHCl}_3/\text{KOH}/\Delta$ produces R-NC

In the light of the above statements, choose the correct answer from the options given below

JEE Main 2026 (Online) 24th January Evening Shift

Options:

A.

Both Statement I and Statement II are false

B.

Statement I is false but Statement II is true

C.

Statement I is true but Statement II is false


D.

Both Statement I and Statement II are true



Answer: B

Solution:

Statement I : False



Statement II : True



Question14

Total number of alkali insoluble solid sulphonamides obtained by reaction of given amines with Hinsberg's reagent is _____.

Aniline, N-Methylaniline, Methanamine, N,N-Dimethylmethanamine, N-Methyl methanamine, Phenylmethanamine, N-propylaniline, N-phenylaniline, N,N-Dimethylaniline, Allyl amine, Isopropyl amine

JEE Main 2026 (Online) 28th January Evening Shift

Options:

A.

8

B.

5

C.

2

D.

4

Answer: D

Solution:

In Hinsberg's test, the sulphonamide formed from a 2° (secondary) amine is alkali insoluble.

This is because a 2° amine has no N-H left in the sulphonamide, so it cannot form a soluble salt with alkali.

So, we have to count the 2° amines from the given list.

Ph – NH – CH₃, Me – NH – Me,

Ph – NH – CH₂ – CH₂ – CH₃, Ph – NH – Ph

Question15

A student performed analysis of aliphatic organic compound 'X' which on analysis gave C = 61.01%, H = 15.25%, N = 23.74%.

This compound, on treatment with HNO₂/H₂O produced another compound 'Y' which did not contain any nitrogen atom. However, the compound 'Y' upon controlled oxidation produced another compound 'Z' that responded to iodoform test.

The structure of 'X' is :


JEE Main 2026 (Online) 28th January Evening Shift

Options:

A.




B.



C.




D.




Answer: D

Solution:



Question16

Consider the three aromatic molecules (P, Q and R) whose structures have been given below :



The correct order regarding the reactivity of these compounds with $\text{Ph}-\text{N} \equiv \text{NCl}^{(-)}$ under optimum but slightly acidic medium is :

JEE Main 2026 (Online) 2nd April Morning Shift

Options:

A.

$P > Q > R$

B.

$R > P > Q$

C.

$R > Q > P$


D.

$P > R > Q$

Answer: A

Question17

Product C of the following reaction sequence will be



JEE Main 2026 (Online) 4th April Morning Shift

Options:

A.

1-Bromo-4-nitrobenzene

B.

1, 3, 5-Tribromo-2-nitrobenzene

C.


4-Bromo-1-nitrobenzene

D.

1, 3, 5-Tribromobenzene

Answer: B

Solution:



Question18

Identify the incorrect statements.

- A. is a stronger base than
- B. can be synthesized by Gabriel phthalimide synthesis.
- C. $\xrightarrow{\text{Br}_2, \text{NaOH}}$ primary aromatic amine
- D. $\xrightarrow[\text{(iii) } \Delta]{\text{(i) NaNO}_2, \text{HCl}, 0^\circ\text{C}}$ product will dissolve in NaOH

Choose the correct answer from the options given below:

JEE Main 2026 (Online) 5th April Morning Shift

Options:

A.

A and D Only

B.

A and C Only

C.


B and C Only

D.


A and B Only

Answer: C






Solution:



delocalized lp is not available for donation and hence weaker base than the above compound.




Aromatic amines can't be synthesized by this process because of follows SN^2 mechanism and aryl halides are difficult to substitute due to partial double bond character.



Question19

Benzyl isocyanide can be obtained from



Choose the correct answer from the options given below :

JEE Main 2026 (Online) 5th April Morning Shift

Options:

A.

A and B Only

B.

A and C Only

C.






B and D Only

D.

D and E Only

Answer: A

Solution:



(A) & (B) gives correct product.

Question20

Given below are two statements :

Statement I : Heating benzamide with bromine in an ethanolic solution of sodium hydroxide will give benzylamine.

Statement II : Nitration of aniline with $\text{HNO}_3/\text{H}_2\text{SO}_4$ at 288 K produces *m*-nitroaniline in higher amount than *o*-nitroaniline (pH adjusted).

In the light of the above statements, choose the correct answer from the options given below :

JEE Main 2026 (Online) 5th April Evening Shift

Options:

A.

Both Statement I and Statement II are true

B.

Both Statement I and Statement II are false

C.

Statement I is true but Statement II is false


D.

Statement I is false but Statement II is true

Answer: D


Solution:

Statement-I



Hence Statement-I is incorrect

Statement-II



Statement-II is correct

Question21

Arrange the following compounds according to increasing order of boiling points.

$n - \text{C}_4\text{H}_9\text{OH}$ (A), $n - \text{C}_4\text{H}_9\text{NH}_2$ (B), $n - \text{C}_4\text{H}_{10}$ (C) and $\text{C}_2\text{H}_5\text{NHC}_2\text{H}_5$ (D).

JEE Main 2026 (Online) 6th April Morning Shift

Options:

A.

$\text{C} < \text{B} < \text{A} < \text{D}$

B.

$\text{D} < \text{C} < \text{B} < \text{A}$

C.

$\text{C} < \text{D} < \text{B} < \text{A}$

D.

$D < B < A < C$

Answer: C

Solution:

(A) $n - C_4H_9OH$ (an alcohol) forms strong intermolecular hydrogen bonding between O – H groups of different molecules. Because of this strong attraction, more heat is needed to separate the molecules, so its boiling point is the highest.

(B) $n - C_4H_9NH_2$ (a primary amine) also shows intermolecular hydrogen bonding due to the presence of N – H bonds. But nitrogen is less electronegative than oxygen, so N – H hydrogen bonding is weaker than O – H hydrogen bonding. Hence, its boiling point is lower than the alcohol.

(C) $n - C_4H_{10}$ is a non-polar hydrocarbon. It cannot form hydrogen bonds. Only weak van der Waals (London dispersion) forces act between its molecules, so it has the lowest boiling point.


(D) $C_2H_5NHC_2H_5$ (a secondary amine) can form hydrogen bonding, but it is weaker than in $n - C_4H_9NH_2$ because it has only one N – H bond (fewer chances for hydrogen bonding). So its boiling point is lower than (B), but higher than the hydrocarbon (C).

Therefore, the increasing order of boiling points is:

$C < D < B < A$

Question22

The number of compounds from the following which can undergo reaction with Br_2/KOH (alcoholic) to give respective products and these respective products can also be obtained separately by Gabriel phthalimide reaction is :



JEE Main 2026 (Online) 6th April Evening Shift

Options:

A.

5

B.

4

C.


3

D.




6

Answer: C


Solution:




Aniline is not formed by Gabriel phthalimide synthesis.



No 1° amine is form




No 1° amine is form



Question23

Consider the following organic reaction sequence. Choose the final product (X) from the following (consider the major product in all intermediate reactions)



JEE Main 2026 (Online) 6th April Evening Shift

Options:

A.

Benzene

B.

Phenol

C.


Propanol

D.

Chlorobenzene

Answer: A

Solution:



Question24

Which of the following amino acid will give violet coloured complex with neutral ferric chloride solution?

JEE Main 2026 (Online) 8th April Evening Shift

Options:

A.

Threonine

B.

Serine

C.

Tyrosine





D.


Cysteine

Answer: C

Solution:

FeCl_3 reacts with phenol to give a violet Coloured complex.





Question25

Which statements are True?

- A. In Hoffmann bromamide degradation, 4 moles of NaOH and 2 moles of Br_2 are consumed per mole of an amide
- B. Hoffmann bromamide reaction is not given by alkyl amides.
- C. Primary amines can be synthesized by Hoffmann bromamide degradation.
- D. Secondary amide on reaction with Br_2 and NaOH will give secondary amine.
- E. The by-products of Hoffmann degradation are Na_2CO_3 , NaBr and H_2O .

Choose the correct answer from the options given below :

JEE Main 2026 (Online) 8th April Evening Shift


Options:

- A.
A, C and E only
- B.
B, C and D only
- C.
C and E only
- D.
C, D and E only

Answer: C

Solution:

Hoffmann bromamide degradation reaction –



Statement (A) – False

Statement (B) – False


Statement (C) – True

Statement (D) – False

Statement (E) – True

Question26

Consider the following sequence of reactions :




Molar mass of the product formed (A) is _____ g mol⁻¹.

JEE Main 2025 (Online) 22nd January Morning Shift

Answer: 154


Solution:



Molar mass of A = $154 \frac{\text{gm}}{\text{mole}}$

Question27

Consider the following sequence of reactions to produce major product (A)




Molar mass of product is .

(Given molar mass in g mol^{-1} of C:12,H:1,O:16,Br:80, N:14,P:31)

JEE Main 2025 (Online) 23rd January Morning Shift

Answer: 171


Solution:



Molar mass of product $(\text{C}_7\text{H}_7\text{Br})(\text{A})$ is 171 g mol^{-1}

Question 28

Consider the following sequence of reactions.




Total number of sp^3 hybridised carbon atoms in the major product C formed is _____.

JEE Main 2025 (Online) 23rd January Evening Shift


Answer: 4

Solution:



Question29

Given below are some nitrogen containing compounds :






Each of them is treated with HCl separately. 1.0 g of the most basic compound will consume _____ mg of HCl .

(Given molar mass in g mol^{-1} C : 12, H : 1, O : 16, Cl : 35.5)


JEE Main 2025 (Online) 29th January Morning Shift

Answer: 341


Solution:



The basic strength order is



Most basic compound (benzyl amine) has localized electron pairs (electrons are not involved in resonance). The basicity of aniline is reduced due to the resonance delocalisation of electrons. $-\text{OCH}_3$ group is also electron withdrawing (less than $-\text{NO}_2$) and hence electron density decrease and basicity decrease.



Mass of benzyl amine given = 1.0 g

Molar mass of benzyl amine = 107.15 g mol⁻¹

Moles of benzyl amine

$$= \frac{\text{mass}}{\text{molar mass}}$$

$$= \frac{1.0 \text{ g}}{107.15 \text{ g mol}^{-1}}$$

$$= 0.0093327 \text{ mol}$$

The stoichiometric ratio between both the reactants is 1 : 1.

Benzyl amine : HCl

1 : 1

So, moles of HCl = 0.0093327 mol

Mass of HCl consumed by benzyl amine is calculated as:

Mass = moles × molar mass

Molar mass of HCl = 36.5 g mol⁻¹

Mass = 0.0093327 mol × 36.5 g mol⁻¹


= 0.34064355 g

= 340.64355 mg

= 340.6 mg ≈ 341 mg






Question30

The products formed in the following reaction sequence are :




JEE Main 2025 (Online) 22nd January Morning Shift

Options:

- A.  , 
- B.  , $\text{CH}_3 - \text{COOH}$
- C.  , $\text{CH}_3 - \text{CHO}$
- D.  , $\text{CH}_3 - \text{CHO}$

Answer: D



Solution:



Question31

Match the Compounds (List - I) with the appropriate Catalyst/ Reagents (List - II) for their reduction into corresponding amines.

	List - I (Compounds)		List - II (Catalyst/Reagents)
(A)	$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$	(I)	NaOH (aqueous)

	List - I (Compounds)		List - II (Catalyst/Reagents)
(B)		(II)	H ₂ /Ni
(C)	R – C ≡ N	(III)	LiAlH ₄ , H ₂ O
(D)		(IV)	Sn, HCl

Choose the correct answer from the options given below :




JEE Main 2025 (Online) 22nd January Evening Shift

Options:

- A. (A)-(II), (B)-(IV), (C)-(III), (D)-(I)
- B. (A)-(III), (B)-(IV), (C)-(II), (D)-(I)
- C. (A)-(III), (B)-(II), (C)-(IV), (D)-(I)
- D. (A)-(II), (B)-(I), (C)-(III), (D)-(IV)





Answer: B

Solution:



Question 32

Which among the following react with Hinsberg's reagent?



Choose the correct answer from the options given below :

JEE Main 2025 (Online) 23rd January Morning Shift

Options:

A. A, C and E only

B. A, B and E only

C. C and D only

D. B and D only

Answer: A

Solution:

B and D are 3° amine which does not have replaceable H on N, So does not react.

Question33

Given below are two statements I and II.

Statement I : Dumas method is used for estimation of "Nitrogen" in an organic compound.

Statement II : Dumas method involves the formation of ammonium sulphate by heating the organic compound with conc H_2SO_4 .

In the light of the above statements, choose the correct answer from the options given below

JEE Main 2025 (Online) 24th January Morning Shift

Options:

A. Statement I is true but Statement II is false

B. Statement I is false but Statement II is true

C. Both Statement I and Statement II are true

D. Both Statement I and Statement II are false


Answer: A

Solution:

In Dumas method nitrogen present in organic compound is converted into N_2 gas whose volumetric analysis gives the percentage of nitrogen atom in the organic compound.

Question34

For reaction



The correct order of set of reagents for the above conversion is :


JEE Main 2025 (Online) 24th January Evening Shift

Options:

- A. Ac_2O , Br_2 , $H_2O(\Delta)$, $NaOH$
- B. H_2SO_4 , Ac_2O , Br_2 , $H_2O(\Delta)$, $NaOH$
- C. Br_2 | $FeBr_3$, $H_2O(\Delta)$, $NaOH$
- D. Ac_2O , H_2SO_4 , Br_2 , $NaOH$

Answer: B

Solution:



Question35

Identify correct statements :

- (A) Primary amines do not give diazonium salts when treated with NaNO_2 in acidic condition.
- (B) Aliphatic and aromatic primary amines on heating with CHCl_3 and ethanolic KOH form carbylamines.
- (C) Secondary and tertiary amines also give carbylamine test.
- (D) Benzenesulfonyl chloride is known as Hinsberg's reagent.
- (E) Tertiary amines reacts with benzenesulfonyl chloride very easily.

Choose the correct answer from the options given below :

Options:

A.

(B) and (C) only

B.


(D) and (E) only

C.

(A) and (B) only

D.

(B) and (D) only

Answer: D**Solution:**

(B)







(C) Only primary amine gives carbyl amine test

(D)

(E) Tertiary amine do not react with $\text{Ph} - \text{SO}_2\text{Cl}$

So correct options are (B) and (D) only

Question36**The steam volatile compounds among the following are :**



Choose the correct answer from the options given below :

JEE Main 2025 (Online) 29th January Morning Shift

Options:

A.

(B) and (D) Only

B.

(A), (B) and (C) Only

C.

(A) and (B) Only

D.

(A) and (C) Only

Answer: C

Solution:


Steam volatile compounds can be distilled using steam distillation. Steam volatility is influenced by the intermolecular forces present in the substance. Steam volatile compounds have lower boiling points and weaker intermolecular forces.

A compound's steam volatility is directly related to hydrogen bonding.


The presence of intramolecular hydrogen bonds make a molecule more volatile. A compound's intermolecular hydrogen bonding decrease its volatility.

So, molecules with intramolecular hydrogen bonds are more likely to be steam volatile:

(A)




Intramolecular hydrogen bonding is present in this molecule.



So, it is steam volatile


(B)



The —NH_2 group and —NO_2 group are at ortho positions (adjacent positions) and hence there is intramolecular hydrogen bonding.


The electronegative oxygen of —NO_2 group and hydrogen of —NH_2 group makes the intramolecular hydrogen bonding. Due to this, intramolecular hydrogen bonding, O-nitroaniline is considered as steam volatile.

(C)





The amino group and hydroxy group are at para positions and the possibility of intramolecular hydrogen bonding is less in this compound. Therefore, this compound is not considered as steam volatile.

(D)



The —NH_2 group and group are far away to form intramolecular hydrogen bonding. As a result, it is not considered as steam volatile.


The steam volatile compounds are



Correct answer option (3) (A) and (B) only.

Question 37

In the following substitution reaction :




Product 'P' formed is :


JEE Main 2025 (Online) 29th January Morning Shift

Options:


A.




B.



C.




D.




Answer: D

Solution:





The reaction is an example of nucleophilic aromatic substitution reaction.



The ethoxide ion ($\text{C}_2\text{H}_5\text{O}^-$) replaces a bromine atom on the aromatic ring.


The nitro group ($-\text{NO}_2$) activates the benzene ring towards nucleophilic aromatic substitution. The Br atom para to the $-\text{NO}_2$ group is more susceptible to substitution due to resonance stabilisation of the intermediate carbanion. Therefore, the $\text{C}_2\text{H}_5\text{O}^-$ ion replaces the Br atom para to the nitro group.



(Product)

Correct answer:

Option (4)







Question38

Which one of the following reaction sequences will give an azo dye?

JEE Main 2025 (Online) 29th January Evening Shift

Options:









Answer: C

Solution:

Azo dyes are organic compounds that contain an azo bond ($-N=N-$)


(1)



There is no azo bond $-N=N-$ formation in this case.

So, azo dye compound is not formed


(2)



Azo dye ($-N=N-$) is not formed in this reaction.

Benzonitrile reaction with H_2SO_4 , gives benzoic acid product. $-CN$ becomes $-COOH$. Then, reaction with PCl_5 gives benzoyl chloride (C_6H_5COCl). Its on reaction with aniline does not give $-N=N-$ type compound. So, azo dye compound is not formed.

(3)





Nitrobenzene on reaction with Sn/HCl undergoes reduction reaction to form aniline. $-NO_2$ is reduced to $-NH_2$.



Aniline on reaction with $NaNO_2/HCl$ undergoes diazotization reaction to form benzene diazonium chloride.

Nitrobenzene on reaction with Sn/HCl undergoes reduction reaction to form aniline. $-NO_2$ is reduced to $-NH_2$.

Aniline on reaction with $NaNO_2/HCl$ undergoes diazotization reaction to form benzene diazonium chloride.




Benzene diazonium chloride (Benzene diazonium salt) on reaction with β -naphthol and $NaOH$ gives an azo compound.



Azo dye is formed in this reaction

(4)



Azo dye is not formed.

Question39

The correct order of basic nature in aqueous solution for the bases

NH_3 , $\text{H}_2\text{N}-\text{NH}_2$, $\text{CH}_3\text{CH}_2\text{NH}_2$, $(\text{CH}_3\text{CH}_2)_2\text{NH}$ and $(\text{CH}_3\text{CH}_2)_3\text{N}$ is :

JEE Main 2025 (Online) 2nd April Morning Shift

Options:

- A. $\text{NH}_3 < \text{H}_2\text{N}-\text{NH}_2 < (\text{CH}_3\text{CH}_2)_3\text{N} < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH}$
- B. $\text{H}_2\text{N}-\text{NH}_2 < \text{NH}_3 < (\text{CH}_3\text{CH}_2)_3\text{N} < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH}$
- C. $\text{NH}_2-\text{NH}_2 < \text{NH}_3 < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_3\text{N} < (\text{CH}_3\text{CH}_2)_2\text{NH}$
- D. $\text{NH}_3 < \text{H}_2\text{N}-\text{NH}_2 < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH} < (\text{CH}_3\text{CH}_2)_3\text{N}$

Answer: C

Solution:

Basic strength of amine depends on hydrogen bonding and electronic inductive effect.



Question40

When a concentrated solution of sulphanilic acid and 1-naphthylamine is treated with nitrous acid (273 K) and acidified with acetic acid, the mass (g) of 0.1 mole of product formed is :

(Given molar mass in gmol^{-1} H : 1, C : 12, N : 14, O : 16, S : 32)


JEE Main 2025 (Online) 2nd April Evening Shift


Options:

- A. 330
- B. 33
- C. 343
- D. 66

Answer: B

Solution:







0.1 mole of red-azo dye (Molar Mass = 327gm/mol) will have 32.7 gm mass. Nearly 33 gm .



Question41

In the following reactions, which one is NOT correct?

JEE Main 2025 (Online) 3rd April Morning Shift


Options:






Answer: C

Solution:



Question42


Identify [A],[B] and [C], respectively in the following reaction sequence :




JEE Main 2025 (Online) 3rd April Morning Shift

Options:


A.




B.



C.




D.



Answer: C

Solution:







Question43

The sequence from the following that would result in giving predominantly 3, 4, 5– Tribromoaniline is :

JEE Main 2025 (Online) 3rd April Evening Shift


Options:






Answer: C

Solution:



Question44


The major product (A) formed in the following reaction sequence is




JEE Main 2025 (Online) 4th April Morning Shift

Options:


A.




B.



C.




D.






Answer: B


Solution:



Question45

Which of the following amine (s) show (s) positive carbylamine test?





Choose the correct answer from the options given below:

JEE Main 2025 (Online) 7th April Morning Shift


Options:

- A. C Only
- B. B, C and D Only
- C. A and E Only
- D. A and C Only

Answer: D

Solution:





Only 1° or primary amines gives positive carbylamines test.



Option (A) and (C) are primary amine and given $\oplus^{\text{ve}}$ carbyl amine test

Question46

Match List - I with List - II.

List - I (Conversion)	List - II (Reagents, Conditions used)
(A) 	(I) Warm H ₂ O
(B) 	(II) (a) NaOH, 368K; (b) H ₃ O ⁺
(C) 	(III) (a) NaOH, 443K; (b) H ₃ O ⁺
(D) 	(IV) (a) NaOH, 623K, 300 atm; (b) H ₃ O ⁺

Choose the correct answer from the options given below :

JEE Main 2025 (Online) 7th April Evening Shift

Options:

A.

(A)-(II), (B)-(III), (C)-(I), (D)-(IV)

B.

(A)-(III), (B)-(IV), (C)-(II), (D)-(I)

C.

(A)-(IV), (B)-(III), (C)-(I), (D)-(II)

D.

(A)-(IV), (B)-(III), (C)-(II), (D)-(I)

Answer: D

Solution:

Aromatic halide give nucleophilic substitution reaction at high temperature or in presence of $-I/-M$ group rate of reaction high even at low temperature.


A-IV

B-III

C-II

D-I

Question47



'A' shows positive Lassaign's test for N and its molar mass is 121.

'B' gives effervescence with aq NaHCO_3 .


'C' gives fruity smell.

Identify A, B and C from the following.


JEE Main 2025 (Online) 8th April Evening Shift

Options:


A.




B.



C.




D.



Answer: C

Solution:



Molar mass = 121


A $\rightarrow$ Benzamide Shows positive Lassaigh's test.

B $\rightarrow$ Benzoic acid gives effervescence with aq. NaHCO_3 .

C $\rightarrow$ Ester gives fruity smell.

Question48


The product A formed in the following reaction is:




[29-Jan-2024 Shift 2]

Options:


A.




B.



C.




D.



Answer: C

Solution:




Question49


Which of the following reaction is correct?
[29-Jan-2024 Shift 2]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question 50


Following is a confirmatory test for aromatic primary amines. Identify reagent (A) and (B)




[30-Jan-2024 Shift 1]

Options:


A.



B.




C.




D.

A = $\text{NaNO}_2 + \text{HCl}$, $0 - 5^\circ\text{C}$;




Answer: D

Solution:



Question51


The products A and B formed in the following reaction scheme are respectively




[30-Jan-2024 Shift 2]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question52

Given below are two statements:

Statement I: Aniline reacts with $\text{con. H}_2\text{SO}_4$ followed by heating at $453 - 473\text{K}$ gives p aminobenzene sulphonic acid, which gives blood red colour in the 'Lassaigne's test'.

Statement II: In Friedel - Craft's alkylation and acylation reactions, aniline forms salt with the AlCl_3 catalyst. Due to this, nitrogen of aniline acquires a positive charge and acts as deactivating group.

In the light of the above statements, choose the correct answer from the options given below :


[31-Jan-2024 Shift 2]

Options:

- A. Statement I is false but statement II is true
- B. Both statement I and statement II are false
- C. Statement I is true but statement II is false
- D. Both statement I and statement II are true


Answer: D

Solution:



Question53

Choose the correct colour of the product for the following reaction.




[24-Jan-2023 Shift 2]

Options:

- A. Yellow
- B. White
- C. Red
- D. Blue

Answer: C

Solution:



Question54

Given below are two statements:

Statement I : Pure Aniline and other arylamines are usually colourless.

Statement II : Arylamines get coloured on storage due to atmospheric reduction.

In the light of the above statements, choose the most appropriate answer from the options given below :

[24-Jan-2023 Shift 2]

Options:

- A. Both Statement I and Statement II are incorrect
- B. Both Statement I and Statement II are correct
- C. Statement I is correct but Statement II is incorrect
- D. Statement I is incorrect but Statement II is correct

Answer: C

Solution:

Statement – 1 is (True)


Pure aniline is colourless liquid

Statement - 2 is (False)

Aniline becomes dark brown due to action of air and light [oxidation]

Question55


Identify the product formed (A and E)




[25-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.



Answer: B

Solution:



Question 56

The correct order in aqueous medium of basic strength in case of methyl substituted amines is :
[25-Jan-2023 Shift 1]

Options:

- A. $\text{Me}_2\text{NH} > \text{MeNH}_2 > \text{Me}_3\text{N} > \text{NH}_3$
- B. $\text{Me}_2\text{NH} > \text{Me}_3\text{N} > \text{MeNH}_2 > \text{NH}_3$
- C. $\text{NH}_3 > \text{Me}_3\text{N} > \text{MeNH}_2 > \text{Me}_2\text{NH}$
- D. $\text{Me}_3\text{N} > \text{Me}_2\text{NH} > \text{MeNH}_2 > \text{NH}_3$

Answer: A

Solution:

In aqueous medium basic strength is dependent on electron density on nitrogen as well as solvation of cation formed after accepting H^+ . After considering all these factors overall basic strength order is $\text{Me}_2\text{NH} > \text{MeNH}_2 > \text{Me}_3\text{N} > \text{NH}_3$

Question 57


Reaction of propanamide with Br_2/KOH (aq) produces:
[29-Jan-2023 Shift 2]

Options:

- A. Ethylnitrile
- B. Propylamine
- C. Propanenitrile
- D. Ethylamine

Answer: D

Solution:



Question58

Match List I with List II

List I		List II	
A	$\text{C}_6\text{H}_5\text{Cl} + \text{CH}_3\text{Cl} \xrightarrow{\text{Na}} \text{C}_6\text{H}_5\text{CH}_3$	I	Fitting reaction
B	$\text{C}_6\text{H}_5\text{Cl} + 2\text{Na} \rightarrow \text{C}_6\text{H}_5\text{C}_6\text{H}_5$	II	Wurtz Fitting reaction
C	$\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^- \xrightarrow{\text{Cu}_2\text{Cl}_2} \text{C}_6\text{H}_5\text{Cl} + \text{N}_2$	III	Finkelstein reaction
D	$\text{C}_2\text{H}_5\text{Cl} + \text{NaI} \rightarrow \text{C}_2\text{H}_5\text{I} + \text{NaCl}$	IV	Sandmeyer reaction




[30-Jan-2023 Shift 1]

Options:

- A. A - II, B - I, C - III, D - IV
- B. A - III, B - II, C - IV, D - I
- C. A - IV, B - II, C - III, D - I
- D. A - II, B - I, C - IV, D - III





Answer: D

Solution:

LIST-I		LIST-II
A.		Wurtz-fitting reaction
B.		Fitting reaction
C.		Sandmeyer reaction
D.	$\text{C}_2\text{H}_5\text{Cl} + \text{NaI} \rightarrow \text{C}_2\text{H}_5\text{I} + \text{NaCl}$	Finkelstein reaction

Question59

Benzyl isocyanide can be obtained by :

- (A) 
- (B) 
- (C) 
- (D) 

Choose the correct answer from the options given below :
[30-Jan-2023 Shift 1]

Options:

A. A and D


B. Only B

C. A and B


D. B and C

Answer: C

Solution:



Question60



In the above conversion of compound (X) to product (Y), the sequence of reagents to be used will be:


[30-Jan-2023 Shift 2]

Options:


- A. (i) Br_2 , Fe (ii) Fe, H^+ (iii) LiAlH_4
- B. (i) Br_2 (aq) (ii) LiAlH_4 (iii) H_3O^+
- C. (i) Fe, H^+ (ii) Br_2 (aq) (iii) HNO_2 (iv) CuBr
- D. (i) Fe, H^+ (ii) Br_2 (aq) (iii) HNO_2 (iv) H_3PO_2

Answer: D

Solution:




Question61




[31-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question62

How many of the transformation given below would result in aromatic amines?


[31-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.




Answer: C

Solution:

Product in the given reactions are as follow-




2. No reactions will be observed as in Gabriel phthalimide synthesis




is poor substrate for S_N^2

3.



4.




+ CH_3COOH

Aromatic amines will be formed in 1, 3 & 4

Ans: 3


Question63

In the following reaction, ' A ' is




'A' Major product.
[1-Feb-2023 Shift 1]


Options:




B.



D.




D.



Answer: B


Solution:

Initially lone pair electron of -NH_2 attack on electrophilic carbon, after then lone pair electron of oxygen attacks leading to formation of cyclic compound.



Question64


The major products A and B from the following reactions are:




[6-Apr-2023 shift 1]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question65


The major product formed in the following reaction is




[6-Apr-2023 shift 1]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question66




Compound P is neutral. Q gives effervescence with NaHCO_3 while R reacts with Hinsbergs reagent to give solid soluble in NaOH.

Compound P is


[6-Apr-2023 shift 1]

Options:


A.




B.



C.




D.



Answer: B

Solution:






Question67

Number of isomeric aromatic amines with molecular formula $C_8H_{11}N$, which can be synthesized by Gabriel Phthalimide synthesis is ____
[6-Apr-2023 shift 2]

Answer: 6


Solution:



(d + 1)

Question 68

Match List I with List II:



is reacted with reagents in List I to form products in List II.

List I (Reagent)	List II (Product)
A. $\text{C}_6\text{H}_5\text{NH}_2$	I. $\text{C}_6\text{H}_5\text{I}$
B. HBF_4, Δ	II. $\text{C}_6\text{H}_5\text{CN}$
C. Cu, HCl	III. $\text{C}_6\text{H}_5\text{N}=\text{N}-\text{C}_6\text{H}_4\text{NH}_2$
D. CuCN/KCN	IV. $\text{C}_6\text{H}_5\text{Cl}$

Choose the correct answer from the options given below:

[8-Apr-2023 shift 1]

Options:

A. A-I, B-III, C-IV, D-II


B. A-III, B-I, C-II, D-IV

C. A-III, B-I, C-IV, D-II

D. A-IV, B-III, C-II, D-I


Answer: C

Solution:



Question69


The product (P) formed from the following multistep reaction is :




[8-Apr-2023 shift 2]

Options:


A.




B.



C.




D.



Answer: D

Solution:



Question 70

Isomeric amines with molecular formula $C_8H_{11}N$ given the following tests

Isomer (P) $\Rightarrow$ Can be prepared by Gabriel phthalimide synthesis


Isomer (Q) $\Rightarrow$ Reacts with Hinsberg's reagent to give solid insoluble in NaOH

Isomer (R) $\Rightarrow$ Reacts with HONO followed by β -naphthol in NaOH to give red dye.


Isomer (P), (Q) and (R) respectively are
[10-Apr-2023 shift 1]

Options:


A.




B.



C.



D.




Answer: B

Solution:

P = Can be prepared by Gabriel phthalimide synthesis it should be 1^o-amine

Q = React with Hinsberg's reagent and insoluble in NaOH it should be 2^o-amine

R = React with HNO_2 followed by B-Napthol in NaOH it give red dye it must be Aromatic Amine




Question71


o-Phenylenediamine $\xrightarrow{\text{HNO}_2}$ **' X ' Major Product ' X ' is**
 [11-Apr-2023 shift 1]

Options:


A.




B.



C.




D.



Answer: C


Solution:

o-Phenylenediamine



Question 72


Compound ' B ' is




[11-Apr-2023 shift 2]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question73

In the reaction given below



'A' is

[13-Apr-2023 shift 1]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question74

Compound A from the following reaction sequence is :




[13-Apr-2023 shift 2]

Options:

- A. Phenol
- B. Benzoic Acid
- C. Aniline
- D. Salicylic Acid


Answer: C

Solution:



Question75


Consider the following sequence of reaction :




The product ' B ' is :
[15-Apr-2023 shift 1]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question 76



In the above conversion the correct sequence of reagents to be added is :

Options :


[15-Apr-2023 shift 1]

Options:

- A. (i) Br_2/Fe , (ii) Fe/H^+ , (iii) KMnO_4 , (iv) Cl_2
- B. (i) Br_2/Fe , (ii) Fe/H^+ , (iii) HONO , (iv) CuCl , (v) KMnO_4
- C. (i) Fe/H^+ , (ii) HONO , (iii) CuCl , (iv) KMnO_4 , (v) Br_2
- D. (i) KMnO_4 , (ii) Br_2/Fe , (iii) Fe/H^+ , (iv) Cl_2 ,


Answer: B

Solution:



Question77


In the given reaction sequence, the major product 'C' is :




[24-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: B

Solution:

$$\text{C}_8\text{H}_{10} \text{ DU} = 9 - 5 = 4$$



Question 78


The conversion of propan-1-ol to n-butylamine involves the sequential addition of reagents. The correct sequential order of reagents is
[24-Jun-2022-Shift-2]

Options:

- A. (i) SOCl_2 (ii) KCN (iii) H_2/Ni , $\text{Na(Hg)}/\text{C}_2\text{H}_5\text{OH}$
- B. (i) HCl (ii) H_2/Ni , $\text{Na(Hg)}/\text{C}_2\text{H}_5\text{OH}$
- C. (i) SOCl_2 (ii) KCN (iii) CH_3NH_2
- D. (i) HCl (ii) CH_3NH_2


Answer: A

Solution:



Question 79

The reaction of



with bromine and KOH gives RNH_2 as the end product. Which one of the following is the intermediate product formed in this reaction?
[25-Jun-2022-Shift-1]


Options:

- A. $\text{R} - \underset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{NH} - \text{Br}$
- B. $\text{R} - \text{NH} - \text{Br}$
- C. $\text{R} - \text{N} = \text{C} = \text{O}$
- D. $\text{R} - \underset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{NBr}_2$

Answer: C

Solution:

The given reaction is Hoffmann-Bromide degradation method.



Question80

During halogen test, sodium fusion extract is boiled with concentrated HNO_3 to

[25-Jun-2022-Shift-2]

Options:

- A. remove unreacted sodium
- B. decompose cyanide or sulphide of sodium
- C. extract halogen from organic compound
- D. maintain the pH of extract.

Answer: B

Solution:


During test for halogen, if nitrogen or sulphur is also present in the compound, then sodium fusion extract is first boiled with concentrated nitric acid to decompose cyanide or sulphide of sodium formed during Lassaigne's test.

Question81


Which of the following ketone will NOT give enamine on treatment with secondary amines? [where t-Bu is $\text{.C(CH}_3\text{)}_3$]
[25-Jun-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: C

Solution:

Enamine formation is an example of nucleophilic addition elimination reaction. Since in ketone



Group is highly sterically hindered hence attack of nucleophile will not be possible.


Question82

Among the following structures, which will show the most stable enamine formation? (Where Me is --CH_3)


[26-Jun-2022-Shift-1]

Options:


A.




B.



C.



D.



Answer: C

Solution:

All these enamines are interconvertible through their resonating structures. So most stable form is 'C' due to steric factor.

Question83


**Which statement is NOT correct for p-toluene sulphonyl chloride?
[26-Jun-2022-Shift-2]**

Options:




- A. It is known as Hinsberg's reagent.
- B. It is used to distinguish primary and secondary amines.
- C. On treatment with secondary amine, it leads to a product, that is soluble in alkali.
- D. It doesn't react with tertiary amines.

Answer: C

Solution:




Hinsberg's reagent



Question84


The final product ' C ' in the following series of reactions




[26-Jun-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question85

Given below are two statements :

Statement I : In Hofmann degradation reaction, the migration of only an alkyl group takes place from carbonyl carbon of the amide to the nitrogen atom.

Statement II : The group is migrated in Hofmann degradation reaction to electron deficient atom.

In the light of the above statements, choose the most appropriate answer from the options given below:

[27-Jun-2022-Shift-1]

Options:

A. Both Statement I and Statement II are correct.

- B. Both Statement I and Statement II are incorrect.
- C. Statement I is correct but Statement II is incorrect.
- D. Statement I is incorrect but Statement II is correct.

Answer: D

Solution:


Hofmann bromamide degradation

In this degradation, the migration of the alkyl/aryl group occurs to the electron-deficient nitrogen (nitrene). Statement (I) is not absolutely correct as it mentions only the alkyl group, whereas migration of aryl groups may also occur depending on migratory aptitude.

Statement (II) is correct as the migration occurs to the electron-deficient atom.

Question86


Identify the major product formed in the following sequence of reactions:




[28-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question87


A primary aliphatic amine on reaction with nitrous acid in cold (273K) and there after raising temperature of reaction mixture to room temperature (298K), gives a/an
[28-Jun-2022-Shift-1]

Options:

- A. nitrile
- B. alcohol
- C. diazonium salt
- D. secondary amine


Answer: B

Solution:



Question88

With respect to the following reaction, consider the given statements :



- (A) o-Nitroaniline and p-nitroaniline are the predominant products.
- (B) p-Nitroaniline and m-nitroaniline are the predominant products.
- (C) HNO_3 acts as an acid.
- (D) H_2SO_4 acts as an acid.

Choose the correct option.

[28-Jun-2022-Shift-2]

Options:

A. (A) and (C) are correct statements.



B. (A) and (D) are correct statements.

C. (B) and (D) are correct statements.


D. (B) and (C) are correct statements.

Answer: C

Solution:




Question89




Consider the above reactions, the product A and product B respectively are
[29-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.




Answer: C

Solution:




Question90




The correct structure of
[25-Jul-2022-Shift-2]



Options:



B.




C.



Answer: A

Solution:



Question91

Given below are two statements : one is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A) : Experimental reaction of CH_3Cl with aniline and anhydrous AlCl_3 does not give o and p methylaniline.

Reason (R): The $-\text{NH}_2$ group of aniline becomes deactivating because of salt formation with anhydrous AlCl_3 and hence yields m-methyl aniline as the product.

In the light of the above statements, choose the most appropriate answer from the options given below :


[26-Jul-2022-Shift-1]

Options:

- A. Both A and R are true and (R) is the correct explanation of (A).
- B. Both A and R are true but (R) is not the correct explanation of (A).
- C. (A) is true, but (R) is false.
- D. (A) is false, but (R) is true.

Answer: C

Solution:



Aniline does not undergo Friedel Craft reaction because the reagent AlCl_3 being electron deficient acts as a Lewis acid.

Question92


Hydrolysis of which compound will give carbolic acid?
[26-Jul-2022-Shift-2]

Options:

- A. Cumene
- B. Benzenediazonium chloride
- C. Benzal chloride
- D. Ethylene glycol ketal


Answer: B

Solution:



Question93

The correct sequential order of the reagents for the given reaction is




[26-Jul-2022-Shift-2]

Options:

- A. HNO_2 , Fe/H^+ , HNO_2 , KI , $\text{H}_2\text{O}/\text{H}^+$
- B. HNO_2 , KI , Fe/H^+ , HNO_2 , $\text{H}_2\text{O}/\text{warm}$
- C. HNO_2 , KI , HNO_2 , Fe/H^+ , $\text{H}_2\text{O}/\text{H}^+$
- D. HNO_2 , Fe/H^+ , KI , HNO_2 , $\text{H}_2\text{O}/\text{warm}$

Answer: B

Solution:



Question94

Match List I with List II.

List I	List II
(A) Benzenesulphonyl chloride	(I) Test for primary amines
(B) Hoffmann bromamide reaction	(II) Anti Saytzeff
(C) Carbylamine reaction	(III) Hinsberg reagent
(D) Hoffmann orientation	(IV) Known reaction of Isocyanates.

Choose the correct answer from the options given below:
[27-Jul-2022-Shift-1]

Options:

A. A-IV, B-III, C-II, D-I

B. A-IV, B-II, C-I, D-III

C. A-III, B-IV, C-I, D-II

D. A-IV, B-III, C-I, D-II

Answer: C

Solution:

(A) Benzene sulphonyl chloride is also known as Hinsberg reagent.

(B) Hoffmann bromamide reaction involves conversion of amide to amine having one C atom less. This reaction involves isocyanate as intermediate.

(C) Carbylamine reaction is a test given by all primary amines.

(D) Hoffmann orientation refers to the addition of molecules to unsymmetrical alkenes according to anti Saytzeff's rule.

Correct match is

A-III, B-IV, C-I, D-II


Question95

An organic compound 'A' contains nitrogen and chlorine. It dissolves readily in water to give a solution that turns litmus red. Titration of compound 'A' with standard base indicates that the molecular weight of 'A' is 131 ± 2 . When a sample of 'A' is treated with aq. NaOH, a liquid separates which contains N but not Cl. Treatment of the obtained liquid with nitrous acid followed by phenol gives orange precipitate. The


compound 'A' is :
[27-Jul-2022-Shift-2]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question96

Identify the correct statement for the below given transformation.




[28-Jul-2022-Shift-1]

Options:

- A. A – $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-\text{CH}_3$, B – $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$, Saytzeff products
- B. A – $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-\text{CH}_3$, B – $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$, Hofmann products
- C. A – $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$, B – $\text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$, Hofmann products
- D. A – $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$, B – $\text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$, Saytzeff products

Answer: C

Solution:



Question97

Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R

Assertion A : Aniline on nitration yields ortho, meta & para nitro derivatives of aniline.

Reason R : Nitrating mixture is a strong acidic mixture.

In the light of the above statements, choose the correct answer from the options given below


[28-Jul-2022-Shift-2]

Options:

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true but R is NOT the correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A

Solution:




Due to formation of anilinium ion in acidic medium meta product is also obtained in significant amount

Question98


Which among the following is the strongest Bronsted base?
[29-Jul-2022-Shift-1]

Options:


A.




B.



C.




D.




Answer: D

Solution:



is the strongest base among the given compounds due to the maximum +I effect and the lone pair of N is not in dynamic state so it can be donated easily.


Question99




Consider the above reaction, the compound A' is :
[29-Jul-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: C


Question100




Which among the following represent reagent 'A'?
[29-Jul-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: A

Solution:




Question101


The Hinsberg reagent is
[29-Jul-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: A



Solution:



B.S.C (Benzene sulphonyl chloride) is known as Hinsberg Reagent.

Question102

Which of the following reaction(s) will not give p-amino azobenzene?



[25 Feb 2021 Shift 1]


Options:

- A. A only
- B. C only
- C. B only
- D. A and B


Answer: C

Solution:

Nitrobenzene in presence of NaBH_4 , NaOH and aniline will not give *p*-amino azobenzene.



In case of (A) and (C), coupling reaction takes place as the medium is quite acidic follows




Question103

The total number of amines among the following which can be synthesised by Gabriel synthesis is
[24 Feb 2021 Shift 2]


Options:

A.




B. $\text{CH}_3\text{CH}_2\text{NH}_2$

C.



D.



Answer: C


Solution:

Gabriel phthalimide synthesis is used to prepare

1° aliphatic or alicyclic amine. Hence, amine which can be synthesised by Gabriel phthalimide synthesis method contains α -carbon.

Aniline ($C_6H_5NH_2$) does not contain α -C cannot be prepared by Gabriel reaction.

Rest amines all contain α -C in its respective position, hence they can easily give Gabriel phthalimide reaction.



$\therefore$ Three amines out of given four amines can be prepared by Gabriel synthesis.

Question 104

A. phenyl methanamine

B. N,N-dimethylaniline

C. N-methyl aniline

D. Benzenamine

Choose the correct order of basic nature of the above amines.

[26 Feb 2021 Shift 2]

Options:

A. $A > C > B > D$

B. $D > C > B > A$


C. $D > B > C > A$

D. $A > B > C > D$

Answer: D


Solution:

In phenyl methanamine, the lone pair on nitrogen of -NH_2 group is localised and does not undergoes resonance as attached to sp^3 -hybridised C-atom.
Phenyl methanamine or benzyl amine




Benzyl group (Aliphatic)

(B), (C) and (D) are aromatic amines in which lone pair of electrons of N-atoms goes in resonance (+ R effect) with the benzene ring. So, Lewis basicity or donation of lone pair of electrons of these amines will be decreased in comparison to (A).




+I effects of two -CH_3 groups are in compensation with -R effect of the ring.



N-methyl aniline (2° -amine)

+I effect of one -CH_3 group is trying to compensate -R effect of the ring.



Benzenamine or aniline (1° amine)

It has no +I effect on N-atom to outweigh -R -effect of the ring.

So, A is purely aliphatic 1° -amine. B is aromatic 3° -amine with more aliphatic nature (for two -CH_3 groups). C is aromatic 2° -amine with less aliphatic nature (for one -CH_3 group).

D is purely aromatic 1° -amine. Hence basicity order is $A > B > C > D$.


Question105

Carbylamines test is used to detect the presence of primary amino group in an organic compound. Which of the following compound is formed when this test is performed with aniline?

[25 Feb 2021 Shift 2]

Options:

A.




Correct statement about the given chemical reaction is [25 Feb 2021 Shift 2]

Options:

- A. - "N H₂ group is ortho and para directing, so product (B) is not possible.
- B. reaction is possible and compound (B) will be the major product.
- C. the reaction will form sulphonated product instead of nitration.
- D. reaction is possible and compound (A) will be major product.

Answer: A

Solution:



During nitration, in strongly acidic medium, aniline gets protonated to form the anilinium ion i.e., (C₆H₅ - NH₃⁺) which is meta-directing as -NH₃⁺ is an electron withdrawing (-I) group. As a result, we get meta-nitro aniline (B) as the major product.


Question107

The diazonium salt of which of the following compounds will form a coloured dye on reaction with β -naphthol in NaOH ?


[24 Feb 2021 Shift 2]

Options:


A.




B.



C.




D.



Answer: C


Solution:

Initially aniline reacts with diazotisation reagent to form diazonium salt. Then β -naphthol react with salt and orange-red dye is obtained. So, diazonium salt of aniline is used to prepare orange-red dye.



Question108

What is the correct sequence of reagents used for converting nitrobenzene into m-dibromo benzene?



[24 Feb 2021 Shift 2]

Options:


- A. $\xrightarrow{\text{NaNO}_2} \text{ / } \xrightarrow{\text{H a}} \text{ / } \xrightarrow{\text{K Br}} \text{ / } \xrightarrow{\text{H}^+}$
- B. $\xrightarrow{\text{Br}_2/\text{Fe}} \text{ / } \xrightarrow{\text{Sn/HCl}} \text{ / } \xrightarrow{\text{NaNO}_2/\text{HCl}} \text{ / } \xrightarrow{\text{CuBrHBr}}$
- C. $\xrightarrow{\text{Sn/HCl}} \text{ / } \xrightarrow{\text{K Br}} \text{ / } \xrightarrow{\text{Br}_2} \text{ / } \xrightarrow{\text{H}^+}$
- D. $\xrightarrow{\text{Sn/HCl}} \text{ / } \xrightarrow{\text{Br}_2} \text{ / } \xrightarrow{\text{NaNO}_2} \text{ / } \xrightarrow{\text{NaBr}}$

Answer: B

Solution:

In first step, NO_2 group is electron withdrawing group, it decreases the electron density in meta position and bromination occur in meta position. In second step, Sn/HCl used for reduction to convert $-\text{NO}_2$ to $-\text{NH}_2$ and form aniline. In next step, diazonium salt is formed and at last bromine replace N_2^+Cl^- group to form 1, 3-dibromobenzene.

Complete reaction is as follows




Question109

An amine on reaction with benzene sulphonyl chloride produces a compound insoluble in alkaline solution. This amine can be prepared by ammonolysis of ethyl chloride. The correct structure of amine is [26 Feb 2021 Shift 1]

Options:

A.



B. $\text{CH}_3\text{CH}_2\text{NH}_2$

C. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NHCH}_3$

D. $\text{CH}_3\text{CH}_2\text{CH}_2\overset{\text{H}}{\underset{|}{\text{N}}} - \text{CH}_2\text{CH}_3$

Answer: D

Solution:

The amine on reaction with benzene sulphonyl chloride (Heisenberg reagent) produces a compound insoluble in alkali. It indicates the amine is a 2° amine. i.e. all options are possible except option (b) which is a 1° amine ($\text{CH}_3\text{CH}_2\text{NH}_2$).

As this 2° amine can be prepared by ammonolysis of ethyl chloride, the 2°-amine should have at least one

ethyl (C_2H_5) group.


(a) $Ph-NH-CH_2CH_2CH_3$ does not have ethyl group.

(c) $CH_3CH_2CH_2-NH-CH_3$ does not have ethyl group.


(d) $CH_3CH_2CH_2-NH-CH_2CH_3$ has one ethyl group.

So, option (d) is the correct answer.

Preparation $CH_3CH_2CH_2-NH-CH_2CH_3$ by ammonolysis of CH_3CH_2Cl




Heisenberg test of $CH_3CH_2-NH-CH_2CH_2CH_3$ or $Et-NH-nPr$



Question 110

In the following reaction the reason why metanitro product also formed is



[24 Feb 2021 Shift 1]

Options:

A. low temperature


B. $-\text{NH}_2$ group is highly meta-directive

C. Formation of anilinium ion

D. $-\text{NO}_2$ substitution always takes place at meta-position

Answer: C


Solution:



The positive anilinium nitrogen deactivates the benzene ring due to $-\text{I}$ effect. It will behave as a m -directing group as $-\text{I}$ effect is more on $-\text{o}$ and $-\text{p}$ position and less on m position. Therefore $-\text{m}$ product is also formed along with $-\text{o}$ and $-\text{p}$ product.

Question 11


'A' and 'B' in the following reactions are :




[24 Feb 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: C

Solution:




Question112

Hofmann bromamide degradation of benzamide gives product A, which upon heating with CHCl_3 and NaOH gives product B. The structures of A


and B are
[17 Mar 2021 Shift 1]

Options:


A.




B.



C.



D.




Answer: C


Solution:

Hofmann bromamide degradation of benzamide gives product aniline (A), which upon heating with CHCl_3 and NaOH gives product benzene isocyanide (B), by carbylamine reaction. Hofmann bromamide degradation method is for preparation of primary amines by treating an amide with bromine in an aqueous or ethanolic solution of sodium hydroxide. The amine formed contains one carbon

less than that present in the amide.




Aliphatic and aromatic primary amines on heating with chloroform and ethanolic potassium hydroxide form isocyanides or carbylamines which are foul smelling substances.



Note Secondary and tertiary amines do not show this reaction. This reaction is known as carbylamine reaction or isocyanide test and is used as a test for primary amines.

Question113



Consider the given reaction, percentage yield of
[18 Mar 2021 Shift 2]

Options:

- A. (C) > (A) > (B)
- B. (B) > (C) > (A)
- C. (A) > (C) > (B)
- D. (C) > (B) > (A)


Answer: C

Solution:

During nitration of aniline, meta-nitroaniline is also formed as product due to formation of NH_3^+ group which is meta directing group under strongly acidic medium. The percentage of p, m and o product is 51%, 47% and


2%, respectively.

Reaction is as follows



% yield order is $\text{C} > \text{B} > \text{A}$.


Question 114




[18 Mar 2021 Shift 1]

Options:


A.




B.



C.




D.




Answer: B

Solution:

Aniline reacts with sodium nitrite and HCl in cold condition to form benzene diazonium chloride (X) and on further reaction, it forms azo dye (Y). Chemical reactions are as follows




Question115




In the above reaction, the structural formula of (A), X and Y respectively are
[17 Mar 2021 Shift 2]

Options:


A.




B.



C.



D.




Answer: A


Solution:

The structural formula of (A), X and Y are p-methoxy benzene diazonium chloride, acetaldehyde and hydrochloric acid respectively.

Reaction involved are as follows



Question116




In the above chemical reaction, intermediate X and reagent / condition A are


[16 Mar 2021 Shift 1]

Options:


A.




B.



C.



D.



Answer: C



Solution:

Aniline reacts with NaNO_2 , HCl at $273 - 278 \text{ }^\circ\text{K}$ to give benzene diazonium salt

$\left(\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^- = \text{X} \right)$ which on hydrolysis

$\left(\text{H}_2\text{O} \xrightarrow{\Delta} \right)$ gives phenol.

Complete reaction is as follows



Question117

In the reaction of hypobromite with amide, the carbonyl carbon is lost as [18 Mar 2021 Shift 2]

Options:


- A. CO_3^{2-}
- B. HCO_3^-
- C. CO_2
- D. CO

Answer: A


Solution:

Hypobromite transforms the primary amide to carbonic acid which further loses carbonyl carbon as CO_3^{2-} , yielding the amine product.


Chemical reaction is as follows



Mechanism



Thus, the carbonyl carbon is lost as CO_3^{2-} in the given reaction. The complete reaction in ionic form looks like.



Question118

An organic compound "A" on treatment with benzene sulphonyl chloride gives compound B . B is soluble in dil. NaOH solution. Compound A is [18 Mar 2021 Shift 2]


Options:

- A. $C_6H_5-N-(CH_3)_2$
- B. $C_6H_5-NHCH_2CH_3$
- C. $C_6H_5-CH_2NHCH_3$
- D. $C_6H_5-\underset{\underset{CH_3}{|}}{C}H-NH_2$

Answer: D

Solution:

Hinsberg reagent (benzene sulphonyl chloride) with 1° amine gives sulphonamide product i.e. N-ethylbenzene sulphonamide and it is soluble in dil. NaOH .



Hence, A is 1° amine i.e. $C_6H_5 - \underset{\substack{| \\ CH_3}}{C} H - NH_2$

Question119

Primary, secondary and tertiary amines can be separated using [17 Mar 2021 Shift 2]

Options:


- A. para-toluene sulphonyl chloride
- B. chloroform and K OH
- C. benzene sulphonic acid
- D. acetyl amide

Answer: A


Solution:

Primary, secondary and tertiary amines can be separated using para-toluene sulphonyl chloride. This test is known as Hinsberg test. Reaction involved are as follows:


Primary (1°) amine Primary amines react with para toluene sulphonyl chloride to form a precipitate that is soluble in NaOH .



Secondary (2°) amine Secondary amines reacts with para toluene sulphonyl chloride to give a precipitate that is insoluble in NaOH .






Tertiary (3°) amine Tertiary amines do not react with para toluene.



Question120

Which of the following is least basic?
[16 Mar 2021 Shift 2]



Options:



Answer: C

Solution:

Basic nature is directly proportional to electron releasing tendency. If lone pair is delocalised, electron density decreases and basic strength decreases.




So, there is delocalisation of electron in both options (a) and (c).

But in option (c) there is delocalisation of electron with both carbonyl group. Hence, it is the weakest base.

There is a subtle interplay of the inductive effect, solvation effect and steric hindrance of the alkyl group, which decides the basic strength of alkyl amines in the aqueous state.

The order of basic strength in case of ethyl substituted amines in aqueous solution is as follows



$\therefore (\text{C}_2\text{H}_5)_2\ddot{\text{N}}\text{H}$ is strong base.

Question121

Ammonolysis of alkyl halides followed by the treatment with NaOH solution can be used to prepare primary, secondary and tertiary amines.

The purpose of NaOH in the reaction is [16 Mar 2021 Shift 2]


Options:

- A. to remove basic impurities
- B. to activate NH_3 used in the reaction
- C. to remove acidic impurities
- D. to increase the reactivity of alkyl halide

Answer: C

Solution:

Ammonolysis of alkyl halides



During the reaction HX (acid) is formed. Hence, we use NaOH to remove this acidic impurities.

Question122

Which of the following is least basic? [16 Mar 2021 Shift 2]



Options:

- A. $(\text{CH}_3\text{CO})\ddot{\text{N}}\text{HC}_2\text{H}_5$
- B. $(\text{C}_2\text{H}_5)_3\ddot{\text{N}}$
- C. $(\text{CH}_3\text{CO})_2\ddot{\text{N}}\text{H}$
- D. $(\text{C}_2\text{H}_5)_2\ddot{\text{N}}\text{H}$

Answer: C

Solution:

Basic nature is directly proportional to electron releasing tendency. If lone pair is delocalised, electron density decreases and basic strength decreases.




So, there is delocalisation of electron in both options (a) and (c).

But in option (c) there is delocalisation of electron with both carbonyl groups. Hence, it is the weakest base.

There is a subtle interplay of the inductive effect, solvation effect and steric hindrance of the alkyl group, which decides the basic strength of alkyl amines in the aqueous state.

The order of basic strength in case of ethyl substituted amines in aqueous solution is as follows



$\therefore (\text{C}_2\text{H}_5)_2\ddot{\text{N}}\text{H}$ is strong base.

Question 123

Ammonolysis of alkyl halides followed by the treatment with NaOH solution can be used to prepare primary, secondary and tertiary amines. The purpose of NaOH in the reaction is [16 Mar 2021 Shift 2]


Options:

- A. to remove basic impurities
- B. to activate NH_3 used in the reaction
- C. to remove acidic impurities
- D. to increase the reactivity of alkyl halide

Answer: C

Solution:

Ammonolysis of alkyl halides



During the reaction HX (acid) is formed. Hence, we use NaOH to remove this acidic impurities.


Question 124

Which of the following reaction does not involve Hoffmann bromamide degradation?


[16 Mar 2021 Shift 1]

Options:


A.




B.



C.




D.




Answer: C

Solution:


Methyl ketone gives haloform reaction to form carboxylic acid salt which on heating with NH_3 forms amide. Amide is reduced to amine by LiAlH_4 .




(a) Benzamide undergoes Hofmann bromamide degradation when treated with bromine and aqueous sodium hydroxide, to produce benzylamine.



(b) Benzene nitrile undergoes alkaline hydrolysis to form acidamide which further gives Hofmann bromamide degradation.



(d) Acid chloride undergoes nucleophilic substitution with NH_3 to form acid amide. Acid amide undergoes Hoffmann bromamide degradation.




Question 125


Which one of the following reactions does not occur ?
[22 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.





Answer: C

Solution:



- (1) Aniline is lewis base give acid base reaction with AlCl_3 and form Anilinium ion
 (2) Anilinium ion has strongest deactivated ring so further Friedel craft Alkylation not occurs.


Question126




In the above reactions, product A and product B respectively are:
[20 Jul 2021 Shift 2]

Options:


A.




B.



C.





D.



Answer: D

Solution:




Question127


**The correct structure of Rhumann's Purple, the compound formed in the reaction of ninhydrin with proteins is :
[20 Jul 2021 Shift 1]**

Options:


A.




B.



C.




D.



Answer: D


Solution:



Ninhydrin Test

Question128


What is the major product "P" of the following reaction ?




[25 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: D

Solution:




Question129




Consider the above reaction, the Product "P" is :
[25 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question130




**In the chemical reactions given above A and B respectively are:
[22 Jul 2021 Shift 2]**

Options:


- A. H_3PO_2 and $\text{CH}_3\text{CH}_2\text{Cl}$
- B. $\text{CH}_3\text{CH}_2\text{OH}$ and H_3PO_2
- C. H_3O_2 and $\text{CH}_3\text{CH}_2\text{OH}$
- D. $\text{CH}_3\text{CH}_2\text{Cl}$ and H_3PO_2

Answer: A

Solution:




Question131




Consider the above reaction, compound B is :
[20 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question 132


What is A in the following reaction ?




[27 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question133


Which one of the products of the following reactions does not react with Hinsberg reagent to form sulphonamide?
[25 Jul 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: B

Solution:




Question134

The number of nitrogen atoms in a semi carbazone molecule of acetone is _____.

[20 Jul 2021 Shift 1]

Answer: 3

Solution:



Semicarbazone molecule of acetone

Question135

Compound A is converted to B on reaction with CHCl_3 and KOH. The compound B is toxic and can be decomposed by C. A, B and C respectively are :


[20 Jul 2021 Shift 1]

Options:

- A. primary amine, nitrile compound, conc. HCl
- B. secondary amine, isonitrile compound, conc. NaOH
- C. primary amine, isonitrile compound, conc. HCl
- D. secondary amine, nitrile compound, conc. NaOH

Answer: C

Solution:



Question136

Given below are two statements :

Statement I : Aniline is less basic than acetamide.

Statement II : In aniline, the lone pair of electrons on nitrogen atom is delocalised over benzene ring due to resonance and hence less available to a proton.

**Choose the most appropriate option ;
[27 Jul 2021 Shift 1]**

Options:

- A. Statement I is true but statement II is false.
- B. Statement I is false but statement II is true.
- C. Both statement I and statement II are true.
- D. Both statement I and statement II are false.

Answer: B

Solution:


Explanation :- aniline is more basic than acetamide because in acetamide, lone pair of nitrogen is delocalised to more electronegative element oxygen.
In Aniline lone pair of nitrogen delocalised over benzene ring.

Question137


**The major products A and B in the following set of reactions are
[31 Aug 2021 Shift 1]**

Options:


A.




B.



C.




D.



Answer: C


Solution:

Nitriles can be converted to 1° amines by reaction with LiAlH_4 .




Mechanism

(i) Hydride nucleophile attacks the electrophilic carbon in nitrile to form imine anion.




(ii) Hydrolysis of amine derivative.




2-hydroxy-2-methyl propionitrile on acidic hydrolysis gives

2-hydroxy-2-methylpropionic acid.




Mechanism



The last step of dehydration takes place in presence of H_2SO_4 to form 2-methylacrylic acid.


2-methylacrylic acid is the product (B).



Hence, correct answer is (c).

Question138


The major product of the following reaction is




[27 Aug 2021 Shift 1]

Options:


A.




B.



C.



D.




Answer: C

Solution:

In first step, ammonia reacts with ethanoyl chloride to give ethanamide and HCl. In the second step degradation of amide takes place that is known as Hoffmann's bromamide degradation reaction. The third step is diazotisation, where an unstable diazonium salt is obtained, which further gives alcohol as a major

product and alkene and alkyl halides.

The complete reaction is



Question139

Which of the following is not a correct statement for primary aliphatic amines?

[27 Aug 2021 Shift 1]

Options:

- A. The intermolecular association in primary amines is less than the intermolecular association in secondary amines.
- B. Primary amines on treating with nitrous acid solution form corresponding alcohols except methyl amine.
- C. Primary amines are less basic than the secondary amines.
- D. Primary amines can be prepared by the gabriel phthalimide synthesis.


Answer: A

Solution:

The intermolecular association is more prominent in case of primary amines as compared to secondary, due to the availability of two hydrogen atom.

Question140


Identify correct A, B and C in the reaction sequence given below.




[31 Aug 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: A

Solution:

Benzene undergoes nitration on reaction with conc. HNO_3 and conc. H_2SO_4 .




Nitrobenzene on reaction with Cl_2 / anhyd. AlCl_3 undergoes electrophilic substitution reaction to give 1-chloro-3-nitro benzene, which on reaction with Fe/HCl gives chloroaniline.



Question141


The major products A and B formed in the following reaction sequence are




[31 Aug 2021 Shift 2]

Options:


A.




B.



C.



D.




Answer: B

Solution:

Aniline reacts with acetic anhydride to give acetanilide.

The acetic acid part is removed and N-acylation takes place by nucleophilic substitution reaction.



Acetanilide is brominated with bromine/acetic acid. As $-\text{NHCOCH}_3$ is ortho/para directing group (since lone pair on N activates the ring).
Bromine attached to para position is the major product.

Question142

The total number of reagents from those given below, that can convert nitrobenzene into aniline is (Integer answer)

<i>I . Sn – HCl</i>	<i>II . Sn – NH_4OH</i>
<i>III . Fe – HCl</i>	<i>IV . Zn – HCl</i>
<i>V . H_2 – Pd</i>	<i>VI . H_2 – Raney nickel</i>

[31 Aug 2021 Shift 1]

Answer: 5

Solution:

The nitrobenzene can be reduced to aniline via following reagents.


(I) Sn-HCl (V) H_2 -Pd

(III) Fe-HCl (VI) H_2 (Raney Ni)

(IV) Zn-HCl

Question143


The correct structures of A and B formed in the following reactions are




[27 Aug 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: D

Solution:

Since, only one equivalent of acetic anhydride is present and $-NH_2$ group is more nucleophilic than $-OH$ group due to lower electronegativity of nitrogen. So, it attacks at the carbonyl center of acetic anhydride forming the product B.




Mechanism



Therefore, the option (d) is correct.


Question144




The major product in the above reaction is
[26 Aug 2021 Shift 2]

Options:


A.




B.



C.




D.



Answer: D

Solution:

Since $\text{—CH}_2\text{—NH}_2$ is more basic, the resulting amide will fail to react further. Therefore, imide formation takes place at only one site (i.e. primary amine). The complete reaction is as follows :




Question145


Which one of the following gives the most stable diazonium salt?
[1 Sep 2021 Shift 2]

Options:


A.




B.



C.




D.



Answer: B

Solution:


Primary amines react with NaNO_2 and HCl to give diazonium salts.



Diazonium salt is not formed rather N-alkylnitroso amine is formed as product.
 $\therefore$ In option (b) the most stable diazonium salt is formed.

Question146


Identify A in the following reaction.




[1 Sep 2021 Shift 2]

Options:


A.




B.



C.



D.




Answer: A

Solution:


Aniline reacts with potassium dichromat ($\text{K}_2\text{Cr}_2\text{O}_7$) to form 1,4 - benzoquinone as the product. The nascent oxygen produced from ($\text{K}_2\text{Cr}_2\text{O}_7$) reacts with aniline to form 1,4 - benzoquinone with the removal of NH_4^+ ion.

This reaction involves oxidation of aniline



Question147


The major product **Z** obtained in the following reaction scheme is:




[Jan. 09,2020(I)]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question148

Consider the following reactions,




The compound [P] is:


[Jan. 09,2020(II)]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question149

The decreasing order of basicity of the following amines is:



[Jan. 09,2020(II)]

Options:

- A. (A) > (C) > (D) > (B)
- B. (C) > (A) > (B) > (D)
- C. (B) > (C) > (D) > (A)
- D. (C) > (B) > (A) > (D)


Answer: D

Solution:

Basic strength of amines depends upon availability of lone pair of electrons. Aliphatic amines are more basic than aromatic amines.

Question150

The increasing order of pK_b for the following compounds will be:



(C)
[Jan. 07,2020 (I)]

Options:

- A. (B) < (C) < (A)
- B. (A) < (B) < (C)

C. (C) < (A) < (B)

D. (B) < (A) < (C)

Answer: D

Solution:


Conjugate acid of guanadine(B) is resonance stabilised and have 2 resonance structure. Similarly conjugate acid of (A) is also resonance stabilised and have one resonance structure. (C) does not exhibit resonance structure.

therefore the basic orderis, k_b : (B) > (A) > (C)

$\therefore pK_b$: (B) < (A) < (C)

Question151

In the following reaction sequence,




the major product B is:

[Jan. 07,2020(II)]

Options:


A.




B.



C.




D.



Answer: A

Solution:




Question152


Which of these will produce the highest yield in Friedel Crafts reaction?
[Jan.09,2020(I)]

Options:


A.




B.



C.



D.




Answer: C

Solution:

Aniline and phenol form complex with lewis acid. Chlorobenzene produces highest yield in Friedel craft reaction among the given options.

Question153

The increasing order of pK_b values of the following compounds is :



[Sep. 06, 2020(I)]

Options:

A. II < IV < III < I


B. I < II < IV < III

C. II < I < III < IV

D. I < II < II < IV

Answer: B

Solution:



-OCH₃ group increases electron density of ring at o - and p - position making (I) most basic. (III) is least basic due to -I effect of -CN group at meta position.

Since, -I effect of -CN > -I effect of -OH group.

Hence, correct basic strength will follow the order

I > II > IV > III

Basic strength $\propto \frac{1}{\text{pK}_b \text{ value}}$

Order of K_b value is, I < II < IV < III.


Question 154

Which of the following compounds can be prepared in good yield by Gabriel phthalimide synthesis?

[Sep. 06, 2020(II)]


Options:

A.




B. CH₃-CH₂-NHCH₃

C.



D.



Answer: A

Solution:

Gabriel phthalimide synthesis gives 1° amine in good yield.

Question155

The most appropriate reagent for conversion of C_2H_5CN into $CH_3CH_2CH_2NH_2$ is:

[Sep. 05, 2020 (I)]

Options:

A. $NaBH_4$


B. CaH_2

C. $LiAlH_4$

D. $Na(CN)BH_3$


Answer: C

Solution:



Question156

The increasing order of basicity of the following compounds is:




[Sep, 05,2020(I)]

Options:

- A. (A) < (B) < (C) < (D)
- B. (B) < (A) < (D) < (C)
- C. (D) < (A) < (B) < (C)
- D. (B) < (A) < (C) < (D)

Answer: B


Solution:



Thus, increasing order of basicity is (B) < (A) < (D) < (C)

Question157


The final major product of the following reaction is:




[Sep. 05, 2020(II)]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question 158


In the following reaction sequence, [C] is:




[Sep. 04, 2020(II)]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question 159


Three isomers A, B and C (mol. formula $\text{C}_8\text{H}_{11}\text{N}$) give the following results :




R has lower boiling point than S $\xrightarrow{\text{C}_6\text{H}_5\text{SO}_2\text{Cl}}$ alkali-insoluble product
 A, B and C, respectively are:
 [Sep. 03, 2020 (II)]

Options:


A.




B.



C.








D.





Answer: B

Solution:



Question160


The major product of the following reaction is:




[Jan. 12, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question161

In the following compound,



the favourable site/s for protonation is/are:
[Jan. 11,2019 (II)]

Options:

- A. (a) and (c)
- B. (b), (c) and (d)
- C. (a) and (d)
- D. (a)


Answer: B

Solution:

Higher the electronegativity ($sp^2 > sp^3$) of an atom, more will be its capacity to take proton. Thus N labelled as (b), (c) and (d) are better protonated than the N at (a) and (e).

Question162


The major product formed in the reaction given below will be:




[Jan. 10,2019(I)]

Options:


A.




B.



C.




D.




Answer: A

Solution:




Question163


Arrange the following amines in the decreasing order of basicity:



I



II



III

[Jan. 9,2019(I)]

Options:

A. I > II > III

B. III > I > II

C. III > II > I

D. I > III > II


Answer: B

Solution:

Compound, III is most basic as the lone pair of nitrogen is easily available for the donation. In case of compound (I) lone pair is not involved in resonance but nitrogen atom is sp^2 hybridised, whereas in compound II the lone pair of nitrogen is involved in aromaticity which makes it least basic.

Question164


The major product obtained in the following reaction is:




[Jan. 9,2019 (II)]

Options:


A.




B.



C.




D.



Answer: D




Solution:

-NH_2 is a stronger base than -OH , hence 1 equivalent of acetic anhydride will acetylate only -NH_2 group.
Reaction involved:



Question 165

The increasing basicity order of the following compounds is:



[Jan. 9, 2019 (II)]

Options:


- A. (D) < (C) < (B) < (A)
- B. (D) < (C) < (A) < (B)
- C. (A) < (B) < (C) < (D)
- D. (A) < (B) < (D) < (C)

Answer: B

Solution:


pK_b	
(A) $EtNH_2$	3.29
(B) $(Et)_2NH$	3.00
(C) Me_3N	4.22
(D) $Ph-NH-Me$	4.7

So, order of basic strength is:



Question166

The increasing order of reactivity of the following compound towards reaction with alkyl halides directly is:




[Jan.12,2019(I)]

Options:

- A. (B) < (A) < (C) < (D)
- B. (A) < (B) < (C) < (D)
- C. (B) < (A) < (D) < (C)
- D. (A) < (C) < (D) < (B)

Answer: A

Solution:




Question 167


The increasing order of the reactivity of the following with LiAlH_4 is :
[Jan. 12, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: C

Solution:

The reactivity order of carboxylic acid derivatives depends on the electrophilicity of the carbonyl carbon and leaving tendency of the leaving group. Leaving ability of the four groups is




Therefore, reactivity order towards LiAlH_4 is



Question168


A compound 'X' on treatment with Br_2/NaOH , provided $\text{C}_3\text{H}_9\text{N}$, which gives positive carbylamine test. Compound 'X' is:
[Jan. 11, 2019 (II)]

Options:




Answer: C

Solution:



Question169


What will be the major product in the following mononitration reaction?




[Jan. 10,2019 (II)]

Options:


A.




B.



C.



D.




Answer: D

Solution:


In the given nitration reaction, major product will be formed as per the activating group, $-\text{NH}$ part of

$-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-$ is activating, while $-\overset{\text{O}}{\parallel}{\text{C}}-$ part is deactivating group.



Question170

The compounds A and B in the following reaction are, respectively:




[Jan. 9,2019 (I)]

Options:

- A. A = Benzyl alcohol, B = Benzyl cyanide
- B. A = Benzyl chloride, B = Benzyl cyanide
- C. A = Benzyl alcohol, B = Benzy 1 isocyanide
- D. A – B enzy chloride, B - Benzyl isocyanide


Answer: D

Solution:



Question171


The major product of the following reaction is:




[Jan. 9,2019 (II)]

Options:


A.




B.



C.




D.



Answer: C


Solution:

Reaction involved for the given reaction:



Question172

The increasing order of the pK_b of the following compounds is:



[April 12, 2019 (I)]

Options:

- A. (A) < (C) < (D) < (B)
- B. (C) < (A) < (D) < (B)
- C. (B) < (D) < (A) < (C)
- D. (B) < (D) < (C) < (A)

Answer: C

Solution:

Electron withdrawing group attached to benzene ring will reduce the basic strength and increase pK_b , while electron donating group decreases pK_b .



Therefore the correct increasing order of pK_b is

(B) < (D) < (A) < (C)

Question173

Ethylamine ($\text{C}_2\text{H}_5\text{NH}_2$) can be obtained from N-ethyl phthalimide on treatment with:
[April 10, 2019(I)]


Options:



Answer: A

Solution:


N-Ethyl phthalimide on treatment with NH_2NH_2 gives ethylamine.



In place of NH_2NH_2 we can also use H_2O in presence of H^+ or OH^- as a catalyst.


Question 174

The major product of the following reaction is:





[April 10, 2019 (I)]

Options:




C.



Answer: B

Solution:



Question175

Which of the following is NOT a correct method of the preparation of benzylamine from cyanobenzene?

[April 10,2019(II)]

Options:

A. H_2/Ni


B. (i) LiAlH_4 (ii) H_3O^+

C. (i) $\text{SnCl}_2 + \text{HCl}$ (gas) (ii) NaBH_4

D. (i) $\text{HCl}/\text{H}_2\text{O}$ (ii) NaBH_4

Answer: D

Solution:



Question 176


The major product ' Y ' in the following reaction is:




[April 10, 2019 (II)]

Options:


A.




B.



C.




D.



Answer: A


Solution:



Question177

Hinsberg's reagent is:
[April 9, 2019 (II)]

Options:



Answer: C

Solution:

Hinsberg's reagent is benzenesulphonyl chloride ($\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$). It is used for detection of primary, secondary and tertiary amines.

Question178

Which of the following amines can be prepared by Gabriel phthalimide reaction?
[April 8,2019(I)]

Options:

A. n -butylamine

B. triethylamine


C. t -butylamine

D. neo-pentylamine

Answer: A

Solution:

Primary amines are prepared by Gabriel phthalimide synthesis



Question179


In the following compounds, the decreasing order of basic strength will be:
[April 8, 2019 (I)]

Options:

- A. $\text{C}_2\text{H}_5\text{NH}_2 > \text{NH}_3 > (\text{C}_2\text{H}_5)_2\text{NH}$
- B. $(\text{C}_2\text{H}_5)_2\text{NH} > \text{NH}_3 > \text{C}_2\text{H}_5\text{NH}_2$
- C. $(\text{C}_2\text{H}_5)_2\text{NH} > \text{C}_2\text{H}_5\text{NH}_2 > \text{NH}_3$
- D. $\text{NH}_3 > \text{C}_2\text{H}_5\text{NH}_2 > (\text{C}_2\text{H}_5)_2\text{NH}$

Answer: C

Solution:



Question180

Polysubstitution is a major drawback in:
[April 8,2019 (II)]

Options:

- A. Friedel Craft's alkylation
- B. Reimer Tiemann reaction
- C. Acetylation of aniline
- D. Friedel Craft's acylation


Answer: A

Solution:

Polysubstitution is a major drawback in Friedel Craft's alkylation because the introduced alkyl group is activating and gives polyalkylated product.

Question181


The major product in the following reaction is:




[April 8, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question182


The major products A and B for the following reactions are, respectively:




[April 9, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question183


The major product obtained in the following reaction is:




[April 8, 2019 (II)]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question184

Benzene diazonium chloride on reaction with aniline in the presence of dilute hydrochloric acid gives:


[April 12, 2019 (II)]

Options:


A.




B.



C.




D.



Answer: C

Solution:




Question185


Aniline dissolved in dilute HCl is reacted with sodium nitrate at 0°C. This solution was added dropwise to a solution containing equimolar mixture of aniline and phenol in dil. HCl. The structure of the major product is:
[April 9, 2019 (I)]

Options:


A.




B.



C.




D.



Answer: C

Solution:

In acidic medium, benzene diazonium chloride reacts with aniline (a basic compound) forming a₃₀ dye (c).




Question186


Coupling of benzene diazonium chloride with 1-naphthol in alkaline medium will give:
[April 8, 2019 (I)]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question 187


Products A and B formed in the following reactions are respectively:




[Online April 16, 2018]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question188

The increasing order of diazotisation of the following compounds is:



[Online April 15, 2018 (II)]

Options:

- A. (D) < (C) < (B) < (A)
- B. (A) < (D) < (B) < (C)
- C. (A) < (B) < (C) < (D)

D. (A) < (D) < (C) < (B)

Answer: D

Solution:


Aromatic diazonium salts are more stable than aliphatic diazonium salts. The higher stability of aryl diazonium salts is due to resonance. Electron donating substituents increase electron density on benzene ring. Hereby they increase the stability of diazonium salts. Electron withdrawing substituents decrease electron density on benzene ring. Hereby they decrease the stability of diazonium salts. $-\text{COCH}_3$ group is electron withdrawing and hence, diazonium salts from (D) is less stable than that from (B). Although $-\text{O}-\text{COCH}_3$ is electron donating substituent, but it is present in meta position. Hence, it will not have significant effect on stability.

Therefore the increasing order of diazotisation is

(A) < (D) < (C) < (B)

Question 189


The major product formed in the following reaction is:




[Online April 15, 2018 (II)]

Options:


A.




B.



C.




D.



Answer: A


Solution:

Note: In the given reaction a molecule of HCl is lost and $\text{C}=\text{C}$ double bond is formed. Thus it is dehydrohalogenation reaction. Nitro group is electron withdrawing group. Hence increases the acidity of H atom (attached to C atom bearing nitro group) which is removed easily. Further the newly formed double bond is in conjugation with nitro group.



Question190

The increasing order of nitration of the following compounds is:



[Online April 15, 2018 (I)]

Options:

A. $(\text{A}) < (\text{B}) < (\text{D}) < (\text{C})$

B. $(\text{A}) < (\text{B}) < (\text{C}) < (\text{D})$

C. $(\text{B}) < (\text{A}) < (\text{C}) < (\text{D})$

D. (B) < (A) < (D) < (C)

Answer: A

Solution:

In the given substituted benzene rings, the substituents methoxy ($-\text{OCH}_3$) and amino ($-\text{NH}_2$) are strongly activating groups, while methyl ($-\text{CH}_3$) is weakly activating and chloro ($-\text{Cl}$) is a deactivating group towards electrophilic aromatic substitution reaction. Since among methyl and methoxy group, methoxy group is more reactive than methyl group, (c) is more reactive than (d). Although amino group is strongly activating group, it gets protonated in presence of acid to form anilinium ion ($-\text{NH}_3^+$) which is strongly deactivating. Hence, (a) is less reactive than (c) and (d).





Chloro group is also deactivating group but less deactivating than ($-\text{NH}_3^+$). Thus order is (a) < (b) < (d) < (c)

Note:

The activating groups increase the electron density on benzene ring and thus increase the rate of electrophilic aromatic substitution reaction. The deactivating groups decrease the electron density on benzene ring and thus decrease the rate of electrophilic aromatic substitution reaction.

Question 191

Among the following compounds, the increasing order of their basic strength is:



[Online April 9, 2017]

Options:

A. (II) < (I) < (IV) < (III)

B. (II) < (I) < (III) < (IV)

C. (I) < (II) < (IV) < (III)

D. (I) < (II) < (III) < (IV)

Answer: A

Solution:


I) Resonance leads to a decrease in basic strength. In aniline, the lone pair electrons of N are in resonance with the benzene ring. Hence, the lone pairs are not easily available to donate, hence it has less basicity.
II) Pyrrole is an aromatic compound, here also lone pair of electrons are involved in the aromaticity, hence it has the least basicity.
III) In the given structure, N is bonded with 1 methyl and 2CH₂ groups. Hence, due to the +I effect, it is the most basic compound among them all.
IV) In the given structure, N is attached with only one CH group, so it has less basic character than III.
Therefore, the basic strength of given compounds is II < I < IV < III.

Question192


Which of the following compounds will form significant amount of metal product during mono-nitration reaction?
[2017]

Options:


A.




B.



C.




D.



Answer: C

Solution:

Nitration takes place in presence of conc. $\text{HNO}_3 + \text{conc. H}_2\text{SO}_4$. In strongly acidic medium, amine is converted into anilinium ion ($-\text{NH}_3^+$); substitution is thus controlled not by $-\text{NH}_2$ group but by $-\text{NH}_3^+$ group which, because of its positive charge, directs the entering group to the meta-position instead of ortho- and para.



Question193

In the Hofmann bromamide degradation reaction, the number of moles of NaOH and Br_2 used per mole of amine produced are :


[2016]

Options:

- A. Two moles of NaOH and two moles of Br_2 .
- B. Four moles of NaOH and one mole of Br_2 .
- C. One mole of NaOH and one mole of Br_2 .
- D. Four moles of NaOH and two moles of Br_2 .

Answer: B

Solution:



Question194

The test to distinguish primary, secondary and tertiary amines is:
[Online April 9, 2016]

Options:

- A. Sandmeyer's reaction
- B. Carbylamine reaction

C. Mustard oil test

D. $\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$


Answer: D

Solution:

Primary, secondary and tertiary amines are distinguished by Hinsberg reagent test. The Hinsberg reagent is an alternative name for benzene sulphonyl chloride ($\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$)

Question195

The " N " which does not contribute to the basicity for the compound is:



[Online April 10,2016]

Options:

A. N 9

B. N x3

C. N |

D. N 7

Answer: A

Solution:

Lone pair of N is involved in aromaticity.

Question196

Fluorination of an aromatic ring is easily accomplished by treating a diazonium salt with HBF_4 . Which of the following conditions is correct about this reaction?

[Online April 10, 2016]

Options:

A. NaF / Cu


B. $\text{Cu}_2\text{O} / \text{H}_2\text{O}$

C. Only heat

D. $\text{NaNO}_2 / \text{Cu}$


Answer: C

Solution:



Question 197


In the reaction




the product E is :
[2015]

Options:


A.




B.



C.





D.



Answer: A

Solution:




Question198


Arrange the following amines in the order of increasing basicity.
[Online April 10,2015]

Options:


A.




B.



C.




D.



Answer: C

Solution:

Aliphatic amines are more basic than aromatic amines thus methylamine is most basic. Electron donating groups increase the basicity whereas electron withdrawing groups decrease the basicity of the aromatic amines. Thus p-methoxyaniline is more basic than aniline which is further more basic than p-nitroaniline.



Question199


On heating an aliphatic primary amine with chloroform and ethanolic potassium hydroxide, the organic compound formed is:
[2014]

Options:

- A. an alkanol
- B. an alkanediol
- C. an alkyl cyanide
- D. an alkyl isocyanide

Answer: D

Solution:



Question200

Considering the basic strength of amines in aqueous solution, which one has the smallest pK_b value?

[2014]

Options:


- A. $(\text{CH}_3)_2\text{NH}$
- B. CH_3NH_2
- C. $(\text{CH}_3)_3\text{N}$
- D. $\text{C}_6\text{H}_5\text{NH}_2$

Answer: A

Solution:


Arylamines are less basic than alkyl amines and even ammonia. This is due to resonance. In aryl amines the lone pair of electrons on N is partly shared with the ring and is thus less available for sharing with a proton. In alkylamines, the electron releasing alkyl group increases the electron density on nitrogen atom and thus also increases the ability of amine for protonation. Hence more the no, of alkyl groups higher should be the basicity of amine. But a slight discrepancy occurs in case of trimethyl amines due to steric effect. Hence the

correct order is



Question201


The major product of the reaction




[Online April 9, 2014]

Options:


A.




B.



C.



D.



Answer: B

Question202

The final product formed when methyl amine is treated with NaNO_2 and HCl is:


[Online April 19,2014]

Options:

- A. Diazomethane
- B. Methylalcohol
- C. Methylcyanide
- D. Nitromethane

Answer: B

Solution:



Question203

Complete reduction of benzene-diazonium chloride with Zn/HCl gives:
[Online April 11, 2014]

Options:

- A. Aniline
- B. Phenylhydrazine
- C. Azobenzene
- D. Hydrazobenzene

Answer: A

Solution:

On reduction with Zn and HCl , $\text{C}_6\text{H}_5\text{N}_2\text{Cl}$ forms aniline as the main product.

Question204

Conversion of benzene diazonium chloride to chlorobenzene is an example of which of the following reactions?


[Online April 12, 2014]

Options:

- A. Claisen
- B. Friedel-craft
- C. Sandmeyer
- D. Wurtz


Answer: C

Solution:



Question205


In a set of reactions p -nitrotoluene yielded a product E.




[Online April 19, 2014]

Options:


A.




B.



C.




D.



Answer: B

Solution:



Question206

A compound with molecular mass 180 is acylated with CH_3COCl to get a compound with molecular mass 390. The number of amino groups present


**per molecule of the former compound is:
[2013]**

Options:

- A. 2
- B. 5
- C. 4
- D. 6

Answer: B

Solution:



Now since the molecular mass increases by 42 unit as a result of the reaction of one mole of CH_3COCl with one-NH₂ group and the given increase in mass is (390 – 180) 210. Hence the number of –NH₂ groups = 210/42 = 5.

Question207

**The order of basicity of amines in gaseous state is:
[Online April 23, 2013]**

Options:

- A. $1^\circ > 2^\circ > 3^\circ > \text{NH}_3$
- B. $3^\circ > 2^\circ > \text{NH}_3 > 1^\circ$
- C. $3^\circ > 2^\circ > 1^\circ > \text{NH}_3$
- D. $\text{NH}_3 > 1^\circ > 2^\circ > 3^\circ$

Answer: C

Solution:

The correct order of relative basicity of amines in the gasphase is $3^\circ > 2^\circ > 1^\circ > \text{NH}_3$. The alkyl group releases electron and thus, tends to disperse the positive charge of the alkyl ammonium ion and therefore stabilises it.

Since, NH_4^+ (from NH_3) has no such alkyl group, it is not stabilised to such an extent as alkylammonium ion.

Question208

Carbylamine forms from aliphatic or aromatic primary amine via which of the following intermediates?



[Online April 25,2013]

Options:

- A. Carbanion
- B. Carbene
- C. Carbocation
- D. Carbon radical


Answer: B

Solution:



Question209

The order of basicity of the compounds



[Online May 19,2012 ; Online May 26,2012]

Options:

- A. IV > I > III > II
- B. I > III > II > IV
- C. III > II > IV > I
- D. II > I > III > IV

Answer: B

Solution:

The order of basicity is I > III > II > IV. The lone pair of electrons on N is more readily available for protonation in I and III than in II. III contains an oxygen atom which has –I effect due to which it is less basic than I. In compound IV lone pair of electrons on N-atom is contributed towards the aromatic sextet formation and hence it is not at all available for protonation.

Question210

The conversion of benzene diazonium chloride to bromobenzene can be accomplished by

[Online May 12,2012]


Options:

- A. Reimer-Tiemann reaction
- B. Friedel-Crafts reaction
- C. Gattermann reaction
- D. Azo-Coupling reaction

Answer: C


Solution:

Diazonium salts react with copper powder and hydrogen bromide to form bromobenzene. This reaction is known as Gattermann reaction.



Question211

In the chemical reactions:



the compounds **A** and **B** respectively are:
[2011 RS]

Options:

- A. Benzene diazonium chloride and benzonitrile
- B. Nitrobenzene and chlorobenzene
- C. Phenol and bromobenzene
- D. Fluorobenzene and phenol


Answer: A

Solution:



Question212

In the chemical reactions,




the compounds 'A' and 'B' respectively are
[2010]

Options:


- A. nitrobenzene and fluorobenzene
- B. phenol and benzene
- C. benzene diazonium chloride and fluorobenzene
- D. nitrobenzene and chlorobenzene

Answer: C

Solution:



The diazonium group can be replaced by fluorine by treating the diazonium salt with fluoroboric acid (HBF₄). The precipitated diazonium fluoroborate is isolated, dried and heated until decomposition occurs to yield the aryl fluoride. This reaction is known as Balz-Schiemann reaction.



Question213

Which one of the following is the strongest base in aqueous solution?
[2007]

Options:

- A. Methylamine
- B. Trimethylamine
- C. Aniline
- D. Dimethylamine.

Answer: D

Solution:


Note: Aromatic amines (e.g. aniline) are less basic than aliphatic amines due to delocalisation of electrons. Among aliphatic amines the order of basicity in aq. solution is $2^\circ > 1^\circ > 3^\circ$. The electron density is decreased in 3° amine due to crowding of alkyl group over N atom which makes the approach and bonding by a proton relatively difficult. Therefore the basicity decreases.

$\therefore$ The correct order of basic strength is

Dimethylamine > Methyl amine > Trimethylamine > Aniline.

Question214

In the chemical reaction,



the compounds (A) and (B) are respectively
[2007]

Options:

A. $\text{C}_2\text{H}_5\text{NC}$ and 3KCl

B. $\text{C}_2\text{H}_5\text{CN}$ and 3KCl

C. $\text{CH}_3\text{CH}_2\text{CONH}_2$ and 3KCl

D. $\text{C}_2\text{H}_5\text{NC}$ and K_2CO_3 .

Answer: A

Solution:

This is carbylamine reaction.



Question215


The most basic compound among the following is
[2005, Online May 7,2012]

Options:

- A. Acetanilide
- B. Benzylamine
- C. p -Nitro aniline
- D. Aniline

Answer: B

Solution:



compound is most basic due to presence of localized lone pair of electron on nitrogen atom while other compounds have delocalized lone pair of electron.

Question216

Which one of the following methods is neither meant for the synthesis nor for separation of amines?
[2005]

Options:

- A. Curtius reaction
- B. Wurtz reaction
- C. Hofmann method
- D. Hinsberg method

Answer: B

Solution:

Wurtz reaction is used for the preparation of hydrocarbons from alkyl halides.
$$RX + 2Na + X'R \rightarrow R-R' + 2NaX$$

Question217

The correct order of increasing basic nature for the bases NH_3 , CH_3NH_2 and $(\text{CH}_3)_2\text{NH}$ is [2003]

Options:



Answer: B

Solution:


The alkyl groups are electron releasing (+ I) groups, thus increase the electron density around the nitrogen thereby increasing the availability of the lone pair of electrons to proton or Lewis acid and making the amine more basic. Hence more the number of alkyl groups, more basic is the amine. Therefore, the correct order is $\text{NH}_3 < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$

Question218


The reaction of chloroform with alcoholic KOH and p-toluidine forms [2003]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question219

Ethyl isocyanide on hydrolysis in acidic medium generates [2003]


Options:

- A. propanoic acid and ammonium salt
- B. ethanoic acid and ammonium salt
- C. methylamine salt and ethanoic acid
- D. ethylamine salt and methanoic acid

Answer: D

Solution:

Ethyl isocyanide on hydrolysis form primary amines.



Question220

When a primary amine reacts with chloroform in ethanolic K OH then the product is
[2002]

Options:

- A. an isocyanide
- B. an aldehyde
- C. a cyanide
- D. an alcohol.

Answer: A

Solution:

