

Alcohols, Phenols and Ethers

Question1


One mole of phenol is treated with dilute HNO_3 at 298 K to give a mixture of products. The mixture is separated by steam distillation. The steam volatile compound (X) is separated. The increase in percentage of oxygen in (X) with respect to phenol is $\text{_____} \times 10^{-1}\%$

(Given molar mass in gmol^{-1} H : 1, C : 12, N : 14, O : 16)

JEE Main 2026 (Online) 5th April Morning Shift

Answer: 175

Solution:



Step 1: Calculate the percentage of oxygen in phenol.

The molecular formula of phenol is $\text{C}_6\text{H}_6\text{O}$. Its molecular mass is:

$$(6 \times 12) + (6 \times 1) + 16 = 94 \text{ g/mol}$$

Percentage of oxygen in phenol:

$$\frac{16}{94} \times 100 = 17.02\%$$

Step 2: Calculate the percentage of oxygen in o-nitrophenol.

When phenol reacts with dilute HNO_3 , the steam-volatile product formed is o-nitrophenol ($\text{C}_6\text{H}_5\text{NO}_3$).

Molecular mass of o-nitrophenol:

$$(6 \times 12) + (5 \times 1) + 14 + (3 \times 16) = 139 \text{ g/mol}$$

Percentage of oxygen in o-nitrophenol:

$$\frac{48}{139} \times 100 = 34.53\%$$

Step 3: Find the percentage increase in oxygen.

Increase in oxygen percentage:

$$34.53 - 17.02 = 17.5$$

Step 4: Express the answer in the required form.

The increase in percentage of oxygen with respect to phenol is:

$$17.5 \times 10^{-1} = 175$$

Question2

4.7 g of phenol is heated with Zn to give product X . If this reaction goes to 60% completion then the number of moles of compound X formed will be


_____ $\times 10^{-2}$. (Nearest Integer)

(Given molar mass in gmol^{-1} : H : 1, C : 12, O : 16)

JEE Main 2026 (Online) 6th April Morning Shift

Answer: 3

Solution:



Given mass of phenol = 4.7 g

Molar mass of phenol ($\text{C}_6\text{H}_6\text{O}$) = $(6 \times 12) + (6 \times 1) + 16 = 94 \text{ g/mol}$

So,

$$\text{Moles of phenol} = \frac{4.7}{94.0} = 0.05 \text{ mol}$$

From the reaction, 1 mole of phenol produces 1 mole of benzene (product X) if the reaction is complete (100% yield).

Therefore, 0.05 mol of phenol will give 0.05 mol of benzene at 100% completion.

The actual yield is 60% of the theoretical yield, so:

$$\text{Moles of X formed} = 0.05 \times 0.60 = 0.03 \text{ mol}$$

$$\text{Moles of X} = 3 \times 10^{-2} \text{ mol}$$

Final Answer: (3)

Question3

Given below are two statements :

Statement I : Phenol on treatment with CHCl_3 /aq. KOH under refluxing condition, followed by acidification produces *p*-hydroxy benzaldehyde as the major product and *o*-hydroxy benzaldehyde as the minor product.

Statement II : The mixture of *p*-hydroxybenzaldehyde and *o* hydroxybenzaldehyde can be easily separated through steam distillation.

In the light of the above statements, choose the correct answer from the options given below

JEE Main 2026 (Online) 22nd January Morning Shift

Options:

A.

Statement I is true but Statement II is false

B.

Both Statement I and Statement II are true

C.


Both Statement I and Statement II are false

D.

Statement I is false but Statement II is true

Answer: D


Solution:



(can be separated by steam distillation)

Question4

3, 3-Dimethyl-2-butanol cannot be prepared by :



Choose the correct answer from the options given below :

JEE Main 2026 (Online) 22nd January Evening Shift

Options:

A.

B and E Only

B.

B, C and E Only

C.


B and C Only

D.

B Only

Answer: A

Solution:




Question5

A mixed ether (P), when heated with excess of hot concentrated hydrogen iodide produces two different alkyl iodides which when treated with aq. NaOH give compounds (Q) and (R). Both (Q) and (R) give yellow precipitate with NaOI . Identify the mixed ether (P) :


JEE Main 2026 (Online) 23rd January Evening Shift

Options:


A.




B.



C.




D.



Answer: B

Solution:



Question6

A hydroxy compound (X) with molar mass 122 g mol^{-1} is acetylated with acetic anhydride, using a large excess of the reagent ensuring complete acetylation of all hydroxyl groups. The product obtained has a molar mass of 290 g mol^{-1} . The number of hydroxyl groups present in compound (X) is:

JEE Main 2026 (Online) 24th January Morning Shift

Options:

A.

2

B.

4

C.


3

D.

5

Answer: B


Solution:



$$\text{No. of OH groups} = \frac{290-122}{42} = 4$$

Question7

From the following, how many compounds contain at least one secondary alcohol?



Choose the correct answer from the options given below :

JEE Main 2026 (Online) 24th January Evening Shift

Options:

A.

Three

B.

Two

C.

Four

D.

Five


Answer: A

Solution:

II, IV & V are secondary alcohol.

Question8

Consider the following reaction sequence



Compound (y) develops characteristic colour with neutral FeCl_3 solution.

Identify the INCORRECT statement from the following for the above sequence.

JEE Main 2026 (Online) 28th January Morning Shift

Options:

A.

Compound y will dissolve in NaHCO_3 and evolve a gas.

B.

Compound x is more acidic than compound y .

C.


Both compounds x and y will dissolve in NaOH .

D.

Both compounds x and y will burn with sooty flame.





Answer: B

Solution:



Question9

The reactions which produce alcohol as the product are :



Choose the correct answer from the options given below :

JEE Main 2026 (Online) 28th January Evening Shift

Options:

A.

B, D and E Only

B.

A and D Only

C.

A, C and E Only




D.

C and D Only

Answer: D

Solution:


Reaction given Alcohol



And. $\rightarrow$ C, D

Question10

Identify the colour of compound ' X ' in the sequence of the reaction.



JEE Main 2026 (Online) 4th April Morning Shift

Options:

A.

Violet

B.

Green

C.

Red

D.

Colourless

Answer: D

Solution:

Compound X is phenolphthalein (an acid–base indicator).



In basic solution, phenolphthalein first shows a pink colour.

But when NaOH is in excess (very strong base), phenolphthalein changes to a form that is colourless.

So, the colour of compound X in excess NaOH is **colourless**.

Question11

Consider compounds A, B and C with following structural formulae



For the conversion of B from A , reagent (D) required is _____ and structural formula of product (E) obtained when C undergoes same reaction using excess reagent (D) is _____ .

JEE Main 2026 (Online) 5th April Morning Shift

Options:

A.

D	E
Conc. H_2SO_4	$\text{CH}_2 = \text{CH} - \text{CH}(\text{OH})\text{CH}_3$

B.

D	E
PCC	$\text{HO} - \text{CH}_2 - \text{CH}_2 - \text{CH} = \text{CH}_2$

C.


D	E
PCC	$\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$

D.

D	E
Conc. H_2SO_4 or H_3PO_4	$\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$


Answer: D

Solution:



Question12

Consider the following sequence of reactions




The major product P is :


JEE Main 2026 (Online) 6th April Morning Shift

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question13

Consider the following reactions. Total number of electrons in the π bonds and lone pair of electrons in the product (X) is :



JEE Main 2026 (Online) 6th April Evening Shift

Options:

A.

12

B.

16

C.


14

D.

18

Answer: D


Solution:



Total number of π and lone pair electrons in final product = 18

Question14

Consider the following reaction.



Statement I : In the above reaction, product formed will be a mixture of benzyl alcohol and iodobenzene.

Statement II : In the above reaction, the $-\text{O}-\text{CH}_2-$ bond is cleaved to give the product.

In the light of the above statements, choose the correct answer from the options given below :

JEE Main 2026 (Online) 6th April Evening Shift

Options:

A.

Both Statement I and Statement II are true

B.

Both Statement I and Statement II are false

C.


Statement I is true but Statement II is false

D.

Statement I is false but Statement II is true

Answer: D

Solution:



Question15

What amount of bromine will be required to convert 2 g of phenol into 2,4,6-tribromophenol?

(Given molar mass in gmol^{-1} of C, H, O, Br are 12, 1, 16, 80 respectively)

JEE Main 2025 (Online) 23rd January Morning Shift

Options:

A. 6.0 g


B. 10.22 g

C. 20.44 g

D. 4.0 g

Answer: B

Solution:



$$\text{Moles of phenol} = \frac{2}{94} = 0.021$$

$$\therefore \text{Moles of bromine} = 0.021 \times 3 = 0.064$$

$$\therefore \text{Mass of bromine} = 0.064 \times 160 = 10.22 \text{ g}$$

Question16

Given below are two statements :

Statement (I) : The boiling points of alcohols and phenols increase with increase in the number of C -atoms.

Statement (II) : The boiling points of alcohols and phenols are higher in comparison to other class of compounds such as ethers, haloalkanes.

In the light of the above statements, choose the correct answer from the options given below :

JEE Main 2025 (Online) 23rd January Evening Shift

Options:

A. Statement I is false but Statement II is true

B. Both Statement I and Statement II are true

C. Statement I is true but Statement II is false

D. Both Statement I and Statement II are false

Answer: B

Solution:

Statement (I):

As the number of carbon atoms in alcohols or phenols increases, the molecule becomes larger, which in turn enhances the van der Waals (dispersion) forces between the molecules.

Stronger intermolecular forces require more energy (i.e., higher temperature) to overcome, leading to higher boiling points.

Therefore, Statement (I) is true.

Statement (II):

Alcohols and phenols have a hydroxyl group ($-OH$) that can form hydrogen bonds, which are significantly stronger than the dipole-dipole interactions typically seen in ethers or the interactions in haloalkanes.

This stronger hydrogen bonding means that alcohols and phenols have higher boiling points compared to ethers and haloalkanes of similar molecular weight.

Hence, Statement (II) is also true.


Based on the above points, the correct answer is:

Option B: Both Statement I and Statement II are true.


Question 17

Given below are two statements :

Statement I :



Statement II : In



, intramolecular substitution takes place first by involving lone pair of electrons on nitrogen.

In the light of the above statements, choose the most appropriate answer from the options given below


JEE Main 2025 (Online) 28th January Morning Shift

Options:

- A. Both Statement I and Statement II are correct
- B. Statement I is incorrect but Statement II is correct
- C. Both Statement I and Statement II are incorrect
- D. Statement I is correct but Statement II is incorrect

Answer: A

Solution:



Rate of (a) is faster than (b) because of its intramolecular substitution.


Question18

Which one of the following, with HBr will give a phenol?


JEE Main 2025 (Online) 29th January Evening Shift

Options:


A.




B.



C.




D.



Answer: A

Solution:

Relation with HBr to give phenol




(1)

The reaction involves two steps,


★) The lone pair of oxygen atom attacks on the hydrogen of HBr.

★) The nucleophile Br⁻ attacks on the methyl group which lead to the formation of phenol along with methyl bromide.


Phenol is formed from this compound on reaction with HBr.



No phenol formation in this case because alkyl chain is attached to the benzene ring and on addition of H-Br, either H⁺ or Br⁻ get attached to this chain and no -OH group is formed as it is directly bonded to the benzene ring.



No phenol formation in this case. -OH group is formed as



So, phenol product is formed in case of option (1) compound only.


Question19

Benzene is treated with oleum to produce compound (X) which when further heated with molten sodium hydroxide followed by acidification produces compound (Y). The compound Y is treated with zinc metal to produce compound (Z). Identify the structure of compound (Z) from the following option.


JEE Main 2025 (Online) 4th April Morning Shift

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question20

Given below are two statements :

Statement I : Dimethyl ether is completely soluble in water. However, diethyl ether is soluble in water to a very small extent.

Statement II : Sodium metal can be used to dry diethyl ether and not ethyl alcohol.

In the light of given statements. choose the correct answer from the options given below

JEE Main 2025 (Online) 7th April Morning Shift

Options:

- A. Statement I is false but Statement II are true
- B. Statement I is true but Statement II is false
- C. Both Statement I and Statement II are true
- D. Both Statement I and Statement II are False

Answer: C

Solution:

St-I - St-I is correct because both given ether are soluble in water $\rightarrow$ Di ethyl ether and butan-1-ol are miscible to almost same extent i.e., 7.5 and 9 gm per 100 ml water due to H -bonding

St-II : - St. II is also correct because sodium metal is not used with ethyl alcohol as H₂ gas release with ethyl a below

Question21

Given below are two statements :

Statement (I) : p-nitrophenol is more acidic than m-nitrophenol and o-nitrophenol.

Statement (II) : Ethanol will give immediate turbidity with Lucas reagent.

In the light of the above statements, choose the correct answer from the options given below :

[27-Jan-2024 Shift 1]


Options:

- A. Statement I is true but Statement II is false
- B. Both Statement I and Statement II are true
- C. Both Statement I and Statement II are false
- D. Statement I is false but Statement II is true

Answer: A

Solution:

Acidic strength



Ethanol give lucas test after long time

Statement (I) $\rightarrow$ correct





Statement (II) $\rightarrow$ incorrect

Question22

Match List-I with List-II.

List – I

(Reaction)



List – II

(Reagent(s))

(I) $\text{Na}_2\text{Cr}_2\text{O}_7, \text{H}_2\text{SO}_4$

(II) (i) NaOH (ii) CH_3Cl

(III) (i) $\text{NaOH}, \text{CHCl}_3$
(ii) NaOH (iii) HCl

(IV) (i) NaOH (ii) CO_2
(iii) HCl

Choose the correct answer from the options given below:
[27-Jan-2024 Shift 2]

Options:

A. (A)-(IV), (B)-(I), (C)-(III), (D)-(II)

B. (A)-(II), (B)-(III), (C)-(I), (D)-(IV)

C. (A)-(II), (B)-(I), (C)-(III), (D)-(IV)

D. (A)-(IV), (B)-(III), (C)-(I), (D)-(II)

Answer: D


Solution:

(A) $\rightarrow$ Kolbe Schmidt Reaction

(B) $\rightarrow$ Reimer Tiemann Reaction

(C) $\rightarrow$ Oxidation of phenol to p-benzoquinone





(D) $\rightarrow \text{PhOH} + \text{NaOH} \rightarrow \text{H}_2\text{O} + \text{PhO}^-$



Question23

Match List-I with List-II.

List – I (Reaction)



List – II (Reagent(s))

(I) $\text{Na}_2\text{Cr}_2\text{O}_7, \text{H}_2\text{SO}_4$

(II) (i) NaOH (ii) CH_3Cl

(III) (i) $\text{NaOH}, \text{CHCl}_3$
(ii) NaOH (iii) HCl

(IV) (i) NaOH (ii) CO_2
(iii) HCl

Choose the correct answer from the options given below:
[27-Jan-2024 Shift 2]

Options:

A. (A)-(IV), (B)-(I), (C)-(III), (D)-(II)

B. (A)-(II), (B)-(III), (C)-(I), (D)-(IV)




C. (A)-(II), (B)-(I), (C)-(III), (D)-(IV)

D. (A)-(IV), (B)-(III), (C)-(I), (D)-(II)

Answer: D

Question24

Major product formed in the following reaction is a mixture of:



[27-Jan-2024 Shift 2]

Options:

A. (1)


B. (2)

C. (3)

D. (4)

Answer: D

Solution:



Question25

**Phenolic group can be identified by a positive:
[27-Jan-2024 Shift 2]**

Options:

A. Phthalein dye test

B. Lucas test

C. Tollen's test

D. Carbylamine test

Answer: A

Solution:

Carbylamine Test-Identification of primary amines

Lucas Test - Differentiation between 1°, 2° and 3° alcohols

Tollen's Test - Identification of Aldehydes

Phthalein Dye Test - Identification of phenols

Question26

**Phenol treated with chloroform in presence of sodium hydroxide, which further hydrolysed in presence of an acid results
[29-Jan-2024 Shift 2]**

Options:

A. Salicylic acid


B. Benzene-1,2-diol

C. Benzene-1, 3-diol

D. 2-Hydroxybenzaldehyde

Answer: D

Solution:




It is Reimer Tiemann Reaction

Question27

Salicylaldehyde is synthesized from phenol, when reacted with
[30-Jan-2024 Shift 2]

Options:

A.




B. CO_2 , NaOH

C. CCl_4 , NaOH

D. HCCl_3 , NaOH

Answer: D

Solution:



Question28

Given below are two statements: One is labelled as Assertion A and the other is labelled as Reason R:

Assertion A: Alcohols react both as nucleophiles and electrophiles.

Reason R: Alcohols react with active metals such as sodium, potassium and aluminum to yield corresponding alkoxides and liberate hydrogen.

In the light of the above statements, choose the correct answer from the options given below:

[31-Jan-2024 Shift 1]

Options:

A. A is false but R is true.

B. A is true but R is false.

C. Both A and R are true and R is the correct explanation of A.

D. Both A and R are true but R is NOT the correct explanation of A

Answer: D

Solution:

As per NCERT, Assertion (A) and Reason (R) is correct but Reason (R) is not the correct explanation.

Question29

Given below are two statements: One is labelled as Assertion A and the other is labelled as Reason R:

Assertion A: pK_a value of phenol is 10.0 while that of ethanol is 15.9.

Reason R: Ethanol is stronger acid than phenol.

In the light of the above statements, choose the correct answer from the options given below:

[31-Jan-2024 Shift 1]

Options:

A. A is true but R is false.

B. A is false but R is true.

C. Both A and R are true and R is the correct explanation of A.

D. Both A and R are true but R is NOT the correct explanation of A.

Answer: A

Solution:

Phenol is more acidic than ethanol because conjugate base of phenoxide is more stable than ethoxide.

Question30

Match List - I with List - II.

List-I (Reactants)	List-II Products
(A) Phenol, Zn/Δ	(I) Salicylaldehyde
(B) Phenol, $CHCl_3$, NaOH, HCl	(II) Salicylic acid
(C) Phenol, CO_2 , NaOH, HCl	(III) Benzene
(D) Phenol, Conc. HNO_3	(IV) Picric acid


Choose the correct answer from the options given below.
[1-Feb-2024 Shift 2]

Options:

- A. (A)-(IV), (B), (II), (C)-(I), (D)-(III)
- B. (A)-(IV), (B)-(I), (C)-(II), (D)-(III)
- C. (A)-(III), (B)-(I), (C)-(II), (D)-(IV)
- D. (A)-(III), (B)-(IV), (C)-(I), (D)-(II)

Answer: C

Solution:



Question31

Which among the following has highest boiling point?
[1-Feb-2024 Shift 2]

Options:

- A. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$
- B. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2 - \text{OH}$
- C. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$
- D. $\text{H}_5\text{C}_2 - \text{O} - \text{C}_2\text{H}_5$


Answer: B

Solution:

Due to H-bonding boiling point of alcohol is High.

Question32


Number of compounds which give reaction with Hinsberg's reagent is_____



[1-Feb-2024 Shift 2]


Answer: 5

Solution:



Question33


'A' and 'B' formed in the following set of reactions are :




[24-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question34


In the cumene to phenol preparation in presence of air, the intermediate is [25-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question35


Find out the major product from the following reaction.




[25-Jan-2023 Shift 2]

Options:


A.




B.



C.




D.



Answer: A

Solution:



Question36

Given below are two statements, one is labelled as Assertion A and the other is labelled as Reason R

Assertion A :- Butylated hydroxyl anisole when added to butter increases its shelf life.

Reason R :- Butylated hydroxyl anisole is more reactive towards oxygen than food.

In the light of the above statements, choose the most appropriate answer from the options given below :-

[25-Jan-2023 Shift 2]

Options:

- A. Both A and R are correct and R is the correct explanation of A.
- B. A is correct but R is not correct.
- C. A is not correct but R is correct.
- D. Both A and R are correct but R is NOT the correct explanation of A.

Answer: A

Solution:

Butylated hydroxyl anisole is an antioxidant.

Question37

The increasing order of pK_a for the following phenols is

[29-Jan-2023 Shift 1]


Options:

- A. 2, 4-Dinitrophenol
- B. 4 - Nitrophenol
- C. 2, 4, 5-Trimethylphenol
- D. Phenol
- E. 3-Chlorophenol

Answer: C

Solution:

Order of acidity for following phenol is




- M and -I increases acidity
+M and +I decreases acidity

Question38


An organic compound 'A' with empirical formula C_6H_6O gives sooty flame on burning. Its reaction with bromine solution in low polarity solvent results in high yield of B. B is [31-Jan-2023 Shift 1]

Options:


A.




B.



C.




D.



Answer: A


Solution:

Aromatic compounds burns with sooty flame



Question39

Decreasing order of dehydration of the following alcohols is



[1-Feb-2023 Shift 1]

Options:

- A. $a > d > b > c$
- B. $b > d > c > a$
- C. $b > a > d > c$
- D. $d > b > c > a$


Answer: B

Solution:

Dehydration of alcohol is directly proportional to the stability of carbocation.

Question40


In the following reaction, ' B ' is




[6-Apr-2023 shift 2]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question41


Find out the major product from the following reaction




[6-Apr-2023 shift 2]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question42


The major product formed in the following reaction is:




[8-Apr-2023 shift 1]

Options:


A.




B.



C.




D.



Answer: D

Solution:



Note: Lithium borohydride is commonly used for selective reduction of esters and lactones to the corresponding alcohol.

Question43

A compound ' X ' when treated with phthalic anhydride in presence of concentrated H_2SO_4 yields ' Y ' ' Y ' is used as an acid/base indicator. ' X ' and ' Y ' are respectively:
[8-Apr-2023 shift 2]

Options:

A. Anisole, methyl orange


B. Toluidine, Phenolphthalein

C. Carboic acid, Phenolphthalein

D. Salicylaldehyde, Phenolphthalein

Answer: C

Solution:



Question44

Suitable reaction condition for preparation of Methyl phenyl ether is [10-Apr-2023 shift 1]


Options:

- A. Benzene, MeBr
- B. $\text{PhO}^{\ominus}\text{Na}^{\oplus}$, MeOH
- C. $\text{Ph}-\text{Br}$, $\text{MeO}^{\ominus}\text{Na}^{\oplus}$
- D. $\text{PhO}^{\ominus}\text{Na}^{\oplus}$, MeBr

Answer: D

Solution:

Williamson's synthesis :-



Question45

Incorrect method of preparation for alcohols from the following is: [10-Apr-2023 shift 2]


Options:

- A. Ozonolysis of alkene.
- B. Hydroboration-oxidation of alkene.
- C. Reaction of alkyl halide with aqueous NaOH.
- D. Reaction of Ketone with RMgBr followed by hydrolysis.


Answer: A

Solution:


1) Ozonolysis of alkene–



2) Hydroboration – oxidation of alkene




3) $R-X + NaOH \longrightarrow R-OH + NaX$



Question46

L-isomer of tetrose X(C₄H₈O₄) gives positive schiff's test and has two chiral carbons. On acetylation, 'X' yields triacetate. 'X' undergoes following reactions




'X' is


[11-Apr-2023 shift 1]

Options:


A.




B.



C.




D.



Answer: D


Solution:



(x) gives positive schiff's test due –CHO group

(x) is L-tetrose.




Question47




Product [X] formed in the above reaction is:

[11-Apr-2023 shift 2]


Options:





Answer: A

Solution:



Question48

Number of compounds from the following which will not produce orange red precipitate with Benedict solution is _____

Glucose, maltose, sucrose, ribose, 2-deoxyribose, amylose, lactose
[11-Apr-2023 shift 2]

Answer: 2

Solution:

Benedict test :

Glucose - ✓

maltose - ✓

sucrose - ✗

ribose - ✓


2-deoxyribose - ✓

amylose - ✗

lactose - ✓

Question49


In the following reaction




[12-Apr-2023 shift 1]

Options:


A.



B.



C.




D.




Answer: A

Solution:



Question50


In the reaction given below




[13-Apr-2023 shift 1]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question51



In the above reaction, left hand side and right hand side rings are named as ' A ' and ' B ' respectively. They undergo ring expansion. The correct statement for this process is:
[13-Apr-2023 shift 1]


Options:

A. Finally both rings will become six membered each.

- B. Ring expansion can go upto seven membered rings
- C. Finally both rings will become five membered each.
- D. Only A will become 6 membered.


Answer: A

Solution:



Question52


For the given reaction




The total number of possible products formed by tertiary carbocation of A is _____.
[13-Apr-2023 shift 1]

Answer: 5

Solution:




Question53




'A' formed in the above reaction is :
[15-Apr-2023 shift 1]

Options:


A.




B.



C.




D.



Answer: A

Solution:



Question54

Choose the correct answer from the options given below:

List-I		List-II	
(A)	A.	I.	Br_2 in CS_2
(B)	B.	II.	$\text{Na}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4$
(C)	C.	III.	Zn
(D)	D.	IV.	$\text{CHCl}_3/\text{NaOH}$

[27-Jun-2022-Shift-2]

Options:

A. A-IV, B-III, C-II, D-I


B. A-IV, B-III, C-I, D-II

C. A-II, B-III, C-I, D-IV

D. A-IV, B-II, C-III, D-I

Answer: A


Solution:



∴ Correct match is (A) - IV, (B) - III, (C) - II, (D) - I

Question55


The major product (P) of the given reaction is (where, Me is $-\text{CH}_3$)




[28-Jun-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: C

Solution:




(1) In compound A, positive charge on O atom is not stable that is why OH_2^+ gets removed.

(2) In Carbocation (B), two α -H present and after 1, 2-methyl shift number of α -H becomes six so it becomes more stable as far carbocation when number of α -H increases stability increases.

(3) Among product D, E and F, D is major product as in case of alkene most stable, alkenes are those which have more α -H. And every reactant produces that product which is more stable.

Question 56


The major product of the following reaction contains _____ bromine atom(s).



[28-Jun-2022-Shift-2]

Answer: 1


Solution:



Here only carbon 12th and 13th is not taking part in the resonance. All other carbons are taking part in resonance and lone pair of O is also taking part in the resonance. As because of resonance stability of compound increases Br will not attack to any carbon which is taking part in the resonance and decrease the stability of compound. So, Br can only attack 12 th carbon or 13 th carbon atom but 12th carbon is a alpha carbon and it can easily donate H^+ ion and add Br^- ion and increase stability.

Question57


In the given conversion the compound A is :




[29-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: B

Solution:



Question58

Given below are two statements:

Statement I : Phenols are weakly acidic.

Statement II : Therefore they are freely soluble in NaOH solution and are weaker acids than alcohols and water.

Choose the most appropriate option :

[29-Jun-2022-Shift-1]

Options:

- A. Both Statement I and Statement II are correct.
- B. Both Statement I and Statement II are incorrect.
- C. Statement I is correct but Statement II is incorrect.
- D. Statement I is incorrect but Statement II is correct.


Answer: C

Solution:

Phenol are weakly acidic. Phenol is more acidic than alcohol & H₂O statement (I) is correct. (II) is incorrect.

Question59


In the given reaction,



the number of sp² hybridised carbon(s) in compound ' X ' is ____
[29-Jun-2022-Shift-2]


Answer: 8

Solution:



Question60


The intermediate X, in the reaction :




[25-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: B

Solution:



Question61


Hex-4-ene-2-ol on treatment with PCC gives 'A'. 'A' on reaction with sodium hypoiodite gives 'B', which on further heating with soda lime gives 'C'. The compound 'C' is :
[24-Jun-2022-Shift-2]

Options:

- A. 2-pentene
- B. propanaldehyde
- C. 2-butene
- D. 4-methylpent-2-ene


Answer: B

Solution:



Question62


In the following reaction :




The compounds A and B respectively are :
[25-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: D


Solution:

Given reaction is cumene-Peroxide method for the preparation of phenol. In this reaction



Question63


Amongst the following, the major product of the given chemical reaction is




[25-Jun-2022-Shift-2]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question64


The major product formed in the following reaction, is




[25-Jun-2022-Shift-2]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question65




Consider the above reaction and identify the intermediate ' X '


[26-Jun-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: A

Solution:



Question66

Oxidation of toluene to benzaldehyde can be easily carried out with which of the following reagents?


[26-Jun-2022-Shift-2]

Options:

- A. CrO_3 /acetic acid, H_3O^+
- B. CrO_3 /acetic anhydride, H_3O^+
- C. KMnO_4 / HCl , H_3O^+
- D. CO / HCl , anhydrous AlCl_3





Answer: B

Solution:



Question67

Which of the following reactions will yield benzaldehyde as a product?

- (A)  c1ccccc1C(=O)O $\xrightarrow[\text{(ii) H}_2/\text{Pd/BaSO}_4]{\text{(i) SOCl}_2, \text{Quinoline}}$
- (B)  c1ccccc1CO $\xrightarrow{\text{CrO}_3/\text{H}_2\text{SO}_4}$
- (C)  c1ccccc1C(=O)OC $\xrightarrow[\text{(ii) PCC}]{\text{(i) NaBH}_4}$
- (D)  Cc1ccccc1 $\xrightarrow[\text{(ii) H}_3\text{O}^+, \Delta]{\text{(i) CrO}_3, (\text{CH}_3\text{CO})_2\text{O}}$


[27-Jun-2022-Shift-1]

Options:

- A. (B) and (C)
 B. (C) and (D)
 C. (A) and (D)
 D. (A) and (C)


Answer: C

Solution:



Question68


Most stable product of the following reaction is:




[25-Jul-2022-Shift-1]

Options:


- A.




B.



C.




D.



Answer: B

Solution:



Question69

Given below are two statements :

Statement I: On heating with KHSO_4 , glycerol is dehydrated and acrolein is formed.

Statement II : Acrolein has fruity odour and can be used to test glycerol's presence.

Choose the correct option.

[25-Jul-2022-Shift-1]

Options:

- A. Both Statement I and Statement II are correct.
- B. Both Statement I and Statement II are incorrect.
- C. Statement I is correct but Statement II is incorrect.
- D. Statement I is incorrect but Statement II is correct.


Answer: B

Solution:

Acrolein has a pungent, suffocating odour.
Acrolein is used to detect presence of glycerol

Question70


Major product of the following reaction is :




[25-Jul-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: D

Solution:



Question71

The difference in the reaction of phenol with bromine in chloroform and bromine in water medium is due to :
[26-Jul-2022-Shift-1]

Options:

- A. Hyperconjugation in substrate
- B. Polarity of solvent
- C. Free radical formation
- D. Electromeric effect the substrate

Answer: B

Solution:



Phenol gives different products with bromine in chloroform and water medium due to the polarity difference between chloroform and water acting as solvent

Question72

A 100 mL solution of $\text{CH}_3\text{CH}_2\text{MgBr}$ on treatment with methanol produces 2.24 mL of a gas at STP. The weight of gas produced is _____ mg. [nearest integer]
[26-Jul-2022-Shift-2]

Answer: 3

Solution:



$$W = n \times M$$

$$= 10^{-4} \times 30 = 3 \text{ mg}$$

Question 73


When ethanol is heated with conc. H_2SO_4 , a gas is produced. The compound formed, when this gas is treated with cold dilute aqueous solution of Baeyer's reagent, is
[29-Jul-2022-Shift-2]

Options:

- A. formaldehyde
- B. formic acid
- C. glycol
- D. ethanoic acid


Answer: C

Solution:



Question 74


Identify the major products A and B for the below given reaction sequence.




[28-Jul-2022-Shift-1]

Options:


- A.




B.



C.




D.




Answer: B

Solution:



Question 75


The major product in the given reaction is




[28-Jul-2022-Shift-2]

Options:

A.




B.



C.




D.




Answer: C

Solution:



Question76


Compound I is heated with Conc. HI to give a hydroxy compound A which is further heated with Zn dust to give compound B. Identify A and B.




[28-Jul-2022-Shift-2]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question77


Consider the above reaction sequence, the Product ' C ' is :




[29-Jul-2022-Shift-1]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question78


A compound 'X' is acidic and it is soluble in NaOH solution, but insoluble in NaHCO₃ solution. Compound 'X' also gives violet colour with neutral FeCl₃ solution. The compound 'X' is :
[29-Jul-2022-Shift-1]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question79


Identify A in the given reaction.




[26 Feb 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: B


Solution:

The given reaction is an application of Darzen's reaction in which an aliphatic alcohol (preferably 1° or 2°) gets converted into the respective alkyl chloride (in pure form) through S_N2 pathway when it uses

pyridine  or as organic base.




Note Phenolic - OH group does not respond to this reaction as +R effect of phenolic —OH group gives double bond character to the C — O bond.




Mechanism


- (i) Alcohol reacts with thionyl chloride to give chlorosulphite and HCl.



- (ii) Pyridine reacts with HCl to give pyridinium chloride.



- (iii) Chlorosulphite reacts with chloride ion via bimolecular nucleophilic substitution reaction to give alkyl chloride.



Question 80

Ceric ammonium nitrate and $CHCl_3$ /alc. KOH are used for the identification of functional groups present in and respectively.
[26 Feb 2021 Shift 2]

Options:


- A. alcohol, phenol
- B. amine, alcohol
- C. alcohol, amine
- D. amine, phenol

Answer: C

Solution:

Ceric ammonium nitrate and $CHCl_3$ /alc. KOH are used for the identification of functional groups present in alcohol and amine respectively.


Ceric ammonium nitrate (CAN) test When an alcohol reacts with few drops of CAN, a red complex ammonium nitrate is developed which confirms presence of alcoholic $-OH$ group.



Test with $CHCl_3$ and alcoholic KOH

[Carbylamine test or Saytzeff's isocyanide test]


When a primary amine (aliphatic or aromatic) is heated with alcoholic KOH and chloroform ($CHCl_3$), the isocyanide (carbylamine) is formed as indicated by a foul order.



Secondary and tertiary amines do not respond to carbylamine test.

Question81


Identify the major products A and B respectively in the following reactions of phenol.




[26 Feb 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: B


Solution:

Phenol on reaction with Br_2 in $\text{CS}_2/273\text{K}$ undergoes an electrophilic substitution reaction by Br^+ (electrophile) in aprotic solvent CS_2 to give 4-bromophenol as the major product.



Phenol on reaction with CHCl_3 , NaOH followed by hydrolysis gives Phenol on reaction with CHCl_3 , Na a salicylaldehyde as a major product.

It is Reimer-Tiemann reaction. It is also an electrophilic substitution reaction of phenol by dichlorocarbene CCl_2 (electrophilic).



Question82

Given below are two statements:

Statement I o-nitrophenol is steam volatile due to intramolecular hydrogen bonding.

Statement II o-nitrophenol has high melting due to hydrogen bonding.

In the light of the above statements, choose the most appropriate answer from the options given below.

[26 Feb 2021 Shift 1]


Options:

- A. Statement I is false but statement II is true
- B. Both statement I and statement II are true
- C. Both statement I and statement II are false
- D. Statement I is true but statement II is false

Answer: D

Solution:

Statement I is true. Because of closer proximity (1, 2-positions) of $-\text{OH}$ and $-\text{NO}_2$ groups, o-nitrophenol shows intramolecular hydrogen bonding.




So, o-nitrophenol exists in monomeric state and becomes steam volatile.

Statement II is false, because, due to the presence of intramolecular hydrogen bonding, boiling point and melting point of o-nitrophenol will be lower.


Note p-nitrophenol is the positional isomer of o-nitrophenol. p-nitrophenol shows intermolecular hydrogen bonding and so, it has

higher boiling point, melting point and water solubility.






Question83

What is ' X ' in the given reaction?



[25 Feb 2021 Shift 2]


Options:



Answer: C


Solution:

When ethylene glycol is heated with oxalic acid at 210°C , first we get an unstable cyclic-diester (ethylene oxalate) which readily decarboxylate to give ethylene as the major product.



Question84


What is the final product (major) 'A' in the given reaction?




[24 Feb 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: A

Solution:




Question85

Which of the following compound gives pink colour on reaction with phthalic anhydride in conc. H_2SO_4 followed by treatment with NaOH ?


[24 Feb 2021 Shift 1]

Options:


A.




B.



C.




D.




Answer: A

Solution:




Question86




[16 Mar 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: D

Solution:


Cyanide undergoes nucleophilic addition with Grignard followed by hydrolysis to give carbonyl compound.



Mechanism Carbon of cyanide is electrophilic and is attacked by nucleophilic phenyl anion.



Question87




Consider the above reaction, and choose the correct statement :
[27 Jul 2021 Shift 2]

Options:


- A. The reaction is not possible in acidic medium
- B. Both compounds A and B are formed equally
- C. Compound A will be the major product
- D. Compound B will be the major product

Answer: C

Solution:




Question88




Consider the above reaction, the major product "P" formed is :-
[27 Jul 2021 Shift 2]

Options:


A.




B.



C.




D.




Answer: B

Solution:




Question89




Consider the above reaction and identify the Product P :
[27 Jul 2021 Shift 1]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question90


Consider the above reaction, the major product 'P' is:




[25 Jul 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question91


In the given reaction 3-Bromo-2, 2 -dimethyl butane $\xrightarrow{\text{C}_2\text{H}_5\text{OH}}$ 'A' Product A is:
[20 Jul 2021 Shift 1]
(Major Product)

Options:

- A. 2 -Ethoxy-3, 3 -dimethyl butane
- B. 1-Ethoxy-3, 3-dimethyl butane
- C. 2-Ethoxy-2, 3 -dimethyl butane
- D. 2 -Hydroxy-3, 3 -dimethyl butane

Answer: C

Solution:



Question92

Given below are two statements.

One is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A) Treatment of bromine water with propene yields 1-bromopropan-2-ol.

Reason (R) Attack of water on bromonium ion follows Markownikoff rule and results in 1-bromopropan-2-ol.

In the light of the above statements, choose the most appropriate answer from the options given below.

[31 Aug 2021 Shift 1]


Options:

- A. Both (A) and (R) are true but (R) is not the correct explanation of (A).
- B. (A) is false but (R) is true.
- C. Both (A) and (R) are true and (R) is the correct explanation of (A).
- D. (A) is true but (R) is false

Answer: C

Solution:

Propene reacts with bromine water to yield 1-bromopropan-2-ol.




Markownikoff's rule states that in an unsymmetrical alkene the electron rich nucleophile adds to the C-atom with lesser number of hydrogen.

Hence, both A and R are true and R is the correct explanation of A.

Hence, correct option is (c).

Question93


The major product formed in the following reaction is




[31 Aug 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: B

Solution:


3,3-dimethylbutan-2-ol reacts with concentrated H_2SO_4 to form but-2,3-diene.



Hence, correct option is (b).

Question94


The major product of the following reaction, if it occurs by S_N2 mechanism is




[27 Aug 2021 Shift 2]

Options:


A.




B.



C.



D.




Answer: D

Solution:


In S_N2 reaction, no formation of carbocation takes place. The nucleophile here which is phenoxide attacks from the back side to the carbon bearing the leaving group (bromide ion).



In S_N2 mechanism



Here, in the question



So, here option (d) is correct.

Question95

Given below are two statements

Statement I Ethyl pent-4-yn-oate on reaction with CH_3MgBr gives a 3° alcohol.

Statement II In this reaction, one mole of ethyl pent-4-yn-oate utilizes two moles of CH_3MgBr . In the light of the above statements, choose the most appropriate answer from the options given below.

[27 Aug 2021 Shift 2]


Options:

- A. Both statement I and statement II are false.
- B. Statement I is false but statement II is true.
- C. Statement I is true but statement II is false.
- D. Both statement I and statement II are true.

Answer: C


Solution:

Ethyl pent-4-yn-oate on reaction with CH_3MgBr gives 3° alcohol.



$\therefore$ Statement I is true.

Mechanism



3 molecules of CH_3MgBr is used for this conversion.

$\therefore$ Statement II is false.


Question96

Which one of the following phenols does not give colour when condensed with phthalic anhydride in presence of conc. H_2SO_4 ?


[26 Aug 2021 Shift 2]

Options:


A.




B.



C.



D.




Answer: B

Solution:

In the electrophilic substitution reaction, phenol gives ortho and para substituted product in which para is the major product. The electrophilic substitution reaction of phthalic anhydride with phenol is taking place at para position of the phenol. All the given molecules have free para position except in p-cresol.


Hence, it does not give colour of phthalic anhydride.

The reaction is as follows :



Question97


The correct options for the products A and B of the following reactions are




[26 Aug 2021 Shift 1]

Options:


A.




B.



C.




D.



Answer: B

Solution:

Bromination of phenol with excess of bromine and water will result in multiple substitution around ring as water ionises phenol to phenoxide ion, which activates the ring for trisubstitution, while less polar solvent, i.e. CS_2 will give mono-substituted compound.



Question98

Given below are two statements : one is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A) Synthesis of ethyl phenyl ether may be achieved by Williamson synthesis.

Reason (R) Reaction of bromobenzene with sodium ethoxide yields ethyl phenyl ether.

In the light of the above statements, choose the most appropriate answer from the options given below

[27 Aug 2021 Shift 1]


Options:

- A. Both (A) and (R) are correct and (R) is the correct explanation of (A)
- B. (A) is correct but (R) is incorrect
- C. (A) is incorrect but (R) is correct
- D. Both (A) and (R) are correct but (R) is not the correct explanation of (A)

Answer: B

Solution:


Williamson's synthesis is used to prepare mixed ethers as well as simple ether. To prepare ethyl phenyl ether following reaction take place.



This reaction is not possible as bromine has partial double bond character with benzene ring, so it cannot be dissociated. So, Assertion is correct but Reason is not correct.

Question99


The major product of the following reaction is:




[Jan. 08,2020(I)]

Options:


A.




B.



C.




D.



Answer: C

Solution:



Question100


Arrange the following compounds in increasing order of C – OH bond length: methanol, phenol, p -ethoxyphenol
[Jan. 08,2020(1)]

Options:

- A. methanol < p -ethoxyphenol < phenol
- B. phenol < methanol < p -ethoxyphenol
- C. phenol < p -ethoxyphenol < methanol
- D. methanol < phenol < p -ethoxyphenol

Answer: C

Solution:





Resonance is a deciding factor to determine the order of bond length in given compounds. Phenol exhibits least C – OH bond length due to resonance, whereas methanol will show maximum bond length due to lack of resonance and p -ethoxyphenol will have some intermediate value of bond length.

Question101



Among the compounds A and B with molecular formula C₉H₁₈O₃, A is having higher boiling point than B. The possible structures of A and B are:
[Jan. 08,2020(II)]

Options:



- A.





B.



C.



D.



Answer: A


Solution:

In (A), -OH group is present, so inter-molecular Hbonding is possible while in (B), due to methoxy group there is no possibility of Inter-molecular H-bonding. So A is having higher boiling point than B.


1-methyl ethylene oxide when treated with an excess of HBr produces:
[Jan. 07,2020(I)]

Options:

A.




B.



C.




D.




Answer: B

Solution:



Question103


The major product of the following reaction is:




[Sep. 05,2020(II)]

Options:


A.




B.



C.




D.




Answer: A

Solution:







Question104

The major product [B] in the following reactions is:




Options:



Answer: B

Solution:




Question105

When neopentyl alcohol is heated with an acid, it slowly converted into an 85: 15 mixture of alkenes A and B, respectively. What are these alkenes?


[Sep. 04,2020(I)]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question106




The number of chiral centres present in [B] is _____.



[Sep. 04,2020(I)]

Answer: 4

Solution:



Number of chiral centres, represented as (*) in the product (B) = 4


Question107

A solution of phenol in chloroform when treated with aqueous NaOH gives compound P as a major product. The mass percentage of carbon in P is (to the nearest integer) (Atomic mass: C = 12; H = 1; O = 16)

[NV, Sep. 06,2020 (II)]

Answer: 69

Solution:




Molecular formula of product 'P' = $\text{C}_7\text{H}_6\text{O}_2$

So, mass % of C in 'P'

$$= \frac{12 \times 7}{84 + 6 + 32} \times 100 = 68.85\% \approx 69\%$$

Question 108

The increasing order of boiling points of the following compounds is :



[Sep. 05,2020 (II)]


Options:

- A. I < III < IV < II
- B. I < IV < III < II
- C. IV < I < II < III
- D. III < I < II < IV


Answer: B

Solution:


(II) and (III) compounds almost have same boiling point. In the given options, option (b) will be the correct answer.




B.P. $\rightarrow 202^{\circ}\text{C}$



B.P. $\rightarrow 279^{\circ}\text{C}$




B.P. $\rightarrow 284^{\circ}\text{C}$



B.P. $\rightarrow 243^{\circ}\text{C}$

Question109

Consider the following reaction:




$\xrightarrow[\text{anhydride}]{\text{Chromic}}$ ► 'p' The product 'P' gives positive ceric ammonium nitrate test. This is because of the presence of which of these –OH group(s)?
[Sep. 03,2020 (II)]

Options:

- A. (ii) only
- B. (iii) and (iv)
- C. (iv) only
- D. (ii) and (iv)

Answer: A


Solution:



Generally CAN test is done for alcohols which give pink or red colour. But for phenols and phenolic compounds it gives brown or black colour. So, this test helps to differentiate phenols from alcohols.

Question110


The major product of the following reaction is :




[Sep. 02,2020 (II)]

Options:


A.




B.



C.




D.



Answer: C


Solution:



It is an example of electrophilic substitution reaction. Position of electrophile is directed by the strong ring activating group ($-\text{OH}$), present in the ring.

Question111


The major aromatic product C in the following reaction sequence will be :




[Sep. 02,2020 (I)]

Options:


A.




B.



C.




D.



Answer: A

Solution:




Question112


An organic compound 'A' (C₉H₁₀O) when treated with conc. HI undergoes cleavage to yield compounds 'B' and 'C'. 'B' gives yellow precipitate with AgN O₃ where as 'C' tautomerizes to 'D'. 'D' gives positive iodoform test. 'A' could be:
[Sep. 02,2020 (II)]

Options:


A.




B.



C.





D.



Answer: C

Solution:



Question113

$\text{CH}_3\text{CH}_2 - \underset{\text{Ph}}{\overset{\text{OH}}{\text{C}}} - \text{CH}_3$ cannot be prepared by:
[Jan. 12,2019 (I)]


Options:

- A. $\text{CH}_3\text{CH}_2\text{COCH}_3 + \text{PhMgX}$
- B. $\text{PhCOCH}_2\text{CH}_3 + \text{CH}_3 \text{MgX}$
- C. $\text{PhCOCH}_3 + \text{CH}_3\text{CH}_2 \text{MgX}$
- D. $\text{HCHO} + \text{PhCH}(\text{CH}_3)\text{CH}_2 \text{MgX}$


Answer: D

Solution:

Tertiary alcohol is prepared by the reaction of Grignard reagent with a ketone (formaldehyde is used to prepare primary alcohol).




Question114




[Jan. 10,2019 (I)]

Options:


A.




B.



C.




D.



Answer: D


Solution:

NaBH_4 is a selective reducing agent, used for the reduction of aldehydes and ketones, it does not affect alkene and ester.



Question115

Which is the most suitable reagent for the following transformation?



[Jan. 10, 2019 (II)]

Options:

A. Tollen's reagent

B. I_2/NaOH


C. $\text{CrO}_2\text{Cl}_2/\text{CS}_2$

D. alkaline KMnO_4

Answer: B


Solution:

The most suitable reagent for the given reaction is I_2/NaOH (Iodoform reaction).



Question116


The major product of the following reaction is:




[Jan. 11,2019(1)]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question117


The major product of the following reaction is:




[Jan. 11,2019(I)]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question118


The major product of the following reaction is:




[Jan. 11,2019(II)]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question119

The increasing order of the pK_a values of the following compounds is:



[Jan. 10,2019(I)]

Options:


- A. $C < B < A < D$
- B. $B < C < D < A$
- C. $D < A < C < B$
- D. $B < C < A < D$

Answer: D

Solution:

Electron withdrawing substituents increase the acidic strength, while electron releasing groups decrease the acidic strength.


$$\therefore \text{Acidic strength} \propto K_a \propto \frac{1}{pK_a}$$



$\text{pK}_a : \text{B} < \text{C} < \text{A} < \text{D}$

Question120


The major product of the following reaction is:




[Jan. 10,2019(II)]

Options:


A.




B.



C.




D.



Answer: B

Solution:

Reaction involved:




Question121


The products formed in the reaction of cumene with O_2 followed by treatment with dil. HCl are:
[Jan. 9,2019(II)]

Options:


A.




B.



C.




D.



Answer: C


Solution:

Reaction involved:



Question122


The major product of the following reaction is:




[Jan. 12,2019(I)]

Options:


A.




B.



C.




D.




Answer: D

Solution:



Question123


The major product in the following conversion is:




[Jan. 12, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: B

Solution:



Question 124


The major product of the following reaction is:




[Jan. 10,2019(I)]

Options:


A.




B.



C.




D.



Answer: B


Solution:




Heating of 2-chloro-1-phenylbutane with EtOK/EtOH gives X as the major product. Reaction of X with $\text{Hg}(\text{OAc})_2/\text{H}_2\text{O}$ followed by NaBH_4 gives Y as the major product. Y is:
[April 12, 2019 (II)]

Options:


A.



B.



C.




D.



Answer: C

Solution:




Question126


What will be the major product when m-cresol is reacted with propargyl bromide ($\text{HC}\equiv\text{C}-\text{CH}_2\text{Br}$) in presence of K_2CO_3 in acetone?
[April 12, 2019 (II)]

Options:


A.




B.



C.




D.




Answer: A

Solution:



Question127


The major product of the following reaction is:




[April 10,2019 (I)]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question128


The major product obtained in the given reaction is:




[April 10,2019 (II)]

Options:


A.




B.



C.




D.




Answer: C

Solution:



Question129


The major product of the following reaction is:




[April 8,2019 (I)]

Options:


A.




B.



C.




D.



Answer: B

Solution:




Question130


Phenol on treatment with CO_2 in the presence of NaOH followed by acidification produces compound X as the major product. X on treatment with $(\text{CH}_3\text{CO})_2\text{O}$ in the presence of catalytic amount of H_2SO_4 produces :
[2018]

Options:


A.




B.



C.




D.



Answer: A

Solution:




Question131

Phenol reacts with methyl chloroformate in the presence of NaOH to form product A. A reacts with Br_2 to form product B. A and B are respectively:


[2018]

Options:


A.




B.



C.






D.




Answer: C

Solution:



Question132


The major product formed in the following reaction is:




[2018]

Options:


A.




B.



C.




D.



Answer: D

Solution:




Question133


Which of the following, upon treatment with tert-BuONa followed by addition of bromine water, fails to decolourize the colour of bromine?
[2017]

Options:


A.




B.



C.











D.




Answer: A

Solution:




The major product of the following reaction is:




[Online April 8,2017]

Options:


A.




B.



C.




D.



Answer: A

Solution:



Question135


The gas evolved on heating CH_3MgBr in methanol is:
[Online April. 9,2016]

Options:

- A. Methane
- B. Ethane
- C. Propane
- D. HBr


Answer: A

Solution:



Question136


Consider the reaction sequence below:




[Online April 10,2016]

Options:


A.




B.



C.




D.



Answer: D

Solution:



Question137

The most suitable reagent for the conversion of $R-CH_2-OH \rightarrow R-CHO$ is:

[2014]

Options:

A. $KMnO_4$

B. $K_2Cr_2O_7$


C. CrO_3

D. PCC (Pyridinium chlorochromate)

Answer: D

Solution:

An excellent reagent for oxidation of 1° alcohols to aldehydes is PCC (consult Q.3 also).



Question138

Which one of the following statements is not correct?

[Online April 11, 2014]

Options:

- A. Alcohols are weaker acids than water
- B. Acid strength of alcohols decreases in the following $\text{RCH}_2\text{OH} > \text{R}_2\text{CHOH} > \text{R}_3\text{COH}$
- C. Carbon-oxygen bond length in methanol, CH_3OH is shorter than that of $\text{C}-\text{O}$ bond length in phenol.
- D. The bond angle in methanol is 108.9° .

Answer: C

Solution:

The $\text{C}-\text{O}$ bond length in alcohols is 142pm and in Phenol it is 136pm. The $\text{C}-\text{O}$ bond length in phenol is shorter than that in methanol due to the conjugation of unshared pair of electrons on oxygen with the ring, which imparts double bond character to the $\text{C}-\text{O}$ bond.

Question139

In the Victor-Meyer's test, the colour given by 1° , 2° and 3° alcohols are respectively:


[Online April 12, 2014]

Options:


- A. Red, colourless, blue
- B. Red, blue, colourless
- C. Colourless, red, blue
- D. Red, blue, violet

Answer: B

Solution:




(ii)



Question 140


Sodium phenoxide when heated with CO_2 under pressure at 125°C yields a product which on acetylation produces C




The major product C would be
[2014]

Options:


A.




B.



C.






D.




Answer: A

Solution:



Question141

The following reaction



is known as:




[Online April 11,2014]

Options:

- A. Perkin reaction
- B. Gatterman-Koch Formylation
- C. Kolbe's reaction
- D. Gattermann reaction

Answer: D


Solution:



Note : Gattermann Koch reaction is a variation of Gattermann reaction, and this reaction involves the use of carbon monoxide instead of HCN.

Question142

Which one of the following substituents at para-position is most effective in stabilizing the phenoxide



[Online April 19,2014]

Options:

- A. $-\text{CH}_3$
- B. $-\text{OCH}_3$
- C. $-\text{COCH}_3$
- D. $-\text{CH}_2\text{OH}$

Answer: C

Solution:

Electron withdrawing group stabilises the benzene ring due to delocalisation of charge.

$-\text{CH}_3$ and $-\text{CH}_2\text{OH}$ are electron donating group and hence decrease the stability of benzene ring $-\text{OCH}_3$ is weaker electron withdrawing group than $-\text{COCH}_3$. Hence COCH_3 group more stabilizes the phenoxide ion at p- position.

Question143

Which one of the following compounds will not be soluble in sodium bicarbonate?

[Online April 19,2014]


Options:

- A. 2,4,6 - Trinitrophenol
- B. Benzoicacid
- C. o - Nitrophenol
- D. Benzene sulphonic acid

Answer: B

Solution:

Due to intramolecular hydrogen bonding this will not be soluble in sodium bicarbonate.



Question144


**Allyl phenyl ether can be prepared by heating:
[Online April 9,2014]**

Options:

- A. $\text{C}_6\text{H}_5\text{Br} + \text{CH}_2 = \text{CH} - \text{CH}_2 - \text{ONa}$
- B. $\text{CH}_2 = \text{CH} - \text{CH}_2 - \text{Br} + \text{C}_6\text{H}_5\text{ONa}$
- C. $\text{C}_6\text{H}_5 - \text{CH} = \text{CH} - \text{Br} + \text{CH}_3 - \text{ONa}$
- D. $\text{CH}_2 = \text{CH} - \text{Br} + \text{C}_6\text{H}_5 - \text{CH}_2 - \text{ONa}$

Answer: B

Solution:



Question145

**Williamson synthesis of ether is an example of:
[Online April 19,2014]**

Options:

- A. Nucleophilic addition
- B. Electrophilic addition
- C. Electrophilic substitution
- D. Nucleophilic substitution

Answer: D

Solution:

This method is suitable for the preparation of a wide variety of unsymmetrical ethers. The nucleophilic substitution of halides with alkoxide leads to desired product.

Question146

**An unknown alcohol is treated with the "Lucas reagent" to determine whether the alcohol is primary, secondary or tertiary. Which alcohol reacts fastest and by what mechanism:
[2013]**

Options:

- A. secondary alcohol by $\text{S}_{\text{N}}1$
- B. tertiary alcohol by $\text{S}_{\text{N}}1$

C. secondary alcohol by S_N^2

D. tertiary alcohol by S_N^2

Answer: B

Solution:

Reaction of alcohols with Lucas reagent proceeds through carbocation formation. Further 3° carbocations (from tertiary alcohols) are highly stable thus reaction proceeds through S_N1 mechanism.

Question147

**Rate of dehydration of alcohols follows the order:
[Online April 9, 2013]**

Options:

A. $2^\circ > 1^\circ > \text{CH}_3\text{OH} > 3^\circ$

B. $3^\circ > 2^\circ > 1^\circ > \text{CH}_3\text{OH}$

C. $2^\circ > 3^\circ > 1^\circ > \text{CH}_3\text{OH}$

D. $\text{CH}_3\text{OH} > 1^\circ > 2^\circ > 3^\circ$

Answer: B

Solution:

The order of dehydration among three type of alcohols is $3^\circ > 2^\circ > 1^\circ > \text{CH}_3\text{OH}$. This behaviour is related to the relative stabilities of carbocations ($3^\circ > 2^\circ > 1^\circ$).

Question148

**Amongst the following alcohols which would react fastest with conc. HCl and ZnCl₂ ?
[Online April 22, 2013]**

Options:

A. pentanol

B. 2-methyl butanol

C. 2-pentanol

D. 2-methyl butan-2-ol

Answer: D


Solution:

$\text{ZnCl}_2 + \text{conc. HCl}$ is Lucas reagent. Lucas reagent reacts fastest with tertiary alcohol.

3° alcohol + Lucas reagent = Immediate turbidity

Question149

Arrange the following compounds in order of decreasing acidity:



[2013]

Options:

A. $\text{II} > \text{IV} > \text{I} > \text{III}$

B. $\text{I} > \text{II} > \text{III} > \text{IV}$


C. $\text{III} > \text{I} > \text{II} > \text{IV}$

D. $\text{IV} > \text{III} > \text{I} > \text{II}$

Answer: C

Solution:

Electron withdrawing substituents like $-\text{NO}_2$, $-\text{Cl}$ increase the acidity of phenol while electron releasing substituents like $-\text{CH}_3$, $-\text{OCH}_3$ decrease acidity. hence the correct order of acidity will be




$(-\text{M}, -\text{I})$ $(-\text{I} > +\text{M})$ $(+\text{I}, +\text{H.C})$ $(+\text{M})$

Question150


What is the structure of the major product when phenol is treated with bromine water?
[Online April 22,2013]

Options:


A.




B.



C.




D.



Answer: A

Solution:

Phenol has activating (electron releasing) – OH group and bromine water supplies Br^+ ion easily, hence under such conditions reaction does not stop at monobromo or dibromo stage but a fully brominated (2, 4, 6, -tribromophenol compound is the final product.



Question151


The reaction of phenol with benzoyl chloride to give phenyl benzoate is known as:
[Online April 23,2013]

Options:

- A. Claisen reaction
- B. Schotten-Baumann reaction
- C. Reimer-Tiemann reaction
- D. Gatterman-Koch reaction


Answer: B

Solution:



Question152


The major product in the following reaction is :




[Online April 25,2013]

Options:


A.




B.



C.



D.



Answer: C

Question153


Phenol on heating with CHCl_3 and NaOH gives salicylaldehyde. The reaction is called:
[Online April 25 ,2013]

Options:

- A. Reimer - Tiemann reaction
- B. Claisen reaction
- C. Cannizzaro's reaction
- D. Hell - Volhard - Zelinsky reaction

Answer: A

Solution:



Question154


An ether (A), $\text{C}_5\text{H}_{12}\text{O}$, when heated with excess of hot concentrated HI produced two alkyl halides which when treated with NaOH yielded compounds (B) and (C) Oxidation of (B) and (C) gave a propanone and an ethanoic acid respectively. The IUPAC name of the ether (A) is :
[Online April 9,2013]

Options:


- A. 2 -ethoxypropane
- B. ethoxypropane
- C. methoxybutane
- D. 2 -methoxybutane

Answer: A

Solution:



Hence the IUPAC name of compound (A) is



Question155

In Williamson synthesis of mixed ether having a primary and a tertiary alkyl group if tertiary halide is used, then:

[Online April 22,2013]


Options:

- A. Rate of reaction will be slow due to slow cleavage of carbon-halogen bond.
- B. Alkene will be the main product.
- C. Simple ether will form instead of mixed ether.
- D. Expected mixed ether will be formed.

Answer: B

Solution:

The tertiary alkyl halide undergo elimination reaction to give alkenes.



Question156

Reagent used to convert allyl alcohol to acrolein is

[Online May 19, 2012]

Options:

- A. MnO_2
- B. H_2O_2
- C. OsO_4
- D. KMnO_4

Answer: A

Solution:

MnO₂ being a mild oxidising agent stops the oxidation of -CH₂OH group at aldehyde stage.

Question157

Consider thiol anion ($\text{RS}^{\ominus}$) and alkoxy anion ($\text{RO}^{\ominus}$). Which of the following statements is correct?

[2011RS]

Options:

- A. $\text{RS}^{\ominus}$ is less basic but more nucleophilic than $\text{RO}^{\ominus}$
- B. $\text{RS}^{\ominus}$ is more basic and more nucleophilic than $\text{RO}^{\ominus}$
- C. $\text{RS}^{\ominus}$ is more basic but less nucleophilic than $\text{RO}^{\ominus}$
- D. $\text{RS}^{\ominus}$ is less basic and less nucleophilic than $\text{RO}^{\ominus}$

Answer: A

Solution:

On moving down a group, the basicity & nucleophilicity are inversely related, i.e. nucleophilicity increases while basicity decreases. i.e $\text{RS}^{\ominus}$ is more nucleophilic but less basic than $\text{RO}^{\ominus}$. This opposite behaviour is because of the fact that basicity and nucleophilicity depends upon different factors. Basicity is directly related to the strength of the H -element bond, while nucleophilicity is indirectly related to the electronegativity of the atom to which proton is attached.

Question158

The correct order of acid strength of the following compounds:

(A) Phenol

(B) p -Cresol

(C) m - Nitrophenol

(D) p -Nitrophenol


[2011 RS]

Options:

- A. $\text{D} > \text{C} > \text{A} > \text{B}$
- B. $\text{B} > \text{D} > \text{A} > \text{C}$
- C. $\text{A} > \text{B} > \text{D} > \text{C}$
- D. $\text{C} > \text{B} > \text{A} > \text{D}$

Answer: A

Solution:



Electron withdrawing substituents increase the acidity of phenols; while electron releasing substituents decrease acidity. Thus the correct order is $D > C > A > B$

Question159

Consider the following reaction :




Among the following, which one cannot be formed as a product under any conditions?
[2011RS]

Options:

- A. Ethylene
- B. Acetylene
- C. Diethyl ether
- D. Ethylhydrogen sulphate

Answer: B

Solution:



Acetylene is not formed under any condition.

Question160

From amongst the following alcohols, the one that would react fastest with conc. HCl and anhydrous ZnCl_2 , is

[2010]

Options:




- A. 2 -Butanol
- B. 2 - Methylpropan-2-ol
- C. 2-Methylpropanol
- D. 1 - Butanol

Answer: B

Solution:

Tertiary alcohols react fastest with conc. HCl and anhydrous ZnCl₂ (Lucas reagent) as its mechanism proceeds through the formation of stable tertiary carbocation.

Mechanism :




Question161

The main product of the following reaction is $\text{C}_6\text{H}_5\text{CH}_2\text{CH}(\text{OH})\text{CH}(\text{CH}_3)_2 \xrightarrow{\text{conc. H}_2\text{SO}_4} ?$


[2010]

Options:


A.




B.



C.




D.



Answer: A

Solution:

Whenever dehydration can produce two different alkenes, major product is formed according to Saytzeff rule i.e. more substituted alkene (alkene having lesser number of hydrogen atoms on the two doubly bonded carbon atoms) is the major product. Such reactions which can produce two or more structural isomers but one of them in greater amounts than the other are called regioselective ; in case a reaction is 100% regioselective, it is termed as regiospecific. In addition to being regioselective, alcohol dehydrations are stereoselective (a reaction in which a single starting material can yield two or more stereoisomeric products, but gives one of them in greater amount than any other).



Question162


The major product obtained on interaction of phenol with sodium hydroxide and carbon dioxide is
[2009]

Options:

- A. salicylaldehyde
- B. salicylic acid
- C. phthalic acid
- D. benzoic acid

Answer: B

Solution:



Question163

Phenol, when it reacts first with concentrated sulphuric acid and then with concentrated nitric acid, gives
[2008]


Options:

- A. 2,4,6 -trinitrobenzene
- B. o -nitrophenol
- C. p -nitrophenol
- D. nitrobenzene

Answer: B


Solution:

Phenol on reaction with conc. H_2SO_4 gives a mixture of o - and p - products (i.e., $-\text{SO}_3\text{H}$ group occupies o-, p⁻ position). At room temperature, o -product is more stable, which on treatment with conc. HNO_3 will yield onitrophenol.



Question164

In the following sequence of reactions,




Options:

- A. propanal
- B. butanal
- C. n -butyl alcohol
- D. n -propyl alcohol.

Answer: D

Solution:





Question165

Among the following the one that gives positive iodoform test upon reaction with I_2 and NaOH is

[2006]

Options:



Answer: B

Solution:

Ethanol and only those 2° alcohols which contain CH(OH)CH_3 group undergo haloform reaction.


Question166

The structure of the compound that gives a tribromo derivative on treatment with bromine water is


[2006]

Options:


A.




B.



C.



D.




Answer: C

Solution:

In (c) both groups are activating and undergo electrophilic substitution in the same positions.



Question167



The electrophile involved in the above reaction is
[2006]

Options:

A. trichloromethyl anion $\left(\overset{\ominus}{\text{C}}\text{Cl}_3\right)$.

B. formyl cation


C. phenoxide ion

D. dichlorocarbene

Answer: D

Solution:

This is Reimer-Tiemann reaction and the electrophile is dichlorocarbene.



Question168

HBr reacts with $\text{CH}_2=\text{CH}-\text{OCH}_3$ under anhydrous conditions at room temperature to give
[2006]

Options:

A. $\text{BrCH}_2-\text{CH}_2-\text{OCH}_3$

B. $\text{H}_3\text{C}-\text{CHBr}-\text{OCH}_3$


C. CH_3CHO and CH_3Br

D. BrCH_2CHO and CH_3OH

Answer: B

Solution:

Methyl vinyl ether under anhydrous condition at room temperature undergoes addition reaction.



Question169


The best reagent to convert pent- 3 – en – 2 – ol into pent- 3 -en2-one is [2005]

Options:

- A. Pyridinium chlorochromate
- B. Chromic anhydride in glacial acetic acid
- C. acidic dichromate
- D. Acidic permanganate

Answer: A

Solution:






Pyridinium chlorochromate (PCC) oxidises 1° and 2° alcohols to aldehyde and ketones.

Question170

Among the following compounds which can be dehydrated very easily? [2004]


Options:



Answer: A

Solution:

3-Methyl pentan-3-ol will be dehydrated most readily since it produces a very stable, tertiary carbonium ion as intermediate.



Question171

During dehydration of alcohols to alkenes by heating with conc. H_2SO_4 , the initiation step is
[2003]

Options:

- A. formation of carbocation
- B. elimination of water
- C. formation of an ester
- D. protonation of alcohol molecule


Answer: D

Solution:


The dehydration of alcohol to form alkene occurs in following three steps.

Step (1) is initiation step.


Step(1) Formation of protonated alcohol.



Step(2) Formation of carbocation



Step(3) Elimination of a proton to form alkene




Question172


Picric acid is:
[2002]

Options:


- A.




B.



C.



D.



Answer: C

Solution:

2,4,6 Trinitrophenol is also known as picric acid.
