

# WEIGHTED ERDŐS–KAC THEOREMS VIA COMPUTING MOMENTS

KAI (STEVE) FAN

**ABSTRACT.** By adapting the moment method developed by Granville and Soundararajan [18], Khan, Milinovich and Subedi [25] recently obtained a weighted version of the Erdős–Kac theorem for  $\omega(n)$  with multiplicative weight  $d_k(n)$ , where  $\omega(n)$  denotes the number of distinct prime divisors of a positive integer  $n$ , and  $d_k(n)$  is the  $k$ -fold divisor function with  $k \in \mathbb{N}$ . In this paper, we generalize their method to study the distribution of additive functions  $f(n)$  weighted by nonnegative multiplicative functions  $\alpha(n)$  in a wide class. In particular, we establish uniform asymptotic formulas for the moments of  $f(n)$  with suitable growth rates. We also prove a qualitative result on the moments which extends a theorem of Delange and Halberstam [9]. As a consequence, we obtain a weighted analogue of the Kubilius–Shapiro theorem.

## CONTENTS

1. Introduction	1
2. Main Results	4
3. Mean Values of Multiplicative Functions	11
4. Computing Moments	21
5. Estimation of $G(z)$ and $D(y, z)$	24
6. Estimation of $E(y, z, w; m)$	29
7. Deduction of Theorems 2.1 and 2.2	33
8. Proof of Theorem 2.3 (sketch)	38
9. Proofs of Theorem 2.4 and Corollary 2.5 (sketch)	41
10. Concluding Remarks	49
References	51

## 1. INTRODUCTION

The celebrated Erdős–Kac theorem, first proved by Erdős and Kac [16] in 1940, states that if  $\omega(n)$  denotes the number of distinct prime divisors of a positive integer  $n$ , then

$$\lim_{x \rightarrow \infty} \frac{1}{x} \cdot \# \left\{ n \leq x : \frac{\omega(n) - \log \log n}{\sqrt{\log \log n}} \leq V \right\} = \Phi(V) \quad (1.1)$$

for any given  $V \in \mathbb{R}$ , where

$$\Phi(V) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^V e^{-v^2/2} dv$$

---

2020 *Mathematics Subject Classification.* Primary: 11N60, 11K65; Secondary: 11N37.

*Key words and phrases.* The Erdős–Kac theorem, Gaussian distribution, method of moments, mean values of multiplicative functions.

is the cumulative distribution function of the standard Gaussian distribution. This statistical result is a direct upgrade of an earlier theorem of Hardy and Ramanujan on the *normal order* of  $\omega$  (see [22] and [23, Theorem 431]), which asserts that given any  $\epsilon > 0$ , the inequality  $|\omega(n) - \log \log n| < \epsilon \log \log n$  holds for all but  $o(x)$  values of  $n \leq x$ . In fact, Erdős and Kac proved in the same paper a more general result in which the function  $\omega$  is replaced by any strongly additive function  $f$  that is bounded on primes and admits an unbounded “variance”  $\sum_{p \leq x} f(p)^2/p$ . Recall that an arithmetic function  $f: \mathbb{N} \rightarrow \mathbb{C}$  is *additive* if  $f(mn) = f(m) + f(n)$  for all positive integers  $m, n \in \mathbb{N}$  with  $\gcd(m, n) = 1$ , and it is *strongly additive* if it also satisfies  $f(p^\nu) = f(p)$  for all prime powers  $p^\nu$ . Thus, strongly additive functions are completely determined by their values at primes, which makes them a particularly nice subclass of additive functions. In fact, it can be shown that (1.1) also holds for  $\Omega(n)$  in place of  $\omega(n)$ , where  $\Omega(n)$  denotes the total number of prime factors of  $n$ , counting multiplicity, by exploiting the fact that  $\omega$  and its cousin  $\Omega$  do not differ very much on average:

$$\sum_{n \leq x} (\Omega(n) - \omega(n)) = O(x). \quad (1.2)$$

The original proof of the Erdős–Kac theorem (1.1) combines the central limit theorem with Brun’s sieve and is quite complicated. Later, LeVeque [26, Theorem 1] introduced some modifications and obtained a quantitative version of (1.1) with a rate of convergence  $O(\log \log \log x / \sqrt{\log \log x})$ . Using a deep analytic approach, Rényi and Turán [29] improved upon LeVeque’s result with a rate of convergence  $O(1/\sqrt{\log \log x})$ , which is best possible in the sense that it cannot be improved to  $o(1/\sqrt{\log \log x})$  without loss of uniformity in  $V$ .

A third approach to (1.1), first suggested by Kac [24], examines the moments of  $\omega$ . In probability theory, the moments of a random variable  $X$  often provides valuable information about its distribution. For example, an application of Markov’s inequality yields  $\mathbb{P}(|X| > c) \leq \mathbb{E}|X|^k / c^k$ . Given all the moments  $\mathbb{E}|X|^k < \infty$ , one may select  $k$  that minimizes this tail estimate. If  $X$  happens to obey a Gaussian law, then it is completely determined by its moments, a direct consequence of [4, Theorem 30.1] or [11, Theorem 3.3.26]. Consequently, by [4, Theorem 30.2], one reduces the proof of (1.1) to that of the asymptotic formula

$$\frac{1}{x} \sum_{n \leq x} (\omega(n) - \log \log x)^m = (\mu_m + o(1)) (\log \log x)^{\frac{m}{2}} \quad (1.3)$$

for every  $m \in \mathbb{N}$ . Here  $\mu_m$  is the  $m$ th moment of a standard Gaussian distribution given by

$$\mu_m = \begin{cases} m! / m!! , & \text{if } 2 \mid m, \\ 0, & \text{otherwise.} \end{cases}$$

The case  $m = 1$  follows from Mertens’ theorem [23, Theorem 427], and the case  $m = 2$  was settled by Turán [35]. Early proofs of (1.3) for all  $k$  are due to Delange [7] in 1953 and Halberstam [19] in 1955, both of which are very complicated. Later, Delange [8] provided an elementary proof of (1.3) for strongly additive functions. Halberstam’s proof was also simplified and rendered more transparent by Billingsley [3] in 1969. In 2007, Granville and Soundararajan [18] derived asymptotic formulas for the moments which hold uniformly in the range  $m \leq (\log \log x)^{1/3}$ . Their method, which may be viewed to some extent as a clever and efficient repackaging of Billingsley’s, is so flexible that it can be modified to study the distribution of values of additive functions in a rather general sieve-theoretic framework.

More generally, one can study the distribution of values of  $\omega(n)$  weighted by certain nonnegative multiplicative functions  $\alpha(n)$ . For instance, Elliott [14] showed, based on the Landau–Selberg–Delange method, that

$$\lim_{x \rightarrow \infty} \left( \sum_{n \leq x} d(n)^c \right)^{-1} \sum_{\substack{n \leq x \\ \omega(n) \leq 2^c \log \log x + V \sqrt{2^c \log \log x}}} d(n)^c = \Phi(V)$$

for any given  $c \in \mathbb{R}$  and  $V \in \mathbb{R}$ , where  $d(n)$  denotes the number of positive divisors of  $n$ . Building on the method of Granville and Soundararajan, Khan, Milinovich and Subedi [25] recently proved an analogue for the weight  $d_k(n)$  with mean  $k \log \log x$  and variance  $k \log \log x$ , where  $d_k(n) := \# \{ (a_1, \dots, a_k) \in \mathbb{N}^k : a_1 \cdots a_k = n \}$  is the  $k$ -fold divisor function. Concrete examples of a different flavor were also obtained by Alladi [1, 2] who proved weighted Erdős–Kac theorems for  $\omega(n)$  with  $n$  running over rough and smooth numbers, respectively. Weighted versions of the Erdős–Kac theorem with general nonnegative multiplicative weights have also been obtained by Elboim and Gorodetsky [12] and Tenenbaum [33, 34]. The argument of Elboim and Gorodetsky adapts Billingsley’s and is simpler, whereas Tenenbaum’s results allow for general additive functions and a larger class of multiplicative weights and provide estimates for the rate of convergence.

The main purpose of this paper is to establish weighted versions of the Erdős–Kac theorem by pushing the method of moments of Granville and Soundararaja to its limit. Our work is the first to apply this method to prove weighted Erdős–Kac theorems with general additive functions and multiplicative weights. We obtain uniform estimates for moments of strength comparable to that of the original estimate of Granville and Soundararajan for  $\omega(n)$ . With our emphasis on the strength of the method, we have refrained ourselves from pursuing the most general theorems at the risk of complicating our exposition. Despite this compromise, our results are already stronger than the corresponding ones obtained in [12] while retaining some of its own advantages over the results in [33, 34]. Our approach is elementary and flexible, and it can be applied to handle certain arithmetic functions of special interests which were only studied previously by different methods. See the comment below Corollary 2.5.

**Definitions and notation.** We introduce some terminologies and notation that will be adopted throughout this paper without further clarification. Given any real or complex valued functions  $f(x)$  and  $g(x)$  with a common domain  $\mathcal{D} \subseteq \mathbb{R}$ , we shall use Landau’s big- $O$  notation  $f(x) = O(g(x))$  and Vinogradov’s notation  $f(x) \ll g(x)$  interchangeably to mean that there exists an absolute constant  $C > 0$  such that  $|f(x)| \leq C|g(x)|$  for all  $x \in \mathcal{D}$ . Likewise, we shall use the notation  $f(x) \gg g(x)$  interchangeably with  $g(x) = O(f(x))$ . If  $f(x) = O(g(x))$  and  $g(x) = O(f(x))$  hold simultaneously, then we adopt the short-hand notation  $f(x) \asymp g(x)$ . If  $\mathcal{D}$  contains a neighborhood of  $\infty$ , then we write  $f(x) = o(g(x))$  when  $f(x)/g(x) \rightarrow 0$  as  $x \rightarrow \infty$  and  $f(x) \sim g(x)$  when  $f(x)/g(x) \rightarrow 1$  as  $x \rightarrow \infty$ . We shall occasionally make use of the characteristic function  $\epsilon_{a,b}$  of the condition  $a \neq b$  for any  $a, b \in \mathbb{R}$ . Equivalently,  $\epsilon_{a,b} = 1 - \delta_{a,b}$ , where  $\delta_{a,b}$  is the Kronecker delta function.

Throughout, the letter  $p$  always denotes a prime, and we write  $\pi(x)$  for the prime counting function, namely,  $\pi(x) = \sum_{p \leq x} 1$ . For any  $x \in \mathbb{R}$ , we write  $[x]$  for the integer part of  $x$  and  $\lceil x \rceil$  for the least integer  $\geq x$ . For every  $n \in \mathbb{N}$ , denote by  $P^-(n)$  and  $P^+(n)$  the least and the greatest prime factor of  $n$ , respectively, with the convention that  $P^-(1) = \infty$  and  $P^+(1) = 1$ .

We say that  $n \in \mathbb{Z} \setminus \{0\}$  is *squareful* if for any prime  $p \mid n$ , one has  $p^2 \mid n$ . Given any prime power  $p^\nu$ , the relation  $p^\nu \parallel n$  means that  $p^\nu \mid n$  but  $p^{\nu+1} \nmid n$ . In addition, we denote by  $R_n$  the *radical* of  $n$ , i.e.,  $R_n := \text{rad}(n) = \prod_{p \mid n} p$ . Finally, we write  $\binom{m}{m_1, \dots, m_k} := \frac{m!}{m_1! \cdots m_k!}$  for the multinomial coefficient of shape  $(m_1, \dots, m_k)$  of size  $m = m_1 + \cdots + m_k$ .

## 2. MAIN RESULTS

The weights  $\alpha: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$  that we shall consider in this paper form a nice subclass  $\mathcal{M}^*$  of nonnegative multiplicative functions, nice in the sense that there exist constants  $A_0, \beta, \sigma_0 > 0$ ,  $\vartheta_0 \geq 0$ ,  $\varrho_0 \in [0, 1)$  and  $r \in (0, 1)$ , such that the following conditions hold:

$$(i) \quad \alpha(p^\nu) \ll p^{(\varrho_0 + \sigma_0 - 1)\nu}, \quad (2.1)$$

$$(ii) \quad \sum_{p \leq x} \frac{\alpha(p) \log p}{p^{\sigma_0 - 1}} = \beta x + O\left(\frac{x}{(\log x)^{A_0}}\right), \quad (2.2)$$

$$(iii) \quad \sum_p' \left( \frac{\alpha(p)^2}{p^{2(r + \sigma_0 - 1)}} + \sum_{\nu \geq 2} \frac{\alpha(p^\nu)}{p^{(r + \sigma_0 - 1)\nu}} \right) < \infty, \quad (2.3)$$

$$(iv) \quad \sum_{\nu \geq 1} \frac{\nu \alpha(p^\nu)}{p^{\sigma_0 \nu}} \ll \frac{(\log \log(p + 1))^{\vartheta_0}}{p}, \quad (2.4)$$

where the sum  $\sum_p'$  is over all but finitely many primes  $p$ . It is not hard to verify that  $\mathcal{M}^*$  is closed under Dirichlet convolution. Moreover,  $\mathcal{M}^*$  properly contains the subclass of multiplicative functions studied by Elboim and Gorodetsky [12]. For instance, the multiplicative function  $\alpha$ , defined by  $\alpha(p) = 1$  for all primes  $p$  and  $\alpha(p^\nu) = p^{\nu/3}$  for all prime powers  $p^\nu$  with  $\nu \geq 2$ , falls into  $\mathcal{M}^*$  but is not covered by the first part of [12, Theorem 1.1].

The class  $\mathcal{M}^*$  contains many familiar multiplicative functions, including the  $\kappa$ -fold divisor function  $d_\kappa(n)$  for  $\kappa > 0$ , the sum-of-divisors function  $\sigma_\lambda(n)$  for  $\lambda > -1$ , Euler's totient function  $\varphi(n)$ , the characteristic function  $\mu(n)^2$  of square-free numbers, and the function  $r_2(n)/4$ , where  $\mu(n)$  is the Möbius function and  $r_2(n) := \#\{(a, b) \in \mathbb{Z}^2 : n = a^2 + b^2\}$ . Less obvious examples include  $\rho_g(n)$ , which denotes the number of zeros of a nonconstant irreducible polynomial  $g \in \mathbb{Z}[x]$  in  $\mathbb{Z}/n\mathbb{Z}$ , and Ramanujan's  $\tau$ -function  $\tau(n)$ , which may be defined as the  $n$ th Fourier coefficient of the modular discriminant. We leave the verification of these claims to the interested reader.

Let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$  and set

$$S(x) = S_\alpha(x) := \sum_{n \leq x} \alpha(n).$$

For any additive function  $f: \mathbb{N} \rightarrow \mathbb{R}$ , define

$$A(x) = A_{\alpha, f}(x) := \sum_{p \leq x} \alpha(p) \frac{f(p)}{p^{\sigma_0}},$$

$$B(x) = B_{\alpha, f}(x) := \sum_{p \leq x} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}}.$$

One may think of  $n$  as a random variable defined on the sample space  $\mathbb{N} \cap [1, x]$  with the natural probability measure induced by  $\alpha$ , that is to say,  $\text{Prob}(n = k) = \alpha(k)/S(x)$  for every  $k \in \mathbb{N} \cap [1, x]$ . We shall show, by estimating the weighted  $m$ th moment defined by

$$M(x; m) = M_{\alpha, f}(x; m) := S(x)^{-1} \sum_{n \leq x} \alpha(n) (f(n) - A(x))^m$$

for every  $m \in \mathbb{N}$ , that for certain additive functions  $f$ , the limiting distribution of the normalization  $(f(n) - A(x))/\sqrt{B(x)}$  is standard Gaussian. To state our results in a coherent manner, we set  $\chi_m := (1 + (-1)^m)/2$ , the characteristic function of even integers, and

$$C_m := \frac{m!}{2^{m/2} \Gamma(m/2 + 1)}$$

for  $m \in \mathbb{N}$ , where  $\Gamma$  is the Gamma function. One quickly notes that  $C_m = \mu_m = (m-1)!!$  for  $m$  even. The numbers  $C_m$  play a nonnegligible role in the error terms of our uniform estimates for  $M(x; m)$ . Our first result is the following theorem.

**Theorem 2.1.** *Let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be a strongly additive function with  $|f(p)| \leq M$  for all  $p$ , where  $M > 0$  is constant, and let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ . If  $\beta = 1$  and  $0 < h_0 < (3/2)^{2/3}$  is arbitrary, and if  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , then we have*

$$M(x; m) = C_m B(x)^{\frac{m}{2}} \left( \chi_m + O \left( \frac{M m^{\frac{3}{2}}}{\sqrt{B(x)}} \right) \right)$$

*uniformly for all sufficiently large  $x$  and all  $1 \leq m \leq h_0(B(x)/M^2)^{1/3}$ . If  $\beta \neq 1$  and if  $B(x)/(\log \log x)^2 \rightarrow \infty$  as  $x \rightarrow \infty$ , then we have*

$$M(x; m) = C_m B(x)^{\frac{m}{2}} \left( \chi_m + O \left( \frac{M m^{\frac{3}{2}} \log \log \log x}{\sqrt{B(x)}} \right) \right)$$

*uniformly for all sufficiently large  $x$  and all  $1 \leq m \ll B(x)^{1/3}/(\log \log \log x)^{2/3}$ . The implicit constants in the error terms of both asymptotic formulas above depend at most on the explicit and implicit constants in the hypotheses except for  $M$ .*

*Remark 2.1.* It may be worth pointing out that as in Theorem 2.1, the implicit constants in the estimates appearing in the rest of the paper depend at most on the explicit and implicit constants in the hypotheses unless stated otherwise.

In the case where  $\alpha = 1$  and  $f = \omega$ , we recover [18, Theorem 1] with a slightly wider range  $1 \leq m \leq h_0(\log \log x)^{1/3}$  compared to the original range  $1 \leq m \leq (\log \log x)^{1/3}$ . Though Theorem 2.1 is formulated for strongly additive functions, similar things can be said about additive functions whose values at prime powers do not grow too rapidly and are hence not expected to contribute very much. A simple example of such functions is  $\Omega(n)$ . Since  $\Omega(p^\nu) = \nu$  for all  $p^\nu$ , one can show, by establishing a weighted version of (1.2), that  $\Omega(n)$  does not differ from its cousin  $\omega(n)$  very much for most values of  $n$ , and so they are expected to have the same distribution. More generally, we shall prove the following variant of Theorem 2.1 for additive functions. For simplicity's sake, we shall focus on a subclass of the multiplicative functions in  $\mathcal{M}^*$ .

**Theorem 2.2.** *Let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be an additive function such that  $f(p^\nu) \leq M\nu^\kappa$  for all prime powers  $p^\nu$ , where  $M > 0$  and  $\kappa \geq 0$  are constant. Let  $\alpha: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$  be a multiplicative function, and suppose that there exist constants  $A_0, \beta, \sigma_0 > 0$ ,  $\vartheta_0 \geq 0$ ,  $\varrho_0 \in [0, 1/2)$  and  $\lambda \in (0, 2^{1-2\varrho_0})$ , such that  $\alpha(n)$  satisfies (2.2), (2.4), and the condition that  $\alpha(p^\nu) = O((\lambda p^{\varrho_0 + \sigma_0 - 1})^\nu)$  for all prime powers  $p^\nu$ . If  $\beta = 1$  and  $0 < h_0 < (3/2)^{2/3}$  is arbitrary, and if  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , then we have*

$$M(x; m) = C_m B(x)^{\frac{m}{2}} \left( \chi_m + O \left( \frac{M m^{\kappa + \frac{3}{2}}}{\sqrt{B(x)}} \right) \right)$$

*uniformly for all sufficiently large  $x$  and all  $m \in \mathbb{N}$  satisfying  $m \leq h_0(B(x)/M^2)^{1/3}$  and  $m \ll B(x)^{1/(2\kappa+3)}$ . If  $\beta \neq 1$  and if  $B(x)/(\log \log \log x)^2 \rightarrow \infty$  as  $x \rightarrow \infty$ , then we have*

$$M(x; m) = C_m B(x)^{\frac{m}{2}} \left( \chi_m + O \left( \frac{M m^{\frac{3}{2}} (\log \log \log x + m^\kappa)}{\sqrt{B(x)}} \right) \right)$$

*uniformly for all sufficiently large  $x$  and all*

$$1 \leq m \ll \min \left( B(x)^{1/(2\kappa+3)}, \frac{B(x)^{1/3}}{(\log \log \log x)^{2/3}} \right).$$

*The implicit constants in the error terms of both asymptotic formulas above depend at most on the explicit and implicit constants in the hypotheses except for  $M$ .*

Theorem 2.2 clearly implies the first part of [12, Theorem 1.1] if we set  $f = \Omega$ ,  $\kappa = 1$ , and  $\vartheta_0 = \varrho_0 = 0$ . It is easy to see that if  $\alpha(p^\nu) = O((\lambda p^{r_0 + \sigma_0 - 1})^\nu)$  for all prime powers  $p^\nu$ , where  $\sigma_0 > 0$ ,  $r_0 \in [0, 1/2)$  and  $\lambda \in (0, 2^{1-2r_0})$  are given constants, then conditions (i) and (iii) are automatically fulfilled with any fixed  $\max(r_0 + \log_2 \lambda, 0) \leq \varrho_0 < 1$ ,  $r_0 + \max(1/2, \log_2 \lambda) < r < 1$ , and of course the same parameter  $\sigma_0$ . Indeed, we shall derive Theorem 2.2 as a corollary of Theorem 2.1.

Let  $g \in \mathbb{Z}[x]$  be a nonconstant irreducible polynomial, and recall that for every  $n \in \mathbb{N}$ ,  $\rho_g(n)$  denotes the number of zeros of  $g$  in  $\mathbb{Z}/n\mathbb{Z}$ . More generally, if  $g \in \mathbb{Q}[x]$  is a nonconstant irreducible polynomial, we may extend the definition above by setting  $\rho_g(n) = 0$  if  $\gcd(n, c_g) > 1$ , where  $c_g \in \mathbb{N}$  is the least positive integer such that  $c_g g(x) \in \mathbb{Z}[x]$ , and insisting that  $\rho_g(n)$  be the number of zeros of  $g(x)$  (or equivalently,  $c_g g(x)$ ) in  $\mathbb{Z}/n\mathbb{Z}$  when  $\gcd(n, c_g) = 1$ . Extended this way with the convention that  $\rho_g(1) = 1$ , the function  $\rho_g(n)$  remains multiplicative. By [21, Lemma 1],  $\rho_g$  is bounded on prime powers and satisfies

$$\sum_{p \leq x} \frac{\rho_g(p)}{p} = \log \log x + M_{\rho_g} + O \left( \frac{1}{\log x} \right).$$

Given a strongly additive function  $f: \mathbb{N} \rightarrow \mathbb{R}$ , we define

$$A_{f,g}(x) := \sum_{p \leq x} \rho_g(p) \frac{f(p)}{p},$$

$$B_{f,g}(x) := \sum_{p \leq x} \rho_g(p) \frac{f(p)^2}{p}.$$

For simplicity's sake, suppose that  $g(\mathbb{N}) \subseteq \mathbb{N}$ . In the case  $g \in \mathbb{Z}[x]$ , Halberstam [20, Theorem 3] showed that if  $B_{f,g}(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , and if  $f(p) = o(\sqrt{B_{f,g}(p)})^1$ , then given  $m \in \mathbb{N}$ ,

$$\frac{1}{x} \sum_{n \leq x} (f(g(n)) - A_{f,g}(x))^m = (\mu_m + o(1)) B_{f,g}(x)^{\frac{m}{2}}$$

holds. Under the stronger condition  $f(p) = O(1)$ , Theorem 2.1 leads to a weighted version of this result in the case  $g(n) = n$ . The remaining cases are captured by the following theorem.

**Theorem 2.3.** *Let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be a strongly additive function with  $|f(p)| \leq M$  for all  $p$ , where  $M > 0$  is constant, and let  $g \in \mathbb{Q}[x]$  be a nonconstant irreducible polynomial such that  $g(0) \neq 0$  and  $g(\mathbb{N}) \subseteq \mathbb{N}$ . Let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ , and fix  $0 < h_0 < (3/2)^{2/3}$  and  $h'_0 > 0$ . For any  $q \in \mathbb{N}$  and  $a \in \mathbb{Z}$  coprime to  $q$ , define*

$$\Delta_\alpha(x; q, a) := \sum_{\substack{n \leq x \\ n \equiv a \pmod{q}}} \alpha(n) - \frac{1}{\varphi(q)} \sum_{\substack{n \leq x \\ \gcd(n, q) = 1}} \alpha(n),$$

If there exist a constant  $\epsilon_0 > 0$  and a function  $\delta(x) \in (0, 1]$  with  $\delta(x)^2 B_{f,g}(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , such that

$$\sum_{\substack{q: \omega(q) \leq m \\ P^+(q) \leq x^{\delta(x)/m}}} \mu(q)^2 \sum_{a \in \mathcal{Z}_g^*(q)} |\Delta_\alpha(x; q, a)| \ll S(x) \exp(-(\log \log x)^{1/3+\epsilon_0}) \quad (2.5)$$

uniformly for all sufficiently large  $x$  and all

$$1 \leq m \leq \min(h_0(B_{f,g}(x)/M^2)^{1/3}, h'_0(\delta(x)^2 B_{f,g}(x)/M^2)^{1/3}), \quad (2.6)$$

where  $\mathcal{Z}_g^*(q)$  is the zero locus of  $g$  in  $(\mathbb{Z}/q\mathbb{Z})^\times$ . Then

$$S(x)^{-1} \sum_{n \leq x} \alpha(n) (f(g(n)) - A_{f,g}(x))^m = C_m B_{f,g}(x)^{\frac{m}{2}} \left( \chi_m + O\left(\frac{M m^{\frac{3}{2}}}{\delta(x) \sqrt{B_{f,g}(x)}}\right) \right)$$

uniformly for all sufficiently large  $x$  and all  $m \in \mathbb{N}$  in the same range (2.6), where the implicit constant in the error term depends at most on the explicit and implicit constants in the hypotheses except for  $M$ .

Theorem 2.3 is applicable to a large class of nonnegative multiplication functions  $\alpha(n)$ , including  $d_k(n)$  for  $k \in \mathbb{N}$  and  $r_2(n)/4$  [5, 27]. Despite the great generality of (2.5), it is oftentimes more convenient to work with the stronger variant

$$\sum_{\substack{q \leq x^{\delta(x)} \\ \omega(q) \ll (\delta(x)^2 \log \log x)^{1/3}}} \mu(q)^2 \rho_g(q) \max_{a \in (\mathbb{Z}/q\mathbb{Z})^\times} |\Delta_\alpha(x; q, a)| \ll S(x) \exp(-(\log \log x)^{1/3+\epsilon_0}) \quad (2.7)$$

<sup>1</sup>Halberstam [20] wrote that for  $g(x) = x$  this pair of conditions contain the condition that  $f(p) = o((\log p)^\epsilon)$  for every given  $\epsilon > 0$ . However, this claim is incorrect. In fact, a simple counterexample may be constructed as follows. Let  $\mathcal{P}$  be an arbitrary infinite subset of odd primes such that  $\sum_{p \in \mathcal{P}} 1/p < \infty$ , and put  $\mathcal{P}(x) := \mathcal{P} \cap [3, x]$ . Define  $f(p) = \sqrt{\log \log p}$  for  $p \in \mathcal{P}$  and  $f(p) = 1$  for  $p \notin \mathcal{P}$ . From partial summation it follows that  $\sum_{p \in \mathcal{P}(x)} f(p)^2/p = o(\log \log x)$ . Then one sees readily that  $f(p) = o((\log p)^\epsilon)$  for any given  $\epsilon > 0$ , while  $f(p) \sim \sqrt{B(p)}$  for large  $p \in \mathcal{P}$ .

This condition may be viewed as an inequality of the *Bombieri–Vinogradov type*, which ensures that the values of  $\alpha(n)$  are well distributed as  $n$  varies over the reduced residue classes  $a \pmod{q}$  for most values of  $q$  and  $a$ . In view of [18, Proposition 4], such a condition arises from a sieving process for the sequence  $\{g(n)\}_{n \geq 1}$ . For this process to work, we need information about the average size of  $\alpha(n)$  subject to the constraint  $d \mid g(n)$  for smooth square-free  $d \in \mathbb{N} \cap [1, x^\delta]$ . If  $d$  is also free of small prime factors up to some constant depending on  $g$ , then this constraint amounts to the congruences  $n \equiv a \pmod{d}$  for  $a \in \mathcal{Z}_g^*(d)$ . So, (2.7) reduces the sieving of  $\{g(n)\}_{n \geq 1}$  to that of  $\mathbb{N}$ .

We shall only sketch the proof of Theorem 2.3, since it is similar to, and in fact, much easier than that of Theorem 2.1. The argument used in the proof may also be modified to study the joint distribution of  $f(n + h_1)$  and  $f(n + h_2)$  with any fixed integers  $h_1 \neq h_2$ .

It is not hard to see that the condition  $f(p) = O(1)$  in Theorem 2.1 can be relaxed, especially when we do not pursue uniformity in  $m$  in the asymptotics for the  $m$ th moment. For instance, in the case  $\alpha = 1$  Delange and Halberstam showed [9, Theorem 1] that if  $f: \mathbb{N} \rightarrow \mathbb{R}$  is a strongly additive function such that  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ ,  $f(p) = O(\sqrt{B(p)})$  for all primes  $p$ , and

$$\sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B(x)}}} \frac{f(p)^2}{p} = o(B(x)) \quad (2.8)$$

for any given  $\epsilon > 0$ , then

$$\frac{1}{x} \sum_{n \leq x} (f(n) - A(x))^m = (\mu_m + o(1)) B(x)^{\frac{m}{2}}$$

for every fixed  $m \in \mathbb{N}$ . The implication of this result on the distribution of  $f$  is slightly weaker than the Kubilius–Shapiro theorem [30, Theorem A] in that the latter asserts that the distribution of an additive function  $f: \mathbb{N} \rightarrow \mathbb{R}$  with an unbounded variance  $B(x)$  which satisfies (2.8) for every given  $\epsilon > 0$  is necessarily Gaussian with mean  $A(x)$  and variance  $B(x)$ . On the other hand, Delange and Halberstam noted that their result no longer holds if one removes the assumption  $f(p) = O(\sqrt{B(p)})$ , which incidentally exposes the limitation of the method of moments compared to the method evolved by Erdős and Kac. Regardless, it will be clear in the sequel that the proof of Theorem 2.1 makes it possible for us to obtain the following natural extension of the result of Delange and Halberstam.

**Theorem 2.4.** *Let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be a strongly additive function, and let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ . Define*

$$B^*(x) := \begin{cases} B(x), & \text{if } \beta = 1, \\ B(x)/(\log \log \log x)^2, & \text{if } \beta \neq 1, \end{cases}$$

*and suppose  $B^*(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . If there exists a constant  $K > 0$  such that  $f(n) = o(\sqrt{B^*(x)})$  for all squarefree  $n \in \mathbb{N} \cap [1, x]$  composed of prime factors  $p$  with  $|f(p)| > K\sqrt{B^*(x)}$ , and if*

$$\sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B^*(x)}}} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} = o(B^*(x))$$

*for any given  $\epsilon > 0$ , then  $M(x; m) = (\mu_m + o(1)) B(x)^{\frac{m}{2}}$  for every fixed  $m \in \mathbb{N}$ .*



The proof of Theorem 2.4, which we shall only outline, is based on the proofs of Theorem 2.1 and [9, Theorem 1]. We shall also obtain as a corollary the following analogue of the Kubilius–Shapiro theorem [30, Theorem C].

**Corollary 2.5.** *Under the notation and hypotheses in Theorem 2.4, we have*

$$\lim_{x \rightarrow \infty} S(x)^{-1} \sum_{\substack{n \leq x \\ f(n) \leq A(x) + V\sqrt{B(x)}}} \alpha(n) = \Phi(V)$$

for any given  $V \in \mathbb{R}$ . The same is true if  $f$  is merely additive.

It is clear that Theorem 2.4 implies Corollary 2.5 when  $f$  is strongly additive. To handle the general case where  $f$  is merely additive, we shall establish a weighted version of [30, Theorem B] which shows that when it comes to the distribution problem, there is no essential difference between strongly additive functions and general additive functions, and thus the distribution of an additive function  $f$  is determined solely by its values at primes.

Corollary 2.5 has many interesting applications. For instance, it implies at once that if  $h: \mathbb{N} \rightarrow \mathbb{R}$  is any *completely additive* function, i.e.,  $h(mn) = h(m) + h(n)$  for all  $m, n \in \mathbb{N}$ , and if  $k > 1$  is a positive integer, then the distribution of  $h(d_k(n))$  weighted by  $\alpha(n)$  is Gaussian with mean  $h(k)\beta \log \log x$  and variance  $h(k)^2\beta \log \log x$ , provided  $h(k) \neq 0$ . In [13] Elliott proved a weighted Erdős–Kac theorem concerning Ramanujan’s  $\tau$ -function. In Remark 9.1 we describe how his result may be derived from Corollary 2.5. Analogues on elliptic holomorphic newforms of weight at least 2 can be obtained in the same way. In a similar fashion, one can also show that if the weight  $\alpha$  in Corollary 2.5 satisfies the additional condition that  $\alpha(p) \sim \beta p^{\sigma_0-1}$  for all but a subset  $E$  of primes  $p$ , where  $\#(E \cap [2, x]) = o(x(\log \log x)^{2-\vartheta_0}/(\log x)^3)$  as  $x \rightarrow \infty$ , then the distribution of  $\Omega(\varphi(n))$  weighted by  $\alpha(n)$  is Gaussian with mean  $\beta(\log \log x)^2/2$  and variance  $\beta(\log \log x)^3/3$ , generalizing an old result of Erdős and Pomerance [17, Theorem 3.1] in an easy manner.

*Remark 2.2.* The condition that  $f(p) = o((\log p)^\epsilon)$  for any given  $\epsilon > 0$ , mentioned by Halberstam [20], does not imply (2.8) in general. To see this, assume for the moment that there exists an infinite subset  $\mathcal{P}$  of primes such that

$$s_{\mathcal{P}}(x) := \sum_{p \in \mathcal{P} \cap [17, x]} \frac{1}{p} = \frac{\log \log x}{\log \log \log x} + c + o(1) \quad (2.9)$$

for sufficiently large  $x$ , where  $c \in \mathbb{R}$  is some constant. Define  $f(p) = (\log p)^{1/(2 \log \log \log p)}$  for  $p \in \mathcal{P}$  and  $f(p) = 1$  for  $p \notin \mathcal{P}$ . Clearly,  $f(p) = o((\log p)^\epsilon)$  for any given  $\epsilon > 0$ . It is easily seen by partial summation that

$$\sum_{p \in \mathcal{P} \cap [17, x]} \frac{f(p)^2}{p} = \int_{17^-}^x (\log t)^{1/\log \log \log t} ds_{\mathcal{P}}(t) = (1 + o(1))(\log x)^{1/\log \log \log x},$$

which implies that

$$B(x) = \sum_{p \in \mathcal{P} \cap [17, x]} \frac{f(p)^2}{p} + O(\log \log x) = (1 + o(1))(\log x)^{1/\log \log \log x}.$$

Let  $y = x^{1/\log \log x}$  and  $\epsilon = 1/2$ . Since

$$\begin{aligned}\log \log y &= \log \log x - \log \log \log x, \\ \log \log \log y &= \left(1 + O\left(\frac{1}{\log \log x}\right)\right) \log \log \log x,\end{aligned}$$

we have

$$(\log y)^{1/\log \log \log y} = \exp\left(\frac{\log \log x}{\log \log \log x} - 1 + O\left(\frac{1}{\log \log \log x}\right)\right).$$

It follows that

$$f(p)^2 > (\log y)^{1/\log \log \log y} > \frac{1}{3}(\log x)^{1/\log \log \log x} > \epsilon^2 B(x)$$

for  $p \in \mathcal{P} \cap (y, x]$  when  $x$  is sufficiently large. Hence, we have

$$\sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B(x)}}} \frac{f(p)^2}{p} \geq \sum_{p \in \mathcal{P} \cap (y, x]} \frac{f(p)^2}{p} > \frac{1}{2}(\log x)^{1/\log \log \log x} > \frac{1}{3}B(x).$$

It remains to construct a set  $\mathcal{P}$  with the desired property (2.9). The following inductive approach was suggested by Prof. Pomerance. Note first that  $\sum_{p \leq x} 1/p = \log \log x + O(1)$  grows slightly faster than our target  $u(x) := \log \log x / \log \log \log x$ , according to Mertens' second theorem [23, Theorem 427]. Moreover, if  $p < p'$  are large consecutive primes, then  $u(p') - u(p) = o(1/\log p)$ , by Bertrand's postulate. Let 17 be the first prime in  $\mathcal{P}$ . Suppose that we have already selected for  $\mathcal{P}$  the primes up to  $q$ , where  $q$  is prime. We put the next prime  $q'$  in  $\mathcal{P}$  if  $s_{\mathcal{P}}(q) < u(q)$  and leave it out of  $\mathcal{P}$  otherwise. Then the running sum  $s_{\mathcal{P}}(x)$  changes by at most  $1/q$  as  $x$  moves from  $q$  to  $q'$ , while the target  $u(x)$  changes by at most  $o(1/\log q)$  as  $x$  moves from  $q$  to  $q'$ . Thus, the difference  $s_{\mathcal{P}}(x) - u(x)$  can be kept within  $o(1/\log x)$ . In particular, (2.9) holds for  $\mathcal{P}$  with  $c = 0$ .

**Overview of the proof of Theorem 2.1.** Before embarking on the proofs of our results, we describe briefly the main steps in the proof of Theorem 2.1. The starting point is an approximation to moments used by Granville, Soundararajan, Khan, Milinovich and Subedi. Though the underlying idea is the same, we need a more complicated version of this approximation (see Lemma 4.1) due to the more general nature of our multiplicative weights  $\alpha$ . To utilize it, we first need to develop an asymptotic formula for the mean value of  $\alpha(n)$  with  $n \leq x$  restricted to squarefree integers  $a \in \mathbb{N} \cap [1, x]$  (see Lemma 3.3). An important feature of this formula is that it holds uniformly for all squarefree integer  $a \in \mathbb{N} \cap [1, x]$ , which is key to both applying the moment approximation and making the moment estimates uniform. This formula will serve as a substitute for the one on  $d_k$  developed by Khan, Milinovich and Subedi. Unlike their proof, which is based on Peron's formula and the complete submultiplicativity of  $d_k$ , our proof uses the mean value estimate for  $\alpha$  supplied by [6, Theorem 2.1] and is completely elementary. This is carried out in the next section.

After applying the moment approximation, we find that the estimation of the main contribution can be worked out as in [18] and [25]. It is the estimation of the error terms that is more involved in our case. In particular, the estimation of the error term in the moment approximation provided by Lemma 4.1 in Section 4 requires separate treatments according as  $\beta = 1$  or  $\beta \neq 1$ . Besides, since the error term in our asymptotic formula for the mean

value of  $\alpha(n)$  over  $a \mid n$  supplied by Lemma 3.3 in Section 3 is weaker than what one can obtain for the special weight  $d_k(n)$  by complex analytic approaches, we need to handle the case  $\beta \in (0, 1)$  with some special care and make a careful selection of parameters accordingly in order to minimize the error terms. With these new technicalities taken care of, we obtain the desired uniform estimates for moments stated in Theorem 2.1.

### 3. MEAN VALUES OF MULTIPLICATIVE FUNCTIONS

Without loss of generality, we may assume  $A_0 \in (0, 1)$  in the sequel. In addition, we shall also make use of the asymptotic formula

$$\sum_{p \leq x} \frac{\alpha(p)}{p^{\sigma_0}} = \beta \log \log x + M_\alpha + O((\log x)^{-A_0}) \quad (3.1)$$

with some constant  $M_\alpha \in \mathbb{R}$ , which follows immediately from (2.2) via partial summation. In view of our assumption that  $f(p) = O(1)$ , this formula implies trivially that  $B(x) \ll \log \log x$ . Moreover, if we define, for every prime  $p$ ,

$$\psi_0(p) := \sum_{\nu \geq 2} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}},$$

then we infer from (2.1), (2.3) and (2.4) that

$$\psi_0(p) \ll \frac{(\log \log(p+1))^{\vartheta_0}}{p}$$

and that  $\sum_p \psi_0(p) < \infty$ .

**Lemma 3.1.** *Let  $\alpha: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$  be a multiplicative function satisfying (2.1) and (2.4) with some  $\sigma_0, \vartheta_0 > 0$  and  $\varrho_0 \in [0, 1)$ . Fix  $h \in \mathbb{R}$ ,  $\epsilon_0 \in (0, 1)$  and  $c_0 \in [1, \epsilon_0^{-1})$ , and define*

$$I_{\alpha, h}(x; a) := \sum_{\substack{q \leq x \\ R_q = a}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3x}{q} \right)^h,$$

where  $a \in \mathbb{N} \cap [1, x]$  is squarefree. Then there exists a constant  $\delta_0 > 0$  such that uniformly for all sufficiently large  $x$ , any  $\delta \in [\delta_0 \log \log x / \log x, 1]$ , and any squarefree  $a \in \mathbb{N} \cap [1, x]$  with  $\omega(a) \leq (1 - \varrho_0) \epsilon_0 \delta^{-1}$ , we have

$$I_{\alpha, h}(x; a) = \left( \tilde{\lambda}_\alpha(a) + O \left( \frac{2^{O(\omega(a))}}{\log x} \left( \frac{1}{x^{c_0 \delta \omega(a)}} + \frac{\epsilon_{h,0} L(a) \log P^+(a)}{a} \right) \right) \right) (\log x)^h,$$

where

$$\tilde{\lambda}_\alpha(a) := \prod_{p|a} \sum_{\nu=1}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}},$$

$$L(a) := \prod_{p|a} (\log \log(p+1))^{\vartheta_0}.$$

*Proof.* Let  $\delta \in (0, 1]$  and fix  $c_1 \in (c_0, \epsilon_0^{-1})$ . Put  $\delta_1 := (1 - \varrho_0)^{-1} c_1 \delta$  and  $y := x^{k\delta_1}$ . For any squarefree  $a = p_1 \cdots p_k \in \mathbb{N} \cap [1, x]$  with  $p_1 < \cdots < p_k \leq x$  and  $k \leq (1 - \varrho_0) \epsilon_0 \delta^{-1}$ , we have  $k\delta_1 \leq c_1 \epsilon_0 < 1$  and

$$I_{\alpha, h}(x; a) = \sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} \leq x \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_k^{\nu_k}} \right)^h.$$

On the one hand, we see that

$$\begin{aligned} & \sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_k^{\nu_k}} \right)^h \\ &= \sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} (\log 3x)^h \left( 1 + O \left( \frac{\epsilon_{h,0}}{\log 3x} \sum_{i=1}^k \nu_i \log p_i \right) \right) \\ &= \sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} (\log x)^h + O \left( \frac{2^{O(k)} \epsilon_{h,0} L(a) \log p_k (\log x)^{h-1}}{a} \right), \end{aligned}$$

by (2.4). From (2.1) it follows that

$$\sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} = \tilde{\lambda}_\alpha(a) + O \left( 2^{O(k)} \sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} > y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{1}{p_1^{(1-\varrho_0)\nu_1} \cdots p_k^{(1-\varrho_0)\nu_k}} \right).$$

The sum in the error term above may be split into two sums according as  $p_2^{\nu_2} \cdots p_k^{\nu_k} \leq y$  or  $p_2^{\nu_2} \cdots p_k^{\nu_k} > y$ . In the first sum we must have  $p_1^{\nu_1} > y/(p_2^{\nu_2} \cdots p_k^{\nu_k})$ . Thus summing over  $\nu_1$  and then over  $\nu_2, \dots, \nu_k$ , we see that the first sum is

$$\ll \frac{1}{y^{1-\varrho_0}} \sum_{\substack{p_2^{\nu_2} \cdots p_k^{\nu_k} \leq y \\ \nu_2, \dots, \nu_k \geq 1}} 1 \leq \frac{2^{O(k)} (\log x)^{k-1}}{x^{c_1 k \delta} (\log p_2) \cdots (\log p_k)}.$$

The second sum is simply

$$\sum_{\substack{p_2^{\nu_2} \cdots p_k^{\nu_k} > y \\ \nu_2, \dots, \nu_k \geq 1}} \frac{1}{p_2^{(1-\varrho_0)\nu_2} \cdots p_k^{(1-\varrho_0)\nu_k}} \sum_{\nu_1 \geq 1} \frac{1}{p_1^{(1-\varrho_0)\nu_1}} \ll \sum_{\substack{p_2^{\nu_2} \cdots p_k^{\nu_k} > y \\ \nu_2, \dots, \nu_k \geq 1}} \frac{1}{p_2^{(1-\varrho_0)\nu_2} \cdots p_k^{(1-\varrho_0)\nu_k}}.$$

It follows that

$$\sum_{\substack{p_1^{\nu_1} \cdots p_k^{\nu_k} > y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{1}{p_1^{(1-\varrho_0)\nu_1} \cdots p_k^{(1-\varrho_0)\nu_k}} \ll \sum_{\substack{p_2^{\nu_2} \cdots p_k^{\nu_k} > y \\ \nu_2, \dots, \nu_k \geq 1}} \frac{1}{p_2^{(1-\varrho_0)\nu_2} \cdots p_k^{(1-\varrho_0)\nu_k}} + \frac{2^{O(k)} (\log x)^{k-1}}{x^{c_1 k \delta} (\log p_2) \cdots (\log p_k)}.$$

Repeating this argument, we obtain

$$\sum_{\substack{p_1^{\nu_1} \dots p_k^{\nu_k} > y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{1}{p_1^{(1-\varrho_0)\nu_1} \dots p_k^{(1-\varrho_0)\nu_k}} \leq \frac{2^{O(k)} (\log x)^{k-1}}{x^{c_1 k \delta} (\log p_2) \dots (\log p_k)},$$

from which we deduce

$$\sum_{\substack{p_1^{\nu_1} \dots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \dots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \dots p_k^{\sigma_0 \nu_k}} = \tilde{\lambda}_\alpha(a) + O\left(\frac{2^{O(k)} (\log x)^{k-1}}{x^{c_1 k \delta} (\log p_2) \dots (\log p_k)}\right). \quad (3.2)$$

On the other hand, we have

$$\begin{aligned} \sum_{x_1 < p^\nu \leq x_2} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} \left( \log \frac{3x_2}{p^\nu} \right)^h &\ll \sum_{\substack{\log_p x_1 < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)\nu}} \left( \log \frac{3x_2}{p^\nu} \right)^h \\ &= - \int_{\log_p x_1}^{\log_p x_2} \left( \log \frac{3x_2}{p^t} \right)^h d \left( \sum_{\substack{t < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)t}} \right) \end{aligned}$$

uniformly for all primes  $p$  and all  $0 < x_1 \leq x_2$ . Using integration by parts, we see that the integral above is equal to

$$-(\log(3x_2/x_1))^h \sum_{\substack{\log_p x_1 < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)\nu}} - \int_{\log_p x_1}^{\log_p x_2} \left( \sum_{\substack{t < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)t}} \right) d \left( \log \frac{3x_2}{p^t} \right)^h.$$

Since

$$\sum_{\substack{t < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)t}} < \frac{1}{p^{(1-\varrho_0)(\lfloor t \rfloor + 1)}} \cdot \frac{p^{1-\varrho_0}}{p^{1-\varrho_0} - 1} \ll \frac{1}{p^{(1-\varrho_0)t}},$$

we have

$$(\log(3x_2/x_1))^h \sum_{\substack{\log_p x_1 < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)\nu}} \ll \frac{(\log(3x_2/x_1))^h}{x_1^{1-\varrho_0}}$$

and

$$\begin{aligned}
\int_{\log_p x_1}^{\log_p x_2} \left( \sum_{\substack{t < \nu \leq \log_p x_2 \\ \nu \in \mathbb{Z}}} \frac{1}{p^{(1-\varrho_0)t}} \right) d \left( \log \frac{3x_2}{p^t} \right)^h &\ll \epsilon_{h,0} \log p \int_{\log_p x_1}^{\log_p x_2} \frac{1}{p^{(1-\varrho_0)t}} \left( \log \frac{3x_2}{p^t} \right)^{h-1} dt \\
&= \frac{\epsilon_{h,0}}{(3x_2)^{1-\varrho_0}} \int_{\log 3}^{\log(3x_2/x_1)} t^{h-1} e^{(1-\varrho_0)t} dt \\
&\ll \frac{\epsilon_{h,0}}{(3x_2)^{1-\varrho_0}} (\log(3x_2/x_1))^{h-1} \left( \frac{3x_2}{x_1} \right)^{1-\varrho_0} \\
&= \frac{\epsilon_{h,0} (\log(3x_2/x_1))^{h-1}}{x_1^{1-\varrho_0}}.
\end{aligned}$$

Hence, it follows that

$$\sum_{x_1 < p^\nu \leq x_2} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} \left( \log \frac{3x_2}{p^\nu} \right)^h \ll \frac{(\log(3x_2/x_1))^h}{x_1^{1-\varrho_0}} \quad (3.3)$$

uniformly for all primes  $p$  and all  $0 < x_1 \leq x_2$ . This inequality implies immediately

$$\begin{aligned}
\sum_{\substack{y < p_1^{\nu_1} \cdots p_k^{\nu_k} \leq x \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_k^{\nu_k}} \right)^h &\leq \frac{2^{O(k)} (\log x)^h}{y^{1-\varrho_0}} \sum_{\substack{p_2^{\nu_2} \cdots p_k^{\nu_k} \leq x \\ \nu_2, \dots, \nu_k \geq 1}} 1 \\
&\leq \frac{2^{O(k)} (\log x)^{k+h-1}}{x^{c_1 k \delta} (\log p_2) \cdots (\log p_k)}.
\end{aligned}$$

Lemma 3.1 now follows upon combining the above with (3.2) and taking  $\delta_0 = 1/(c_1 - c_0)$  with the range  $\delta \geq \delta_0 \log \log x / \log x$  in mind.  $\square$

Let  $\alpha$  be a multiplicative function as in Theorem 2.1 with  $A_0 \in (0, 1)$ . Suppose first that (2.3) holds with the restricted sum  $\sum_p'$  replaced by the full sum  $\sum_p$ . For  $\sigma_0 = 1$  De la Bretèche and Tenenbaum [6, Theorem 2.1] showed

$$\sum_{n \leq x} \alpha(n) = \frac{1}{\Gamma(\beta)} \prod_p \left( 1 - \frac{1}{p} \right)^\beta \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^\nu} x (\log x)^{\beta-1} \left( 1 + O \left( \frac{1}{(\log x)^{A_0}} \right) \right),$$

where the implicit constant in the error term depends at most on the explicit and implicit constants in the hypotheses. For the general case where  $\sigma_0 > 0$  is arbitrary, it is easy to show, by applying the above to  $\alpha(n)/n^{\sigma_0-1}$  and employing partial summation as in the proof of [12, Corollary 3.3], that

$$S(x) = \sum_{n \leq x} \alpha(n) = \lambda_\alpha x^{\sigma_0} (\log x)^{\beta-1} \left( 1 + O \left( \frac{1}{(\log x)^{A_0}} \right) \right), \quad (3.4)$$

where

$$\lambda_\alpha := \frac{1}{\sigma_0 \Gamma(\beta)} \prod_p \left( 1 - \frac{1}{p} \right)^\beta \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}}. \quad (3.5)$$

Suppose now that (2.3) holds with the restricted sum  $\sum_p'$  being the sum  $\sum_{p>Q_0}$ , where  $Q_0 \geq 1$  is some constant. Let  $P_0 := \prod_{p \leq Q_0} p$  and  $\mathbf{1}_{P_0}(n)$  the indicator function of the set  $\{n \in \mathbb{N} : \gcd(n, P_0) = 1\}$ . Then  $\alpha(n)\mathbf{1}_{P_0}(n)$  is a nonnegative multiplicative function satisfying (2.1)–(2.4) with the sum  $\sum_p'$  in (2.3) replaced by the full sum  $\sum_p$ . In particular, (3.4) is applicable to  $\alpha(n)\mathbf{1}_{P_0}(n)$ . Thus, we obtain

$$\sum_{\substack{n \leq x \\ \gcd(n, P_0)=1}} \alpha(n) = \lambda_\alpha(P_0) x^{\sigma_0} (\log 3x)^{\beta-1} \left( 1 + O\left( \frac{1}{(\log 3x)^{A_0}} \right) \right), \quad (3.6)$$

where

$$\lambda_\alpha(P_0) := \frac{1}{\sigma_0 \Gamma(\beta)} \prod_{p \leq Q_0} \left( 1 - \frac{1}{p} \right)^\beta \cdot \prod_{p > Q_0} \left( 1 - \frac{1}{p} \right)^\beta \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}}.$$

Examining the proof of Lemma 3.1, we find that for every given  $h \in \mathbb{R}$ ,

$$\sum_{\substack{q \leq x \\ R_q | P_0}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3x}{q} \right)^h = \prod_{p \leq Q_0} \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} (\log x)^h \left( 1 + O\left( \frac{1}{\log x} \right) \right)$$

for all sufficiently large  $x$ . Combining this with (3.6) gives

$$S(x) = \sum_{\substack{q \leq x \\ R_q | P_0}} \alpha(q) \sum_{\substack{n' \leq x/q \\ \gcd(n', P_0)=1}} \alpha(n') = \lambda_\alpha x^{\sigma_0} (\log x)^{\beta-1} \left( 1 + O\left( \frac{1}{(\log x)^{A_0}} \right) \right),$$

which is the same as (3.4).

For our applications, we will need an asymptotic formula for

$$S(x; a) = S_\alpha(x; a) := \sum_{\substack{n \leq x \\ \gcd(n, a)=1}} \alpha(n)$$

uniform in  $a \in \mathbb{N} \cap [1, x]$ . One may be tempted to apply (3.4) to the function  $\alpha(n)\mathbf{1}_a(n)$ , where  $\mathbf{1}_a(n)$  is the indicator function of the set  $\{n \in \mathbb{N} : \gcd(n, a) = 1\}$ . However, it is not immediately clear whether the implied constant in the error term obtained via this naive approach is independent of  $a \in \mathbb{N} \cap [1, x]$ . Fortunately, the following lemma provides the desired estimate for  $S(x; a)$  under the hypotheses (i)–(iv).

**Lemma 3.2.** *For any  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ , we have*

$$S(x; a) = \lambda_\alpha(a) x^{\sigma_0} (\log x)^{\beta-1} \left( 1 + O\left( \frac{1}{(\log x)^{A_0}} \right) \right)$$

*uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ , where*

$$\lambda_\alpha(a) := \frac{1}{\sigma_0 \Gamma(\beta)} \prod_{p|a} \left( 1 - \frac{1}{p} \right)^\beta \cdot \prod_{p \nmid a} \left( 1 - \frac{1}{p} \right)^\beta \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}},$$

*The implicit constant in the error term depends at most on the explicit and implicit constants in the hypotheses.*

*Proof.* Let  $a \in \mathbb{N} \cap [1, x]$ . For simplicity of notation, we write  $\sum^a$  for sums in which the indices take values coprime to  $a$ . As we have demonstrated above, there is no loss of generality by assuming that  $\sigma_0 = 1$  and that (2.3) holds with the restricted sum  $\sum_p'$  replaced by the full sum  $\sum_p$ . We start by determining the relation between  $\lambda_\alpha(a)$  and  $\lambda_\alpha$ . Note that condition (iv) implies that  $\alpha(p) \ll (\log \log(p+1))^{\vartheta_0}$ , from which it follows that

$$\sum_{p|a} \alpha(p) \log p \ll (\log \log x)^{\vartheta_0} \log a \leq (\log \log x)^{\vartheta_0} \log x.$$

By (2.2) we have

$$\sum_{p \leq x}^a \alpha(p) \log p = \beta x + O\left(\frac{x}{(\log x)^{A_0}}\right),$$

from which we deduce

$$\sum_{p \leq x}^a \frac{\alpha(p)}{p} = \beta \log \log x + M_\alpha + O((\log x)^{-A_0}).$$

Combining the above with (3.1), we see that

$$\begin{aligned} \lambda_\alpha(a) &= \lambda_\alpha \prod_{p|a} \left( \sum_{\nu=0}^{\infty} \frac{\alpha(p^\nu)}{p^\nu} \right)^{-1} = \lambda_\alpha \exp \left( - \sum_{p|a} \frac{\alpha(p)}{p} + O(1) \right) \\ &= \lambda_\alpha \exp \left( - \sum_{p \leq x} \frac{\alpha(p)}{p} + \sum_{p \leq x}^a \frac{\alpha(p)}{p} + O(1) \right) \\ &\asymp \lambda_\alpha \asymp 1. \end{aligned}$$

This assures us that there is no need to differentiate  $\lambda_\alpha(a)$  and  $\lambda_\alpha$  in the error terms.

Next, we connect  $S(x; a)$  with

$$T(x; a) = T_\alpha(x; a) := \sum_{n \leq x}^a \frac{\alpha(n)}{n}.$$

It is clear from (3.4) that  $S(x; a) \leq S(x; 1) \ll x(\log x)^{\beta-1}$  and  $T(x; a) \leq T(x; 1) \ll (\log x)^\beta$ . Moreover, it is shown in the the proof of [6, Theorem 2.1] that

$$T(x; 1) = \left( 1 + O\left(\frac{1}{\log x}\right) \right) \frac{\lambda_\alpha}{\beta} (\log x)^\beta. \quad (3.7)$$



Following the proof of [6, Theorem 2.1], we find

$$\begin{aligned}
S(x; a) \log x &= \sum_{n \leq x}^a \alpha(n) \log n + \sum_{n \leq x}^a \alpha(n) \log \frac{x}{n} \\
&= \sum_{k \leq x}^a \alpha(k) \sum_{\substack{p^\nu \leq x/k \\ p \nmid k}}^a \alpha(p^\nu) \log p^\nu + \int_{1^-}^x \frac{S(t; a)}{t} dt \\
&= \sum_{k \leq x}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p \nmid k}}^a \alpha(p) \log p + O \left( \sum_{k \leq x}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p \nmid k}}^a \alpha(p) \log p \right) \\
&\quad + O \left( \sum_{k \leq x}^a \alpha(k) \sum_{\substack{p^\nu \leq x/k \\ \nu \geq 2}}^a \alpha(p^\nu) \log p^\nu \right) + O(x(\log x)^{\beta-1}) \\
&= \beta x T(x, a) - \sum_{k \leq x}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p \mid a}}^a \alpha(p) \log p + O \left( x \sum_{k \leq x}^a \frac{\alpha(k)}{k(\log(3x/k))^{A_0}} \right) \\
&\quad + O(x(\log x)^{\beta-1}). \tag{3.8}
\end{aligned}$$

By partial summation we have

$$\begin{aligned}
\sum_{k \leq x} \frac{\alpha(k)}{k(\log(3x/k))^{A_0}} &= \frac{S(x)}{x(\log 3)^{A_0}} + \int_{1^-}^x \frac{\log(3x/t) - A_0}{t^2(\log(3x/t))^{A_0+1}} S(t) dt \\
&\ll (\log x)^{\beta-1} + \int_1^x \frac{(\log 3t)^{\beta-1}}{t(\log(3x/t))^{A_0}} dt \\
&= (\log x)^{\beta-1} + \int_0^{\log x} \frac{(\log 3 + t)^{\beta-1}}{(\log 3x - t)^{A_0}} dt \\
&\ll (\log x)^{\beta-1} + \frac{1}{(\log x)^{A_0}} \int_0^{(\log x)/2} (\log 3 + t)^{\beta-1} dt \\
&\quad + (\log x)^{\beta-1} \int_{(\log x)/2}^{\log x} \frac{1}{(\log 3x - t)^{A_0}} dt \\
&\ll (\log x)^{\beta-A_0}. \tag{3.9}
\end{aligned}$$

Let  $x_1 := x/(\log x)^2$ . For  $k \leq x_1$  we see that

$$\sum_{\substack{p \leq x/k \\ p \mid a}} \alpha(p) \log p \ll (\log \log x)^{\vartheta_0} \log a \ll \frac{x}{k(\log(x/k))^{A_0}},$$

so that

$$\sum_{k \leq x_1}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p \mid a}}^a \alpha(p) \log p \ll x \sum_{k \leq x}^a \frac{\alpha(k)}{k(\log(3x/k))^{A_0}} \ll x(\log x)^{\beta-A_0}. \tag{3.10}$$

On the other hand, we have by (2.2) that

$$\begin{aligned} \sum_{x_1 < k \leq x}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p|a}} \alpha(p) \log p &\ll x \sum_{x_1 < k \leq x} \frac{\alpha(k)}{k} \ll x ((\log x)^\beta - (\log x_1)^\beta + O((\log x)^{\beta-1})) \\ &\ll x(\log x)^{\beta-1} \log \log x, \end{aligned} \quad (3.11)$$

where we have used (3.7) to estimate the sum over  $k$  and the mean value theorem to get

$$(\log x)^\beta - (\log x_1)^\beta = \beta \xi^{\beta-1} \log \frac{x}{x_1} \ll (\log x)^{\beta-1} \log \log x$$

for some  $\xi \in (\log x_1, \log x)$ . Combining (3.10) with (3.11), we obtain

$$\sum_{k \leq x}^a \alpha(k) \sum_{\substack{p \leq x/k \\ p|a}} \alpha(p) \log p \ll x(\log x)^{\beta-A_0}.$$

Inserting this and (3.9) into (3.8) yields

$$S(x; a) = \frac{\beta x}{\log x} T(x; a) + O(x(\log x)^{\beta-1-A_0}) \quad (3.12)$$

uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ .

To estimate  $T(x; a)$ , we repeat the argument above with  $\alpha(n)$  replaced by  $\alpha(n)/n$ . From (2.2) it follows that

$$\sum_{p \leq x} \frac{\alpha(p)}{p} \log p = \beta \log x + O((\log x)^{1-A_0}). \quad (3.13)$$

Thus, we have

$$\begin{aligned} T(x; a) \log x &= \sum_{n \leq x}^a \frac{\alpha(n)}{n} \log n + \sum_{n \leq x}^a \frac{\alpha(n)}{n} \log \frac{x}{n} \\ &= \sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{\substack{p^\nu \leq x/k \\ p \nmid k}}^a \frac{\alpha(p^\nu)}{p^\nu} \log p^\nu + U(x; a) \\ &= \sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{p \leq x/k}^a \frac{\alpha(p)}{p} \log p + O \left( \sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{\substack{p \leq x/k \\ p|k}}^a \frac{\alpha(p)}{p} \log p \right) \\ &\quad + O \left( \sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{\substack{p^\nu \leq x/k \\ \nu \geq 2}}^a \frac{\alpha(p^\nu)}{p^\nu} \log p^\nu \right) + U(x; a) \\ &= (\beta + 1)U(x; a) - \sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{\substack{p \leq x/k \\ p|a}}^a \frac{\alpha(p)}{p} \log p + O((\log x)^{1-A_0} T(x; a)), \end{aligned}$$

where

$$U(x; a) := \sum_{n \leq x}^a \frac{\alpha(n)}{n} \log \frac{x}{n} = \int_{1^-}^x \frac{T(t; a)}{t} dt. \quad (3.14)$$

In view of (3.13), we have

$$\begin{aligned} \sum_{\substack{p \leq x/k \\ p|a}} \frac{\alpha(p)}{p} \log p &\leq \sum_{p \leq (\log x)^2} \frac{\alpha(p)}{p} \log p + \sum_{\substack{(\log x)^2 < p \leq x \\ p|a}} \frac{\alpha(p)}{p} \log p \\ &\ll \log \log x + (\log \log x)^{\vartheta_0} \sum_{\substack{(\log x)^2 < p \leq x \\ p|a}} \frac{\log p}{p} \\ &\ll \log \log x + (\log \log x)^{\vartheta_0} \omega(a) \frac{\log \log x}{(\log x)^2} \ll \log \log x, \end{aligned}$$

so that

$$\sum_{k \leq x}^a \frac{\alpha(k)}{k} \sum_{\substack{p \leq x/k \\ p|a}} \frac{\alpha(p)}{p} \log p \ll (\log \log x) T(x; a).$$

It follows that

$$T(x; a) \log x = (\beta + 1) U(x; a) + O((\log x)^{1-A_0} T(x; a)).$$

Hence, there exists a function  $\epsilon(x; a)$  such that  $\epsilon(x; a) = O((\log x)^{-A_0})$  and

$$T(x; a) = \frac{1}{1 - \epsilon(x; a)} \cdot \frac{\beta + 1}{\log x} U(x; a) \quad (3.15)$$

uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ .

Finally, we estimate  $U(x; a)$  and  $T(x; a)$  by following the proof of [31, Theorem A]. For  $y \geq 2$  and  $a \in \mathbb{N} \cap [1, y]$ , let

$$V(y; a) := \log \left( \frac{\beta + 1}{(\log y)^{\beta+1}} U(y; a) \right).$$

In light of (3.14) and (3.15), we have

$$\begin{aligned} \frac{d}{dy} V(y; a) &= -\frac{\beta + 1}{y \log y} + \frac{1}{U(y; a)} \cdot \frac{d}{dy} U(y; a) \\ &= -\frac{\beta + 1}{y \log y} + \frac{T(y; a)}{U(y; a) y} \\ &= \frac{\beta + 1}{y \log y} \cdot \frac{\epsilon(y; a)}{1 - \epsilon(y; a)} \ll \frac{1}{y (\log y)^{A_0+1}} \end{aligned}$$

uniformly for all sufficiently large  $y$  and all  $a \in \mathbb{N} \cap [1, y]$ , which implies that

$$V_a := \int_2^\infty \frac{d}{dy} V(y; a) dy < \infty.$$

Since

$$V(x; a) - V(2; a) = V_a - \int_x^\infty \frac{d}{dy} V(y; a) dy = V_a + O((\log x)^{-A_0})$$

uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ , it follows that

$$\frac{\beta + 1}{(\log x)^{\beta+1}} U(x; a) = \exp(V(x; a)) = \exp(V_a + V(2; a)) (1 + O((\log x)^{-A_0})).$$

Combining this estimate with (3.15), we infer

$$T(x; a) = \exp(V_a + V(2; a)) (\log x)^\beta (1 + O((\log x)^{-A_0})) \quad (3.16)$$

uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ . The leading coefficient can be made explicit by arguing as in the proof of [31, Theorem A]. Alternatively, we can also take advantage of (3.6). Fixing  $a \in \mathbb{N}$ , we have by (3.6) with  $\sigma_0 = 1$  that

$$T(x; a) = \frac{\lambda_\alpha(a)}{\beta} (\log x)^\beta (1 + O((\log x)^{-A_0}))$$

for all sufficiently large  $x$ . Comparing this with (3.16) shows that  $\exp(V_a + V(2; a)) = \lambda_\alpha(a)/\beta$ . Carrying this back into (3.16), we obtain

$$T(x; a) = \frac{\lambda_\alpha(a)}{\beta} (\log x)^\beta (1 + O((\log x)^{-A_0}))$$

uniformly for all sufficiently large  $x$  and all  $a \in \mathbb{N} \cap [1, x]$ . Inserting the above into (3.12) completes the proof Lemma 3.2.  $\square$

The next result, which is key to the computation of moments, is a direct corollary of Lemmas 3.1 and 3.2.

**Lemma 3.3.** *Fix  $\epsilon_0 \in (0, 1)$ , and let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ . Then there exist constants  $\delta_0 > 0$  and  $Q_0 \geq 2$ , such that uniformly for all sufficiently large  $x$ , any  $\delta \in [\delta_0 \log \log x / \log x, 1]$ , and any square-free  $a \in \mathbb{N} \cap [1, x]$  with  $\omega(a) \leq (1 - \rho_0)\epsilon_0\delta^{-1}$ ,  $P^-(a) > Q_0$  and  $P^+(a) \leq x^\delta$ , we have*

$$\sum_{\substack{n \leq x \\ a|n}} \alpha(n) = \lambda_\alpha \left( F(\sigma_0, a) + O \left( \frac{2^{O(\omega(a))} L(a)}{a} \left( \frac{1}{(\log x)^{A_0}} + \frac{\epsilon_{\beta,1} \log P^+(a)}{\log x} \right) \right) \right) x^{\sigma_0} (\log x)^{\beta-1},$$

where  $L(a)$  is defined as in Lemma 3.1,

$$F(\sigma_0, a) := \prod_{p|a} \left( 1 - \left( \sum_{\nu=0}^{\infty} \alpha(p^\nu) p^{-\sigma_0 \nu} \right)^{-1} \right),$$

and  $\lambda_\alpha$  is defined by (3.5).

*Proof.* Suppose that  $\delta_0 > 0$  is a constant for which Lemma 3.1 holds when  $c_0 = 1$  and  $h \in \{\beta - 1, \beta - 1 - A_0\}$ . Let  $Q_0 \geq 2$  be such that

$$\sum_{\nu=1}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} \leq \frac{1}{2}$$

for all  $p > Q_0$ . Then we have

$$F(\sigma_0, p) = \sum_{\nu=1}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} + O \left( \left( \sum_{\nu=1}^{\infty} \frac{\alpha(p^\nu)}{p^{\sigma_0 \nu}} \right)^2 \right) = \frac{\alpha(p)}{p^{\sigma_0}} + O \left( \psi_0(p) + \frac{\alpha(p)^2}{p^{2\sigma_0}} \right) \quad (3.17)$$

for all  $p > Q_0$ . For any square-free integer  $a \in [1, x]$  with  $\omega(a) \leq (1 - \varrho_0)\epsilon_0\delta^{-1}$ ,  $P^-(a) > Q_0$  and  $P^+(a) \leq x^\delta$ , we have by Lemma 3.2 that

$$\sum_{\substack{n \leq x \\ \gcd(n, a) = 1}} \alpha(n) = \lambda_\alpha(a) x^{\sigma_0} (\log 3x)^{\beta-1} \left( 1 + O\left(\frac{1}{(\log 3x)^{A_0}}\right) \right). \quad (3.18)$$

Note that

$$\sum_{\substack{n \leq x \\ a|n}} \alpha(n) = \sum_{\substack{q \leq x \\ R_q = a}} \alpha(q) \sum_{\substack{n' \leq x/q \\ \gcd(n', a) = 1}} \alpha(n').$$

By (3.18), the main term of the inner sum contributes

$$\lambda_\alpha(a) x^{\sigma_0} \sum_{\substack{q \leq x \\ R_q = a}} \alpha(q) \left( \log \frac{3x}{q} \right)^{\beta-1},$$

which, by Lemma 3.1, is equal to

$$\begin{aligned} & \lambda_\alpha(a) x^{\sigma_0} \left( \tilde{\lambda}_\alpha(a) + O\left(\frac{2^{O(\omega(a))}}{\log x} \left( \frac{1}{x^{\delta\omega(a)}} + \frac{\epsilon_{\beta,1} L(a) \log P^+(a)}{a} \right) \right) \right) (\log x)^{\beta-1} \\ &= \lambda_\alpha \left( F(\sigma_0, a) + O\left(\frac{2^{O(\omega(a))}}{\log x} \left( \frac{1}{a} + \frac{\epsilon_{\beta,1} L(a) \log P^+(a)}{a} \right) \right) \right) x^{\sigma_0} (\log x)^{\beta-1}, \end{aligned}$$

since  $a \leq x^{\delta\omega(a)}$ . Analogously, the contribution from the error term of the inner sum is

$$\begin{aligned} & \ll \lambda_\alpha \left( F(\sigma_0, a) + \frac{2^{O(\omega(a))} L(a) \log P^+(a)}{a \log x} \right) x^{\sigma_0} (\log x)^{\beta-1-A_0} \\ & \ll \frac{\lambda_\alpha 2^{O(\omega(a))} L(a)}{a} x^{\sigma_0} (\log x)^{\beta-1-A_0}, \end{aligned}$$

where we have used the estimate  $F(\sigma_0, a) \ll 2^{O(\omega(a))} L(a)/a$ , which follows directly from (2.4) and (3.17). Combining these estimates completes the proof of Lemma 3.3.  $\square$

*Remark 3.1.* We point out that the lower bound  $Q_0$  for  $\omega(a)$  in the lemma above is by and large an artificial thing, whose value is insignificant for our applications. However, we need it because (3.17) may not hold for small primes. As we shall see later, having such a lower bound also frees us from dealing with minor contributions from small primes.

#### 4. COMPUTING MOMENTS

By rescaling the strongly additive function  $f$  in Theorem 2.1, we may assume, without loss of generality, that  $|f(p)| \leq 1$  for all primes  $p$ . Note that  $0 \leq F(\sigma_0, p) < 1$  for all primes  $p$ . For every  $p$  we define  $f_p: \mathbb{N} \rightarrow \mathbb{R}$  by

$$f_p(n) := \begin{cases} f(p)(1 - F(\sigma_0, p)), & \text{if } p \mid n, \\ -f(p)F(\sigma_0, p), & \text{otherwise.} \end{cases}$$

Given any  $q \in \mathbb{N}$  we may also extend  $f_p$  via complete multiplicativity by setting

$$f_q(n) := \prod_{p^\nu \parallel q} f_p(n)^\nu.$$

It is clear that  $|f_q(n)| \leq 1$ . The following result provides an approximation of the moments of  $f$  in terms of those of  $f_p$ .

**Lemma 4.1.** *Let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ , and let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be a strongly additive function with  $|f(p)| \leq 1$  for all  $p$ . Then there exists a constant  $Q_0 \geq 2$ , such that*

$$\sum_{n \leq y} \alpha(n) (f(n) - A(x))^m = \sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m + O(E(y, z, w; m))$$

holds uniformly for all sufficiently large  $x \geq z$ , any  $y \geq 1$ , and all  $m \in \mathbb{N}$ , where

$$E(y, z, w; m) := \sum_{\substack{a+b+c=m \\ 0 \leq a < m \\ b, c \geq 0}} \binom{m}{a, b, c} 2^{O(m-a)} (\log(v+2))^c \sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \omega(n; z, w)^b,$$

$v := \log x / \log z$ ,  $w := x^{1/\log(v+2)}$ , and

$$\omega(n; z, w) := \sum_{\substack{z < p \leq w \\ p|n}} 1.$$

*Proof.* Let  $Q_0 \geq 2$  be a constant for which (3.17) holds. Suppose that  $z > Q_0$  is sufficiently large. By (2.4), (3.17) and the fact that  $\sum_p \psi_0(p) < \infty$ , we find

$$\sum_{Q_0 < p \leq x} f(p) F(\sigma_0, p) = A(x) + O(1).$$

We compute

$$\begin{aligned} f(n) - A(x) &= \sum_{\substack{p|n \\ p > Q_0}} f(p) - \sum_{Q_0 < p \leq x} f(p) F(\sigma_0, p) + O(1) \\ &= \sum_{\substack{Q_0 < p \leq z \\ p|n}} f(p) + \sum_{\substack{p > z \\ p|n}} f(p) - \sum_{Q_0 < p \leq z} f(p) F(\sigma_0, p) - \sum_{z < p \leq x} f(p) F(\sigma_0, p) + O(1) \\ &= \sum_{Q_0 < p \leq z} f_p(n) + \sum_{\substack{p > z \\ p|n}} f(p) - \sum_{z < p \leq x} f(p) F(\sigma_0, p) + O(1). \end{aligned}$$

By (3.1) we have

$$\left| \sum_{z < p \leq x} f(p) F(\sigma_0, p) \right| \leq \sum_{z < p \leq x} \frac{\alpha(p)}{p^{\sigma_0}} + O(1) \leq \beta \log(v+2) + O(1).$$

Since

$$\sum_{\substack{p > z \\ p|n}} |f(p)| \leq \sum_{\substack{z < p \leq x \\ p|n}} 1 < \omega(n; z, w) + \log(v+2),$$

it follows that

$$f(n) - A(x) = \sum_{Q_0 < p \leq z} f_p(n) + O(\omega(n; z, w) + \log(v + 2)).$$

We have therefore proved

$$\sum_{n \leq y} \alpha(n) (f(n) - A(x))^m = \sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) + O(\omega(n; z, w) + \log(v + 2)) \right)^m.$$

Opening the  $m$ th power on the right-hand side by means of the multinomial theorem completes the proof of Lemma 4.1.  $\square$

Let  $z = x^{1/v}$  and  $w = x^{1/\log(v+2)}$  be as in Lemma 4.1, where  $v \geq 1$  is a function of  $x$  and  $m$  to be chosen later. Fix  $\epsilon_0 \in (0, 1)$  and  $\eta_0 \in (0, 1]$ , and suppose that  $y \in [x^{\eta_0}, x]$ . Under the hypotheses in Theorem 2.1, we seek to estimate the weighted moments

$$\sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m,$$

appearing in Lemma 4.1. Expanding out the  $m$ th power we see that

$$\sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m = \sum_{Q_0 < p_1, \dots, p_m \leq z} \sum_{n \leq y} \alpha(n) f_{p_1 \dots p_m}(n). \quad (4.1)$$

This suggests that we study the sum

$$\sum_{n \leq y} \alpha(n) f_q(n)$$

for  $q \in \mathbb{N}$  with  $\omega(q) \leq m$ ,  $P^-(q) > Q_0$  and  $P^+(q) \leq z$ . A key observation is that  $f_q(n) = f_q(\gcd(n, R_q))$ . From this we deduce

$$\sum_{n \leq y} \alpha(n) f_q(n) = \sum_{a|R_q} f_q(a) \sum_{\substack{n \leq y \\ \gcd(n, R_q) = a}} \alpha(n) = \sum_{ab|R_q} f_q(a) \mu(b) \sum_{\substack{n \leq y \\ ab|n}} \alpha(n).$$

Note that  $\log y / \log z \in [\eta_0 v, v]$ . By Lemma 3.3, there exists a constant  $v_0 > 0$ , independent of  $Q_0$  and  $\eta_0$ , such that

$$\sum_{n \leq y} \alpha(n) f_q(n) = \lambda_\alpha \left( G(\sigma_0, q) + O(2^{O(m)} E_y(q)) \right) y^{\sigma_0} (\log y)^{\beta-1} \quad (4.2)$$

holds uniformly for all sufficiently large  $x$ , any  $y \in [x^{\eta_0}, x]$  and  $v \in [\eta_0^{-1}, v_0 \log x / \log \log x]$ , and all  $m \leq (1 - \varrho_0) \epsilon_0 \log y / \log z$ , where

$$G(\sigma_0, q) := \sum_{ab|R_q} f_q(a) \mu(b) F(\sigma_0, ab),$$

$$E_y(q) := \sum_{ab|R_q} \frac{|f_q(a)| L(ab)}{ab} \left( \frac{1}{(\log y)^{A_0}} + \frac{\epsilon_{\beta,1} \log P^+(ab)}{\log y} \right).$$

Combining (4.2) with (4.1) gives

$$\sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m = \lambda_\alpha \left( G(z) + O\left(2^{O(m)} D(y, z)\right) \right) y^{\sigma_0} (\log y)^{\beta-1}, \quad (4.3)$$

where

$$G(z) := \sum_{Q_0 < p_1, \dots, p_m \leq z} G(\sigma_0, p_1 \cdots p_m),$$

$$D(y, z) := \sum_{Q_0 < p_1, \dots, p_m \leq z} E_y(p_1 \cdots p_m).$$

## 5. ESTIMATION OF $G(z)$ AND $D(y, z)$

It is easy to see that  $G(\sigma_0, q)$  is multiplicative as a function of  $q$ . Indeed, given any  $q_1, q_2 \in \mathbb{N}$  with  $\gcd(q_1, q_2) = 1$ , we have

$$\begin{aligned} G(\sigma_0, q_1)G(\sigma_0, q_2) &= \sum_{\substack{a_1 b_1 | R_{q_1} \\ a_2 b_2 | R_{q_2}}} f_{q_1}(a_1) f_{q_2}(a_2) \mu(b_1) \mu(b_2) F(\sigma_0, a_1 b_1) F(\sigma_0, a_2 b_2) \\ &= \sum_{\substack{a_1 b_1 | R_{q_1} \\ a_2 b_2 | R_{q_2}}} f_{q_1}(a_1 a_2) f_{q_2}(a_1 a_2) \mu(b_1 b_2) F(\sigma_0, a_1 a_2 b_1 b_2) \\ &= \sum_{\substack{a_1 b_1 | R_{q_1} \\ a_2 b_2 | R_{q_2}}} f_{q_1 q_2}(a_1 a_2) \mu(b_1 b_2) F(\sigma_0, a_1 a_2 b_1 b_2) \\ &= \sum_{ab | R_{q_1 q_2}} f_{q_1 q_2}(a) \mu(b) F(\sigma_0, ab) = G(\sigma_0, q_1 q_2). \end{aligned}$$

Furthermore, we have

$$\begin{aligned} G(\sigma_0, p^\nu) &= f_{p^\nu}(1) + f_{p^\nu}(p) F(\sigma_0, p) - f_{p^\nu}(1) F(\sigma_0, p) \\ &= (-f(p) F(\sigma_0, p))^\nu + (f(p)(1 - F(\sigma_0, p)))^\nu F(\sigma_0, p) - (-f(p) F(\sigma_0, p))^\nu F(\sigma_0, p) \\ &= f(p)^\nu F(\sigma_0, p)(1 - F(\sigma_0, p)) \left( (-1)^\nu F(\sigma_0, p)^{\nu-1} + (1 - F(\sigma_0, p))^{\nu-1} \right) \end{aligned}$$

for all prime powers  $p^\nu$ . Note that  $G(\sigma_0, p) = 0$ ,  $|G(\sigma_0, p^\nu)| \leq 1/4$ , and  $G(\sigma_0, p^\nu) \geq 0$  when  $2 \mid \nu$ . In addition, we have by (3.17) that

$$G(\sigma_0, p^2) = f(p)^2 F(\sigma_0, p)(1 - F(\sigma_0, p)) = \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} + O\left(\psi_0(p) + \frac{\alpha(p)^2}{p^{2\sigma_0}}\right) \quad (5.1)$$

and that

$$|G(\sigma_0, p^\nu)| \leq |f(p)|^\nu F(\sigma_0, p) \leq \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} + O\left(\psi_0(p) + \frac{\alpha(p)^2}{p^{2\sigma_0}}\right) \quad (5.2)$$

for all  $p^\nu$  with  $p > Q_0$  and  $\nu \geq 2$ .

Now we proceed to estimate  $G(z)$  in the main term of (4.3). Recall that  $y \in [x^{\eta_0}, x]$  and  $z = x^{1/v}$ . We shall suppose in this section that  $1 \leq m \leq \min(v, h_0 B(x)^{1/3})$ ,  $\log(v+2) = o(B(x))$ , and  $m \log(v+2) \ll B(x)$ , where  $0 < h_0 < (3/2)^{2/3}$  is any given constant, and



obtain a uniform treatment for  $G(z)$  and  $D(y, z)$  under this more general assumption. Since  $G(\sigma_0, q)$  is multiplicative in  $q$  and  $G(\sigma_0, p) = 0$  for all  $p > Q_0$ , we have

$$G(z) = \sum_{\substack{Q_0 < p_1, \dots, p_m \leq z \\ p_1 \cdots p_m \text{ squareful}}} G(\sigma_0, p_1 \cdots p_m). \quad (5.3)$$

When  $2 \mid m$ , the main contribution arises from

$$\frac{m!}{(m/2)! 2^{m/2}} \sum_{\substack{Q_0 < p_1, \dots, p_{m/2} \leq z \\ p_1, \dots, p_{m/2} \text{ distinct}}} G(\sigma_0, p_1^2 \cdots p_{m/2}^2) = C_m \sum_{\substack{Q_0 < p_1, \dots, p_{m/2} \leq z \\ p_1, \dots, p_{m/2} \text{ distinct}}} \prod_{i=1}^{m/2} G(\sigma_0, p_i^2), \quad (5.4)$$

since the number of ways to partition a set of  $m$  elements into  $m/2$  two-element equivalence classes is

$$\frac{m!}{(m/2)! 2^{m/2}} = \frac{m!}{m!!} = C_m.$$

The sum on the right-hand side of (5.4) can be rewritten as

$$\sum_{\substack{Q_0 < p_1, \dots, p_{m/2-1} \leq z \\ p_1, \dots, p_{m/2-1} \text{ distinct}}} \prod_{i=1}^{m/2-1} G(\sigma_0, p_i^2) \sum_{\substack{Q_0 < p_{m/2} \leq z \\ p_{m/2} \neq p_1, \dots, p_{m/2-1}}} G(\sigma_0, p_{m/2}^2).$$

By (5.1) and (3.1), the inner sum over  $p_{m/2}$  is equal to

$$\sum_{Q_0 < p \leq z} G(\sigma_0, p^2) - \sum_{i=1}^{m/2-1} G(\sigma_0, p_i^2) = B(z) + O\left(\sum_{Q_0 < p \leq q_N} \frac{\alpha(p)}{p^{\sigma_0}}\right) = B(z) + O(\log \log(m+2)),$$

where  $N = m/2 + \pi(Q_0)$  and  $q_N$  is the  $N$ th prime. Repeating this argument we obtain

$$\sum_{\substack{Q_0 < p_1, \dots, p_{m/2} \leq z \\ p_1, \dots, p_{m/2} \text{ distinct}}} \prod_{i=1}^{m/2} G(\sigma_0, p_i^2) = (B(z) + O(\log \log(m+2)))^{m/2}.$$

But

$$B(x) - B(z) = \sum_{z < p \leq x} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} \leq \beta \log(v+2) + O(1).$$

Hence when  $m$  is even, the main contribution to  $G(z)$  is given by

$$C_m (B(x) + O(\log(v+2)))^{m/2} = C_m B(x)^{\frac{m}{2}} (1 + O(mB(x)^{-1} \log(v+2))).$$

The remaining contribution to  $G(z)$  comes from

$$\sum_{s < m/2} \sum_{Q_0 < p_1 < \cdots < p_s \leq z} \sum_{\substack{k_1 + \cdots + k_s = m \\ k_1, \dots, k_s \geq 2}} \binom{m}{k_1, \dots, k_s} \prod_{i=1}^s G(\sigma_0, p_i^{k_i}). \quad (5.5)$$

Since (5.5) vanishes when  $m \leq 2$ , we may suppose  $m \geq 3$ . By (5.2) we see that

$$\prod_{i=1}^s |G(\sigma_0, p_i^{k_i})| \leq \prod_{i=1}^s \left( \alpha(p_i) \frac{f(p_i)^2}{p_i^{\sigma_0}} + O\left(\psi_0(p_i) + \frac{\alpha(p_i)^2}{p_i^{2\sigma_0}}\right) \right).$$

Thus, we have

$$\sum_{Q_0 < p_1 < \dots < p_s \leq z} \prod_{i=1}^s |G(\sigma_0, p_i^{k_i})| \leq \frac{1}{s!} (B(x) + O(1))^s = \frac{1}{s!} B(x)^s (1 + O(sB(x)^{-1})) \ll \frac{B(x)^s}{s!}.$$

Since

$$\sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \geq 2}} \binom{m}{k_1, \dots, k_s} \leq \frac{m!}{2^s} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \geq 2}} 1 = \frac{m!}{2^s} \binom{m-s-1}{s-1},^2$$

(5.5) is

$$\ll m! \sum_{s < m/2} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} B(x)^s.$$

To estimate the sum above, we put  $m_1 := \lfloor (m-1)/2 \rfloor$  and observe that

$$\begin{aligned} \sum_{s < m/2} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} B(x)^s &= B(x)^{m_1} \sum_{s \leq m_1} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} B(x)^{s-m_1} \\ &\leq B(x)^{m_1} m^{-3m_1} \sum_{s \leq m_1} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} h_0^{3(m_1-s)} m^{3s}, \end{aligned}$$

where we have used the assumption that  $B(x) \geq m^3/h_0^3$  with some  $0 < h_0 < (3/2)^{2/3}$ . Let

$$e_m := \begin{cases} 1, & \text{if } 2 \mid m, \\ 1/2, & \text{otherwise.} \end{cases}$$

Then  $m_1 = m/2 - e_m$ . Note that

$$\begin{aligned} m^{-3m_1} \sum_{s \leq m/4} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} h_0^{3(m_1-s)} m^{3s} &\leq m^{-3m_1} \sum_{s \leq m/4} \frac{1}{s! (s-1)!} \left(\frac{9}{4}\right)^{m_1-s} \left(\frac{m^4}{2}\right)^s \\ &\ll m^{-3m_1} \left(\frac{9}{4}\right)^{m_1} \left(\frac{m^4}{2}\right)^{m/4} \ll \frac{C_m}{m!} m^{3e_m}, \end{aligned}$$

since

$$\frac{C_m}{m!} = \frac{1}{2^{m/2} \Gamma(m/2 + 1)} \asymp m^{-\frac{m+1}{2}} e^{\frac{m}{2}}$$

by Stirling's formula. Next, we have

$$\begin{aligned} m^{-3m_1} \sum_{m/4 < s \leq m/3} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} h_0^{3(m_1-s)} m^{3s} &\leq 2^{O(m)} m^{-3m_1} \sum_{m/4 < s \leq m/3} \frac{1}{s! (s-1)!} \left(\frac{m^4}{2}\right)^s \\ &\leq \frac{2^{O(m)} m^{-3m_1}}{m^{m/2}} \sum_{m/4 < s \leq m/3} \left(\frac{m^4}{2}\right)^s \\ &\leq \frac{2^{O(m)} m^{-3m_1}}{m^{m/2}} m^{4m/3} \\ &= 2^{O(m)} m^{-2m/3+3e_m} \ll \frac{C_m}{m!} m^{3e_m}. \end{aligned}$$

---

<sup>2</sup>We have corrected the binomial coefficient in [18, Equ. (11)].

Finally, we observe that

$$\begin{aligned}
& m^{-3m_1} \sum_{m/3 < s \leq m_1} \frac{1}{s! 2^s} \binom{m-s-1}{s-1} h_0^{3(m_1-s)} m^{3s} \\
&= m^{-3m_1} \sum_{m/3 < s \leq m_1} \frac{1}{s! 2^s} \binom{m-s-1}{m-2s} h_0^{3(m_1-s)} m^{3s} \\
&\leq m^{-3m_1} \sum_{m/3 < s \leq m_1} \frac{1}{s! 2^s} (m-s)^{m-2s} h_0^{3(m_1-s)} m^{3s} \\
&\leq \frac{m^{-3m_1}}{m_1!} \sum_{m/3 < s \leq m_1} \frac{m_1!}{s! 2^s} \left(\frac{2m}{3}\right)^{m-2s} h_0^{3(m_1-s)} m^{3s} \\
&\leq \frac{m^{-3m_1}}{m_1!} \sum_{m/3 < s \leq m_1} \frac{1}{2^s} \left(\frac{m}{2}\right)^{m_1-s} \left(\frac{2m}{3}\right)^{m-2s} h_0^{3(m_1-s)} m^{3s} \\
&\ll \frac{m^{m-2m_1}}{m_1! 2^{m/2}} \sum_{m/3 < s \leq m_1} \left(\frac{2h_0^{3/2}}{3}\right)^{m-2s} \ll \frac{C_m}{m!} m^{3e_m}.
\end{aligned}$$

Collecting the estimates above, we see that the contribution to  $G(z)$  from (5.5) is

$$\ll C_m m^{3e_m} B(x)^{m_1} = C_m B(x)^{\frac{m}{2}} \left(\frac{m^3}{B(x)}\right)^{e_m} \leq C_m B(x)^{\frac{m}{2}} \frac{m^{\frac{3}{2}}}{\sqrt{B(x)}}.$$

We can therefore conclude that

$$G(z) = C_m B(x)^{\frac{m}{2}} \left( \chi_m \left( 1 + O\left(\frac{m \log(v+2)}{B(x)}\right) \right) + O\left(\frac{m^{\frac{3}{2}}}{\sqrt{B(x)}}\right) \right). \quad (5.6)$$

Next, we estimate  $D(y, z)$  in the error term of (4.3). By definition, we have

$$D(y, z) = \sum_{s \leq m} \sum_{Q_0 < p_1 < \dots < p_s \leq z} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} E_y(p_1^{k_1} \dots p_s^{k_s}).$$

Let

$$H(\sigma_0, q) := \sum_{ab \mid R_q} \frac{|f_q(a)| L(ab)}{ab}.$$

Then  $H(\sigma_0, q)$  is multiplicative in  $q$ . Moreover, we have

$$E_y(q) \leq H(\sigma_0, q) \left( \frac{1}{(\log y)^{A_0}} + \frac{\epsilon_{\beta,1} \log P^+(q)}{\log y} \right).$$

It follows that  $D(y, z) \leq D_1(y, z) + \epsilon_{\beta,1} D_2(y, z)$ , where

$$D_1(y, z) := \frac{1}{(\log y)^{A_0}} \sum_{s \leq m} \sum_{Q_0 < p_1 < \dots < p_s \leq z} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \prod_{i=1}^s H(\sigma_0, p_i^{k_i}),$$

$$D_2(y, z) := \frac{1}{\log y} \sum_{s \leq m} \sum_{Q_0 < p_1 < \dots < p_s \leq z} \log p_s \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \prod_{i=1}^s H(\sigma_0, p_i^{k_i}).$$

By Mertens' theorems [23, Theorems 425, 427] we have, for any  $t \geq 3$ , that

$$\sum_{p \leq t} \frac{(\log \log(p+1))^{\vartheta_0}}{p} = \frac{1}{\vartheta_0 + 1} (\log \log t)^{\vartheta_0 + 1} + O(1) \quad (5.7)$$

and that

$$\sum_{p \leq t} \frac{(\log \log(p+1))^{\vartheta_0} \log p}{p} = \left(1 + O\left(\frac{1}{\log t} + \frac{\vartheta_0}{\log \log t}\right)\right) (\log \log t)^{\vartheta_0} \log t. \quad (5.8)$$

Furthermore, let

$$T_n(t) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} t^k$$

denote the  $n$ th Touchard polynomial, where

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} := \frac{1}{k!} \sum_{\substack{n_1 + \dots + n_k = n \\ n_1, \dots, n_k \in \mathbb{N}}} \binom{n}{n_1, \dots, n_k}$$

is the  $k$ th Stirling number of the second kind of size  $n$ . The sequence  $\{T_n(t)\}_{n=0}^{\infty}$  of the Touchard polynomials is known to satisfy the recurrence relation

$$T_{n+1}(t) = t \sum_{i=0}^n \binom{n}{i} T_i(t),$$

from which one verifies readily by induction that

$$T_n(t) \leq \left(t + \frac{n-1}{2}\right)^n \quad (5.9)$$

for all  $n \geq 1$  and  $t \geq 0$ . Since

$$\begin{aligned} H(\sigma_0, p^\nu) &= |f(p)| F(\sigma_0, p) \left(1 + \frac{L(p)}{p}\right) + \frac{|f(p)| L(p)}{p} (1 - F(\sigma_0, p)) \\ &= |f(p)| \left(F(\sigma_0, p) + \frac{(\log \log(p+1))^{\vartheta_0}}{p}\right) \end{aligned}$$

for any prime powers  $p^\nu$  with  $p > Q_0$ , we obtain, from (3.17), (5.7), (5.8) and (5.9), that

$$\begin{aligned} D_1(y, z) &\leq \frac{2^{O(m)}}{(\log x)^{A_0}} \sum_{s \leq m} \frac{1}{s!} (\log \log z)^{s(\vartheta_0+1)} \sum_{\substack{k_1+\dots+k_s=m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \\ &\leq \frac{2^{O(m)}}{(\log x)^{A_0}} T_m((\log \log z)^{\vartheta_0+1}) \leq \frac{2^{O(m)}}{(\log x)^{A_0}} (\log \log x)^{m(\vartheta_0+1)}, \end{aligned}$$

and that

$$\begin{aligned} D_2(y, z) &\leq \frac{2^{O(m)} \log z}{\log x} \sum_{s \leq m} \frac{1}{(s-1)!} (\log \log z)^{s(\vartheta_0+1)-1} \sum_{\substack{k_1+\dots+k_s=m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \\ &\leq \frac{2^{O(m)}}{v \log \log z} T_m((\log \log z)^{\vartheta_0+1}) \leq \frac{2^{O(m)}}{v} (\log \log x)^{m(\vartheta_0+1)-1}. \end{aligned}$$

Hence, we conclude that

$$D(y, z) \leq 2^{O(m)} (\log \log x)^{m(\vartheta_0+1)-1} \left( \frac{\log \log x}{(\log x)^{A_0}} + \frac{\epsilon_{\beta,1}}{v} \right). \quad (5.10)$$

## 6. ESTIMATION OF $E(y, z, w; m)$

In this section, we seek to bound the function  $E(y, z, w; m)$  introduced in Lemma 4.1 under the assumptions in Theorem 2.1. We start with the case  $\beta = 1$ . Suppose that  $1 \leq m \leq h_0 B(x)^{1/3}$ , where  $0 < h_0 < (3/2)^{2/3}$  is any given constant. Recall that  $y \in [x^{\eta_0}, x]$ ,  $z = x^{1/v}$  and  $w = x^{1/\log(v+2)}$ . With the choice  $v = (1 - \varrho_0)^{-1} \epsilon_0^{-1} \eta_0^{-1} m$ , we clearly have  $v \in [\eta_0^{-1}, v_0 \log x / \log \log x]$  and  $m \leq (1 - \varrho_0) \epsilon_0 \log y / \log z$ . Inputting (5.6) and (5.10) into (4.3), we obtain

$$\sum_{n \leq y} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m = \lambda_\alpha C_m B(x)^{\frac{m}{2}} \left( \chi_m + O \left( \frac{m^{\frac{3}{2}}}{\sqrt{B(x)}} \right) \right) y^{\sigma_0}. \quad (6.1)$$

The key lies in the estimation of the sum

$$\sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \omega(n; z, w)^b. \quad (6.2)$$

In the present case, we may simply use the trivial bound  $\omega(n; z, w) \ll v \ll m$ , so that (6.2) is bounded above by

$$2^{O(b)} m^b \sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a.$$

It is clear that we can use (6.1) to handle the sum above. If  $a$  is even, then this sum is  $\ll \lambda_\alpha C_a B(x)^{\frac{a}{2}} y^{\sigma_0}$ ; if  $a$  is odd, then it is

$$\begin{aligned} &\leq \left( \sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^{a-1} \right)^{1/2} \left( \sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^{a+1} \right)^{1/2} \\ &\ll \lambda_\alpha \sqrt{C_{a-1} C_{a+1}} B(x)^{\frac{a}{2}} y^{\sigma_0} \end{aligned}$$

by the Cauchy-Schwarz inequality. The sequence  $\{C_\ell\}_{\ell=1}^\infty$  is strictly increasing, which can be easily seen from the identity

$$\frac{C_{\ell+1}}{C_\ell} = \frac{\ell+1}{\sqrt{2}} \cdot \frac{\Gamma(\ell/2+1)}{\Gamma((\ell+1)/2+1)} = \sqrt{2} \cdot \frac{\Gamma(\ell/2+1)}{\Gamma((\ell/2+1/2))}$$

and the fact that  $\Gamma(y)$  is strictly increasing on  $[3/2, \infty)$ . Moreover, we have by Stirling's formula that

$$\frac{C_\ell}{C_{\ell+1}} \ll \frac{1}{\ell+1} \cdot \frac{((\ell+1)/2)^{\ell/2+1} e^{-(\ell+1)/2}}{(\ell/2)^{(\ell+1)/2} e^{-\ell/2}} \ll \frac{1}{\sqrt{\ell+1}},$$

which implies that

$$C_a \leq 2^{O(m-a)} C_m \sqrt{\frac{a!}{m!}} \leq 2^{O(m-a)} C_m \sqrt{\frac{a^a}{m^m}} \leq \frac{2^{O(m-a)} C_m}{(\sqrt{m})^{m-a}}$$

for all  $0 \leq a \leq m$ . Hence, (6.2) is bounded above by

$$\frac{2^{O(m-a)} \lambda_\alpha C_m m^b}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} y^{\sigma_0} \leq 2^{O(m-a)} \lambda_\alpha C_m (\sqrt{m})^{m-a} B(x)^{\frac{a}{2}} y^{\sigma_0}.$$

Inputting this inequality into the definition of  $E(y, z, w; m)$ , we conclude that

$$E(y, z, w; m) \leq \lambda_\alpha C_m y^{\sigma_0} \sum_{a=0}^{m-1} \binom{m}{a} B(x)^{\frac{a}{2}} (O(\sqrt{m}))^{m-a} \ll \lambda_\alpha C_m m^{\frac{3}{2}} B(x)^{\frac{m-1}{2}} y^{\sigma_0}. \quad (6.3)$$

Now we consider the case  $\beta \neq 1$ . Suppose that  $1 \leq m \ll B(x)^{1/3}/(\log \log \log x)^{2/3}$  and that  $B(x)/(\log \log \log x)^2 \rightarrow \infty$  as  $x \rightarrow \infty$ . In this case we take  $v = (\log \log x)^{m(\vartheta_0+2)}$ , so that  $v \in [2\eta_0^{-1}, v_0 \log x / \log \log x]$  and  $m \leq (1 - \varrho_0)\epsilon_0 \log t / \log z$  for any  $t \in [x^{\eta_0/2}, x]$  when  $x$  is sufficiently large. Inserting (5.6) and (5.10) into (4.3) leads to

$$\sum_{n \leq t} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(n) \right)^m = \lambda_\alpha C_m B(x)^{\frac{m}{2}} \left( \chi_m + O\left( \frac{m^{\frac{3}{2}}}{\sqrt{B(x)}} \right) \right) t^{\sigma_0} (\log t)^{\beta-1} \quad (6.4)$$

uniformly for all  $t \in [x^{\eta_0/2}, x]$ . Again, we need to estimate (6.2) uniformly for  $y \in [x^{\eta_0}, x]$ . Note that (6.2) can be rewritten as

$$\sum_{k=1}^b \sum_{z < p_1 < \dots < p_k \leq w} \sum_{\substack{l_1 + \dots + l_k = b \\ l_1, \dots, l_k \geq 1}} \binom{b}{l_1, \dots, l_k} \sum_{\substack{n \leq y \\ p_1 \dots p_k | n}} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a.$$

Observe that

$$\begin{aligned} \sum_{\substack{n \leq y \\ p_1 \cdots p_k | n}} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a &= \sum_{\substack{q \leq y \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{\substack{n' \leq y/q \\ \gcd(n', q) = 1}} \alpha(n') \left| \sum_{Q_0 < p \leq z} f_p(n') \right|^a \\ &\leq \sum_{\substack{q \leq y \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a, \end{aligned}$$

since  $p_1, \dots, p_k > p$ . If  $q = p_1^{\nu_1} \cdots p_k^{\nu_k} > \sqrt{y}$  with given  $z < p_1 < \cdots < p_k \leq w$ , then we have the trivial estimate

$$\sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \leq \pi(z)^a \sum_{n \leq 3y/q} \alpha(n) \ll \lambda_\alpha \pi(z)^a \left( \frac{y}{q} \right)^{\sigma_0} \left( \log \frac{3y}{q} \right)^{\beta-1}$$

by (3.4) and the fact that  $|f_p(n)| \leq 1$ . By the proof of Lemma 3.1, and particularly by (3.3), we find that

$$\begin{aligned} \sum_{\substack{\sqrt{y} < q \leq y \\ R_q = p_1 \cdots p_k}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3y}{q} \right)^{\beta-1} &= \sum_{\substack{\sqrt{y} < p_1^{\nu_1} \cdots p_k^{\nu_k} \leq y \\ \nu_1, \dots, \nu_k \geq 1}} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_k^{\nu_k})}{p_1^{\sigma_0 \nu_1} \cdots p_k^{\sigma_0 \nu_k}} \left( \log \frac{3y}{p_1^{\nu_1} \cdots p_k^{\nu_k}} \right)^{\beta-1} \\ &\ll \frac{2^{O(k)} (\log y)^{k+\beta-2}}{(\sqrt{y})^{1-\varrho_0}}, \end{aligned}$$

from which it follows that

$$\sum_{\substack{\sqrt{y} < q \leq y \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \ll \lambda_\alpha \pi(z)^a \frac{2^{O(k)} y^{\sigma_0} (\log y)^{k+\beta-2}}{(\sqrt{y})^{1-\varrho_0}}.$$

Summing the above over all  $z < p_1 < \cdots < p_k \leq w$  yields immediately

$$\begin{aligned} &\sum_{z < p_1 < \cdots < p_k \leq w} \sum_{\substack{\sqrt{y} < q \leq y \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \\ &\leq \lambda_\alpha \pi(z)^a \pi(w)^k \frac{2^{O(k)} y^{\sigma_0} (\log y)^{k+\beta-2}}{k! (\sqrt{y})^{1-\varrho_0}} \leq \frac{\lambda_\alpha y^{\sigma_0} (\log y)^{\beta-1}}{k! (\sqrt[3]{y})^{1-\varrho_0}} \end{aligned} \quad (6.5)$$

for sufficiently large  $x$ , since  $y \in [x^{\eta_0}, x]$ ,  $a + k \leq m \ll (\log \log x)^{1/3} / (\log \log \log x)^{2/3}$ , and

$$\pi(z)^a \pi(w)^k \leq \left( \frac{w}{\log w} + O\left( \frac{w}{(\log w)^2} \right) \right)^m \ll \left( \frac{w}{\log w} \right)^m \leq \frac{x^{1/\log \log \log x} (m \log \log \log x)^m}{(\log x)^m}.$$

If  $q = p_1^{\nu_1} \cdots p_k^{\nu_k} \leq \sqrt{y}$ , then  $x^{\eta_0/2} \leq \sqrt{y} \leq y/q \leq y \leq x$ . Thus, we can apply (6.4) with  $t = y/q$  to handle

$$\sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a.$$

If  $a$  is even, then this sum is

$$\ll \lambda_\alpha C_a B(x)^{\frac{a}{2}} \left(\frac{y}{q}\right)^{\sigma_0} \left(\log \frac{y}{q}\right)^{\beta-1} \leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} \left(\frac{y}{q}\right)^{\sigma_0} \left(\log \frac{y}{q}\right)^{\beta-1};$$

if  $a$  is odd, then it is

$$\begin{aligned} &\leq \left( \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^{a-1} \right)^{1/2} \left( \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^{a+1} \right)^{1/2} \\ &\ll \lambda_\alpha \sqrt{C_{a-1} C_{a+1}} B(x)^{\frac{a}{2}} \left(\frac{y}{q}\right)^{\sigma_0} \left(\log \frac{y}{q}\right)^{\beta-1} \\ &\leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} \left(\frac{y}{q}\right)^{\sigma_0} \left(\log \frac{y}{q}\right)^{\beta-1} \end{aligned}$$

by Cauchy–Schwarz. It follows that

$$\begin{aligned} &\sum_{\substack{q \leq \sqrt{y} \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \\ &\leq \frac{2^{O(m-a)} \lambda_\alpha C_m y^{\sigma_0}}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} \sum_{\substack{q \leq \sqrt{y} \\ R_q = p_1 \cdots p_k}} \frac{\alpha(q)}{q^{\sigma_0}} \left(\log \frac{y}{q}\right)^{\beta-1} \\ &\leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1} \prod_{i=1}^k \sum_{\nu=1}^{\infty} \frac{\alpha(p_i^\nu)}{p_i^{\sigma_0 \nu}} \\ &= \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1} \prod_{i=1}^k \left( \frac{\alpha(p_i)}{p_i^{\sigma_0}} + \psi_0(p_i) \right) \end{aligned}$$

for all  $0 \leq a < m$ . Since (3.1) implies that

$$\sum_{z < p_1 < \cdots < p_k \leq w} \prod_{i=1}^k \left( \frac{\alpha(p_i)}{p_i^{\sigma_0}} + \psi_0(p_i) \right) \leq \frac{1}{k!} \left( \sum_{z < p \leq w} \left( \frac{\alpha(p)}{p^{\sigma_0}} + \psi_0(p) \right) \right)^k \leq \frac{2^{O(k)}}{k!} (\log v)^k,$$

we obtain

$$\begin{aligned} &\sum_{z < p_1 < \cdots < p_k \leq w} \sum_{\substack{q \leq \sqrt{y} \\ R_q = p_1 \cdots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \\ &\leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{k! (\sqrt{m})^{m-a}} (\log v)^k B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1}. \end{aligned} \tag{6.6}$$



Combining (6.6) with (6.5) and extending the inner sum over  $q$  to the entire range, we conclude that

$$\begin{aligned} & \sum_{z < p_1 < \dots < p_k \leq w} \sum_{\substack{q \leq y \\ R_q = p_1 \dots p_k}} \alpha(q) \sum_{n \leq y/q} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \\ & \leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{k! (\sqrt{m})^{m-a}} (\log v)^k B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1}. \end{aligned}$$

Hence, (6.2) is bounded above by

$$\begin{aligned} & \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1} \sum_{k=1}^b \frac{(\log v)^k}{k!} \sum_{\substack{l_1 + \dots + l_k = b \\ l_1, \dots, l_k \geq 1}} \binom{b}{l_1, \dots, l_k} \\ & = \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} y^{\sigma_0} (\log y)^{\beta-1} \sum_{k=1}^b \left\{ \begin{matrix} b \\ k \end{matrix} \right\} (\log v)^k \\ & \leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} T_b(\log v) y^{\sigma_0} (\log y)^{\beta-1}. \end{aligned}$$

It follows by (5.9) that the above does not exceed

$$\frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} (\log v)^b y^{\sigma_0} (\log y)^{\beta-1},$$

where we have used the observation that  $\log v > m \log \log \log x > m \geq b$ . In other words, we have shown that

$$\sum_{n \leq y} \alpha(n) \left| \sum_{Q_0 < p \leq z} f_p(n) \right|^a \omega(n; z, w)^b \leq \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} (\log v)^b y^{\sigma_0} (\log y)^{\beta-1}.$$

Inputting this inequality into the definition of  $E(y, z, w; m)$ , we conclude that

$$\begin{aligned} E(y, z, w; m) & \leq \lambda_\alpha C_m y^{\sigma_0} (\log y)^{\beta-1} \sum_{a=0}^{m-1} \binom{m}{a} B(x)^{\frac{a}{2}} \left( O \left( \frac{\log v}{\sqrt{m}} \right) \right)^{m-a} \\ & \ll \lambda_\alpha C_m \sqrt{m} (\log v) B(x)^{\frac{m-1}{2}} y^{\sigma_0} (\log y)^{\beta-1} \\ & \ll \lambda_\alpha C_m m^{\frac{3}{2}} (\log \log \log x) B(x)^{\frac{m-1}{2}} y^{\sigma_0} (\log y)^{\beta-1}. \end{aligned} \tag{6.7}$$

## 7. DEDUCTION OF THEOREMS 2.1 AND 2.2

Theorem 2.1 now follows immediately upon combining (6.1) and (6.4) with (6.3) and (6.7) and invoking Lemma 4.1 and (3.4). In fact, we have shown that the same asymptotic formulas which hold for  $M(x; m)$  also hold for

$$S(y)^{-1} \sum_{n \leq y} \alpha(n) (f(n) - A(x))^m \tag{7.1}$$

uniformly in the range  $y \in [x^{\eta_0}, x]$ , where  $\eta_0 \in (0, 1]$  is any fixed constant.

Now we prove Theorem 2.2. Recall that under the hypotheses in Theorem 2.2, the multiplicative function  $\alpha(n)$  satisfies conditions (i)–(iv). We shall again suppose  $A_0 \in (0, 1)$  throughout the proof. Define the strongly additive function  $\tilde{f}: \mathbb{N} \rightarrow \mathbb{R}$ , called the *strongly additive contraction of  $f$* , by  $\tilde{f}(p) = f(p)$  for all primes  $p$ . Then

$$\sum_{n \leq x} \alpha(n) (f(n) - A(x))^m = \sum_{k=0}^m \binom{m}{k} \sum_{n \leq x} \alpha(n) \left( \tilde{f}(n) - A(x) \right)^k \left( f(n) - \tilde{f}(n) \right)^{m-k} \quad (7.2)$$

for every  $m \in \mathbb{N}$ . The term corresponding to  $k = m$  can be estimated directly using Theorem 2.1. Hence, it remains to deal with

$$\sum_{n \leq x} \alpha(n) \left( \tilde{f}(n) - A(x) \right)^k \left( f(n) - \tilde{f}(n) \right)^l \quad (7.3)$$

for  $0 \leq k < m$  and  $l = m - k$ . Note that

$$\begin{aligned} & \left| \sum_{n \leq x} \alpha(n) \left( \tilde{f}(n) - A(x) \right)^k \left( f(n) - \tilde{f}(n) \right)^l \right| \\ & \leq \sum_{n \leq x} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k \left| \sum_{p^\nu \parallel n, \nu \geq 2} (f(p^\nu) - f(p)) \right|^l \\ & \leq \sum_{p_1, \dots, p_l \leq \sqrt{x}} \sum_{\substack{p_1^{\nu_1}, \dots, p_l^{\nu_l} \leq x \\ \nu_1, \dots, \nu_l \geq 2}} |f(p_1^{\nu_1}) - f(p_1)| \cdots |f(p_l^{\nu_l}) - f(p_l)| \sum_{\substack{n \leq x \\ p_1^{\nu_1}, \dots, p_l^{\nu_l} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k. \end{aligned}$$

Since  $f(p^\nu) = O(\nu^\kappa)$  for all  $p^\nu$ , the last expression above does not exceed

$$2^{O(l)} \sum_{s \leq l} \sum_{p_1 < \dots < p_s \leq \sqrt{x}} \sum_{\substack{l_1 + \dots + l_s = l \\ l_1, \dots, l_s \in \mathbb{N}}} \binom{l}{l_1, \dots, l_s} \sum_{\substack{p_1^{\nu_1} \dots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \sum_{\substack{n \leq x \\ p_1^{\nu_1}, \dots, p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k.$$

If we write  $n = p_1^{\nu_1} \cdots p_s^{\nu_s} n'$  with  $\gcd(n', p_1 \cdots p_s) = 1$ , then it is clear that

$$\left| \tilde{f}(n) - A(x) \right|^k = \left| \tilde{f}(n') - A(x) + \sum_{i=1}^s f(p_i) \right|^k \leq \sum_{a=0}^k \binom{k}{a} \left| \tilde{f}(n') - A(x) \right|^a \left| \sum_{i=1}^s f(p_i) \right|^{k-a}.$$

Thus, the innermost sum of  $\alpha(n) |\tilde{f}(n) - A(x)|^k$  is

$$\leq \alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s}) \sum_{a=0}^k \binom{k}{a} \left| \sum_{i=1}^s f(p_i) \right|^{k-a} \sum_{n \leq x / (p_1^{\nu_1} \cdots p_s^{\nu_s})} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^a, \quad (7.4)$$

where we have dropped the superscript of  $n$  for simplicity of notation. Since the right-hand side of the above clearly vanishes if  $p_1 \cdots p_s > \sqrt{x}$ , we may assume  $p_1 \cdots p_s \leq \sqrt{x}$  instead. Let  $\lambda' := 1 - \varrho_0 - \log_2 \lambda > \rho_0$ , and choose a constant  $\max(1/2, \sqrt{\varrho_0/\lambda'}) < \delta_0 < 1$ , so that  $1 - \varrho_0 + \delta_0^2 \lambda' > 1$ . Let  $x_s := x / (p_1 \cdots p_s)$  and  $y_s := x_s^{\delta_0}$ . Then  $x_s \geq \sqrt{x} \geq p_1 \cdots p_s$ . If

$p_1^{\nu_1} \cdots p_s^{\nu_s} > p_1 \cdots p_s y_s$  with given  $p_1 < \cdots < p_s$ , then we use the trivial estimate

$$\begin{aligned} \sum_{n \leq x / (p_1^{\nu_1} \cdots p_s^{\nu_s})} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^a &\ll 2^{O(a)} (\log x)^a \sum_{n \leq x / (p_1^{\nu_1} \cdots p_s^{\nu_s})} \alpha(n) \\ &\ll \lambda_\alpha 2^{O(a)} (\log x)^a \left( \frac{x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\sigma_0} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1}. \end{aligned}$$

Thus, (7.4) is

$$\ll \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \lambda_\alpha 2^{O(k)} x^{\sigma_0} (\log x)^k.$$

Since  $\alpha(p^\nu) = O((\lambda p^{\varrho_0 + \sigma_0 - 1})^\nu)$  for all  $p^\nu$ , we have

$$\begin{aligned} &\sum_{\substack{p_1 \cdots p_s y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \\ &\leq 2^{O(l)} \sum_{\substack{p_1 \cdots p_s y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \\ &\leq \frac{2^{O(l)}}{(p_1 \cdots p_s)^{1-\varrho_0}} \sum_{\substack{y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x_s \\ \nu_1, \dots, \nu_s \geq 1}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \left( \log \frac{3x_s}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1}. \end{aligned}$$

It is not hard to see that the proof of (3.3) also gives

$$\sum_{z_1 < p^\nu \leq z_2} \left( \frac{\lambda}{p^{1-\varrho_0}} \right)^\nu \left( \log \frac{3z_2}{p^\nu} \right)^{\beta-1} \ll \frac{(\log(3z_2/z_1))^{\beta-1}}{z_1^{1-\varrho_0-\log_p \lambda}}$$

uniformly for all primes  $p$  and all  $0 < z_1 \leq z_2$ . Thus, we have

$$\begin{aligned} &\sum_{\substack{y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x_s \\ \nu_1, \dots, \nu_s \geq 1}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \left( \log \frac{3x_s}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \\ &\leq 2^{O(l)} (\log x)^{\kappa l} \sum_{\substack{y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x_s \\ \nu_1, \dots, \nu_s \geq 1}} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \left( \log \frac{3x_s}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \\ &\leq 2^{O(l)} (\log x)^{\kappa l} \sum_{\substack{p_2^{\nu_2} \cdots p_s^{\nu_s} \leq x_s \\ \nu_2, \dots, \nu_s \geq 1}} \left( \frac{\lambda}{\log_{p_1} \lambda} \right)^{\nu_2} \cdots \left( \frac{\lambda}{p_s^{\log_{p_1} \lambda}} \right)^{\nu_s} \frac{(\log(3x_s/y_s))^{\beta-1}}{y_s^{1-\varrho_0-\log_{p_1} \lambda}} \\ &\leq \frac{2^{O(l)} (\log x)^{(\kappa+1)m+\beta-2}}{x^{\delta_0(1-\delta_0)\lambda'/2} (p_1 \cdots p_s)^{\delta_0^2 \lambda'}} \leq \frac{2^{O(l)} (\log x)^{\beta-1}}{x^{(1-\delta_0)\lambda'/5} (p_1 \cdots p_s)^{\delta_0^2 \lambda'}}, \end{aligned}$$

where the penultimate inequality follows from the previous line together with the observations that  $p_i^{\log p_i \lambda} > \lambda$  for all  $2 \leq i \leq s$ , that  $x^{(1+\delta_0)/2} \geq (p_1 \cdots p_s)^{1+\delta_0}$ , and that

$$y_s^{1-\varrho_0-\log p_1 \lambda} \geq y_s^{\lambda'} = \left( x^{(1-\delta_0)/2} \cdot \frac{x^{(1+\delta_0)/2}}{p_1 \cdots p_s} \right)^{\delta_0 \lambda'} \geq x^{\delta_0(1-\delta_0)\lambda'/2} (p_1 \cdots p_s)^{\delta_0^2 \lambda'}.$$

It follows that

$$\begin{aligned} & \sum_{\substack{p_1 \cdots p_s y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \left( \log \frac{3x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1} \\ & \leq \frac{2^{O(l)}}{(p_1 \cdots p_s)^{1-\varrho_0+\delta_0^2 \lambda'}} x^{-(1-\delta_0)\lambda'/5} (\log x)^{\beta-1}, \end{aligned}$$

from which we deduce that

$$\begin{aligned} & \sum_{p_1 < \cdots < p_s \leq \sqrt{x}} \sum_{\substack{p_1 \cdots p_s y_s < p_1^{\nu_1} \cdots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \sum_{\substack{n \leq x \\ p_1^{\nu_1}, \dots, p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k \\ & \leq 2^{O(m)} \lambda_\alpha x^{\sigma_0 - (1-\delta_0)\lambda'/5} (\log x)^{k+\beta-1} \sum_{p_1 < \cdots < p_s \leq \sqrt{x}} \frac{1}{(p_1 \cdots p_s)^{1-\varrho_0+\delta_0^2 \lambda'}} \\ & \leq \frac{1}{s!} \lambda_\alpha x^{\sigma_0 - (1-\delta_0)\lambda'/6} (\log x)^{\beta-1}. \end{aligned} \tag{7.5}$$

On the other hand, if  $p_1^{\nu_1} \cdots p_s^{\nu_s} \leq p_1 \cdots p_s y_s$ , then  $x^{(1-\delta_0)/2} \leq x/(p_1^{\nu_1} \cdots p_s^{\nu_s}) \leq x$ . Thus, we can apply the asymptotic formulas for (7.1) with  $\eta_0 = (1-\delta_0)/2$  and  $y = x/(p_1^{\nu_1} \cdots p_s^{\nu_s})$ , in conjunction with the Cauchy–Schwarz inequality, to estimate the inner sum in (7.4). As a consequence, we have

$$\sum_{n \leq x/(p_1^{\nu_1} \cdots p_s^{\nu_s})} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^a \ll \frac{2^{O(m-a)} \lambda_\alpha C_m}{(\sqrt{m})^{m-a}} B(x)^{\frac{a}{2}} \left( \frac{x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\sigma_0} \left( \log \frac{x}{p_1^{\nu_1} \cdots p_s^{\nu_s}} \right)^{\beta-1}.$$

Inserting this into (7.4) shows that the sum

$$\sum_{\substack{n \leq x \\ p_1^{\nu_1}, \dots, p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k$$

is

$$\begin{aligned} & \leq \frac{2^{O(m-k)} \lambda_\alpha C_m}{(\sqrt{m})^{m-k}} \cdot \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \left( \sqrt{B(x)} + O \left( \frac{1}{\sqrt{m}} \sum_{i=1}^s |f(p_i)| \right) \right)^k x^{\sigma_0} (\log x)^{\beta-1} \\ & = \frac{2^{O(l)} \lambda_\alpha C_m}{m^{l/2}} \cdot \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} B(x)^{\frac{k}{2}} \left( 1 + O \left( \sqrt{\frac{m}{B(x)}} \right) \right) x^{\sigma_0} (\log x)^{\beta-1} \\ & \leq \frac{2^{O(l)} \lambda_\alpha C_m}{m^{l/2}} \cdot \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1}. \end{aligned}$$

Note that

$$\begin{aligned}
& \sum_{\substack{p_1^{\nu_1} \cdots p_s^{\nu_s} \leq p_1 \cdots p_s y_s \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \\
& \leq 2^{O(l)} \sum_{\substack{p_1^{\nu_1} \cdots p_s^{\nu_s} \leq p_1 \cdots p_s y_s \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \\
& \leq \frac{2^{O(l)}}{(p_1 \cdots p_s)^{1-\varrho_0}} \sum_{\substack{p_1^{\nu_1} \cdots p_s^{\nu_s} \leq y_s \\ \nu_1, \dots, \nu_s \geq 1}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \left( \frac{\lambda}{p_1^{1-\varrho_0}} \right)^{\nu_1} \cdots \left( \frac{\lambda}{p_s^{1-\varrho_0}} \right)^{\nu_s} \\
& \leq \frac{2^{O(l)}}{(p_1 \cdots p_s)^{1-\varrho_0}} \prod_{i=1}^s \text{Li}_{-\lceil \kappa l_i \rceil} (\lambda/p_i^{1-\varrho_0}),
\end{aligned}$$

where

$$\text{Li}_{-\ell}(\zeta) := \sum_{n=1}^{\infty} n^{\ell} \zeta^n$$

is the polylogarithm function of order  $-\ell$  and complex argument  $\zeta$  with  $|\zeta| < 1$ , where  $\ell \geq 0$  is any integer. For example,  $\text{Li}_0(\zeta) = \zeta/(1-\zeta)$  and  $\text{Li}_{-1}(\zeta) = \zeta/(1-\zeta)^2$ . The function  $\text{Li}_{-\ell}(\zeta)$  can be expressed in terms of the Eulerian polynomial  $A_{\ell}(\zeta)$ :

$$\text{Li}_{-\ell}(\zeta) = \frac{\zeta A_{\ell}(\zeta)}{(1-\zeta)^{\ell+1}},$$

where

$$A_{\ell}(\zeta) := \sum_{j=0}^{\ell} \left\langle \begin{matrix} \ell \\ j \end{matrix} \right\rangle \zeta^j$$

is the  $\ell$ th Eulerian polynomial, and

$$\left\langle \begin{matrix} \ell \\ j \end{matrix} \right\rangle := \sum_{a=0}^j (-1)^a \binom{\ell+1}{a} (j+1-a)^{\ell}$$

is the  $j$ th Eulerian number of size  $\ell$ . Combinatorially, it is known that, for every  $\ell \geq 1$ ,

$$\left\langle \begin{matrix} \ell \\ j \end{matrix} \right\rangle = \#\{\tau \in S_{\ell}: \tau \text{ has exactly } j \text{ ascents}\},$$

where  $S_{\ell}$  is the set of all permutations of  $\{1, \dots, \ell\}$ . Using this combinatorial interpretation one finds that  $A_{\ell}(1) = \#S_{\ell} = \ell!$ . Since  $l_1 + \cdots + l_s = l \leq m$ , we have

$$\prod_{i=1}^s \text{Li}_{-\lceil \kappa l_i \rceil} (\lambda/p_i^{1-\varrho_0}) \leq \frac{2^{O(l)} \lceil \kappa l_1 \rceil! \cdots \lceil \kappa l_s \rceil!}{(p_1 \cdots p_s)^{1-\varrho_0}} = \frac{2^{O(l)} (l_1^{l_1} \cdots l_s^{l_s})^{\kappa}}{(p_1 \cdots p_s)^{1-\varrho_0}} \leq \frac{2^{O(l)} m^{\kappa l}}{(p_1 \cdots p_s)^{1-\varrho_0}},$$

by Stirling's formula. Hence, we obtain

$$\sum_{\substack{p_1^{\nu_1} \cdots p_s^{\nu_s} \leq p_1 \cdots p_s y_s \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \cdots \nu_s^{\kappa l_s} \frac{\alpha(p_1^{\nu_1}) \cdots \alpha(p_s^{\nu_s})}{p_1^{\sigma_0 \nu_1} \cdots p_s^{\sigma_0 \nu_s}} \leq \frac{2^{O(l)} m^{\kappa l}}{(p_1 \cdots p_s)^{2(1-\varrho_0)}}.$$

It follows that

$$\begin{aligned} & \sum_{\substack{p_1^{\nu_1} \dots p_s^{\nu_s} \leq p_1 \dots p_s y_s \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \dots \nu_s^{\kappa l_s} \sum_{\substack{n \leq x \\ p_1^{\nu_1} \dots p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k \\ & \leq \frac{2^{O(l)} \lambda_\alpha C_m m^{\kappa l}}{m^{l/2} (p_1 \dots p_s)^{2(1-\varrho_0)}} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1}. \end{aligned}$$

Summing the above over  $p_1 < \dots < p_s \leq \sqrt{x}$ , we arrive at

$$\begin{aligned} & \sum_{p_1 < \dots < p_s \leq \sqrt{x}} \sum_{\substack{p_1^{\nu_1} \dots p_s^{\nu_s} \leq p_1 \dots p_s y_s \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \dots \nu_s^{\kappa l_s} \sum_{\substack{n \leq x \\ p_1^{\nu_1} \dots p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k \\ & \leq 2^{O(l)} \lambda_\alpha C_m m^{(\kappa-1/2)l} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1} \sum_{p_1 < \dots < p_s \leq \sqrt{x}} \frac{1}{(p_1 \dots p_s)^{2(1-\varrho_0)}} \\ & \leq \frac{2^{O(l)}}{s!} \lambda_\alpha C_m m^{(\kappa-1/2)l} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1}, \end{aligned}$$

since  $\varrho_0 \in [0, 1/2)$ . Combining this estimate with (7.5), we obtain

$$\begin{aligned} & \sum_{p_1 < \dots < p_s \leq \sqrt{x}} \sum_{\substack{p_1^{\nu_1} \dots p_s^{\nu_s} \leq x \\ \nu_1, \dots, \nu_s \geq 2}} \nu_1^{\kappa l_1} \dots \nu_s^{\kappa l_s} \sum_{\substack{n \leq x \\ p_1^{\nu_1} \dots p_s^{\nu_s} \parallel n}} \alpha(n) \left| \tilde{f}(n) - A(x) \right|^k \\ & \leq \frac{2^{O(l)}}{s!} \lambda_\alpha C_m m^{(\kappa-1/2)l} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1}. \end{aligned}$$

Therefore, (7.3) is bounded above by

$$\begin{aligned} & 2^{O(l)} \lambda_\alpha C_m m^{(\kappa-1/2)l} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1} \sum_{s \leq l} \frac{1}{s!} \sum_{\substack{l_1 + \dots + l_s = l \\ l_1, \dots, l_s \in \mathbb{N}}} \binom{l}{l_1, \dots, l_s} \\ & \leq 2^{O(l)} \lambda_\alpha C_m m^{(\kappa-1/2)l} B(x)^{\frac{k}{2}} x^{\sigma_0} (\log x)^{\beta-1} T_l(1) \\ & \leq 2^{O(l)} C_m m^{(\kappa+1/2)l} B(x)^{\frac{k}{2}} S(x), \end{aligned}$$

which allows us to conclude that

$$\sum_{k=0}^{m-1} \binom{m}{k} \sum_{n \leq x} \alpha(n) \left( \tilde{f}(n) - A(x) \right)^k \left( f(n) - \tilde{f}(n) \right)^{m-k} \ll C_m m^{\kappa + \frac{3}{2}} B(x)^{\frac{m-1}{2}} S(x),$$

provided that in addition, we also have  $1 \leq m \ll B(x)^{1/(2\kappa+3)}$ . Inserting the above estimate and the estimate for the term corresponding to  $k = m$  into (7.2) completes the proof of Theorem 2.2.

## 8. PROOF OF THEOREM 2.3 (SKETCH)

Now we outline the proof of Theorem 2.3. The first step is to redefine  $f_q(n)$  introduced in Section 4. Again, let us suppose that  $A_0 \in (0, 1)$  and that  $|f(p)| \leq 1$  for all primes  $p$ . For

every  $q \in \mathbb{N}$  we define

$$\begin{aligned}\bar{F}(\sigma_0, q) &:= \prod_{p|q} (1 - F(\sigma_0, p)), \\ \tilde{F}(\sigma_0, q) &:= \frac{\rho_g(q)}{\varphi(q)} \bar{F}(\sigma_0, q).\end{aligned}$$

For each prime  $p$  we put

$$f_p(n) := \begin{cases} f(p)(1 - \tilde{F}(\sigma_0, p)), & \text{if } p \mid n, \\ -f(p)\tilde{F}(\sigma_0, p), & \text{otherwise.} \end{cases}$$

And as before, we set

$$f_q(n) := \prod_{p^\nu \parallel q} f_p(n)^\nu$$

for any  $q \in \mathbb{N}$ . In addition, let  $c_g \in \mathbb{N}$  be the least positive integer such that  $c_g g(x) \in \mathbb{Z}[x]$ , and let  $Q_0 > c_g |g(0)| \geq 1$  be such that (3.17) holds. Then for each  $q \in \mathbb{N}$  with  $P^-(q) > Q_0$  we have  $\mathcal{Z}_g(q) = \mathcal{Z}_g^*(q) \subseteq (\mathbb{Z}/q\mathbb{Z})^\times$  and  $\rho_g(q) = \#\mathcal{Z}_g^*(q)$ , where  $\mathcal{Z}_g(q)$  denotes the zero locus of  $g$  in  $\mathbb{Z}/q\mathbb{Z}$ . In particular, we have  $0 \leq \rho_g(q) \leq \varphi(q)$ , which implies that  $0 \leq \tilde{F}(\sigma_0, q) \leq 1$  and that  $|f_q(n)| \leq 1$  for all  $n \in \mathbb{N}$ .

Next, we need an analogue of Lemma 4.1. Let  $x$  be sufficiently large and set  $z := x^{\delta(x)/m} > Q_0$ . Then we have

$$\sum_{Q_0 < p \leq x} f(p) \tilde{F}(\sigma_0, p) = A_{f,g}(x) + O(1)$$

by (2.1), (3.17), and the facts that  $\rho_g$  is bounded on prime powers and that  $\sum_p \psi_0(p) < \infty$ . It is easily seen that

$$f(g(n)) - A_{f,g}(x) = \sum_{Q_0 < p \leq z} f_p(g(n)) + \sum_{\substack{p > z \\ p|g(n)}} f(p) - \sum_{z < p \leq x} f(p) \tilde{F}(\sigma_0, p) + O(1).$$

Note that

$$\sum_{z < p \leq x} f(p) \tilde{F}(\sigma_0, p) = A_{f,g}(x) - A_{f,g}(z) + O(1) \ll \log \left( \frac{m}{\delta(x)} + 1 \right).$$

Since  $1 \leq g(n) \ll n^{d_g}$  uniformly for all  $n \in \mathbb{N}$ , where  $d_g := \deg g \geq 1$ , we have

$$\sum_{\substack{p > z \\ p|g(n)}} f(p) \ll \frac{m}{\delta(x)}.$$

It follows that

$$\sum_{n \leq x} \alpha(n) (f(g(n)) - A_{f,g}(x))^m = \sum_{n \leq x} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(g(n)) \right)^m + O(E_g(x; m)),$$

where

$$E_g(x; m) := \sum_{k=0}^{m-1} \binom{m}{k} 2^{O(m-k)} (m\delta(x)^{-1})^{m-k} \sum_{n \leq x} \alpha(n) \left| \sum_{p \leq z} f_p(g(n)) \right|^k.$$

Now we turn to the estimation of

$$\sum_{n \leq x} \alpha(n) \left( \sum_{Q_0 < p \leq z} f_p(g(n)) \right)^m = \sum_{Q_0 < p_1, \dots, p_m \leq z} \sum_{n \leq x} \alpha(n) f_{p_1 \dots p_m}(g(n)).$$

Let  $q \in \mathbb{N} \cap [1, x^{\delta(x)}]$  with  $\omega(q) \leq m$ ,  $P^-(q) > Q_0$  and  $P^+(q) \leq z$ . Then we have

$$\sum_{n \leq x} \alpha(n) f_q(g(n)) = \sum_{ab|R_q} f_q(a) \mu(b) \sum_{\substack{n \leq x \\ ab|g(n)}} \alpha(n) = \sum_{ab|R_q} f_q(a) \mu(b) \sum_{c \in \mathcal{Z}_g^*(ab)} \sum_{\substack{n \leq x \\ n \equiv c \pmod{ab}}} \alpha(n).$$

Thus in place of Lemma 3.3, we need to input in our analysis the information about the distribution of values of  $\alpha(n)$  with  $n$  restricted to reduced residue classes. By Lemma 3.2, the innermost sum differs from

$$\begin{aligned} \frac{1}{\varphi(ab)} \sum_{\substack{n \leq x \\ \gcd(n, ab)=1}} \alpha(n) &= \frac{1}{\varphi(ab)} \lambda_\alpha(ab) x^{\sigma_0} (\log x)^{\beta-1} \left( 1 + O\left(\frac{1}{(\log x)^{A_0}}\right) \right) \\ &= \frac{1}{\varphi(ab)} \lambda_\alpha \bar{F}(\sigma_0, ab) x^{\sigma_0} (\log x)^{\beta-1} \left( 1 + O\left(\frac{1}{(\log x)^{A_0}}\right) \right) \\ &= \frac{1}{\varphi(ab)} \lambda_\alpha x^{\sigma_0} (\log x)^{\beta-1} \left( \bar{F}(\sigma_0, ab) + O\left(\frac{1}{(\log x)^{A_0}}\right) \right) \end{aligned}$$

by the amount  $\Delta_\alpha(x; ab, c)$ . Hence, we obtain

$$\sum_{n \leq x} \alpha(n) f_q(g(n)) = \lambda_\alpha \left( \tilde{G}_1(\sigma_0, q) + O\left(\frac{\tilde{G}_2(\sigma_0, q)}{(\log x)^{A_0}}\right) \right) x^{\sigma_0} (\log x)^{\beta-1} + J_q(x),$$

where

$$\begin{aligned} \tilde{G}_1(\sigma_0, q) &:= \sum_{ab|R_q} f_q(a) \mu(b) \tilde{F}(\sigma_0, ab), \\ \tilde{G}_2(\sigma_0, q) &:= \sum_{ab|R_q} \frac{\rho_g(ab)}{\varphi(ab)} |f_q(a)|, \\ J_q(x) &:= \sum_{ab|R_q} f_q(a) \mu(b) \sum_{c \in \mathcal{Z}_g^*(ab)} \Delta_\alpha(x; ab, c). \end{aligned}$$

It is clear that  $\tilde{G}_1$  and  $\tilde{G}_2$  are both multiplicative in  $q$ . Easy calculation shows that

$$\tilde{G}_1(\sigma_0, p^\nu) = f(p)^\nu \tilde{F}(\sigma_0, p) \left( 1 - \tilde{F}(\sigma_0, p) \right) \left( (-1)^\nu \tilde{F}(\sigma_0, p)^{\nu-1} + (1 - \tilde{F}(\sigma_0, p))^{\nu-1} \right)$$

for any prime power  $p^\nu$ . In particular, we have  $\tilde{G}_1(\sigma_0, p) = 0$ ,  $|\tilde{G}_1(\sigma_0, p^\nu)| \leq 1/4$ , and  $\tilde{G}_1(\sigma_0, p^\nu) \geq 0$  when  $2 \mid \nu$ . Moreover, we have that

$$\tilde{G}_1(\sigma_0, p^2) = f(p)^2 \tilde{F}(\sigma_0, p) \left( 1 - \tilde{F}(\sigma_0, p) \right) = \rho_g(p) \frac{f(p)^2}{p} + O\left(\frac{F(\sigma_0, p)}{p} + \frac{1}{p^2}\right),$$

and that

$$|\tilde{G}_1(\sigma_0, p^\nu)| \leq |f(p)|^\nu \tilde{F}(\sigma_0, p) \leq \rho_g(p) \frac{f(p)^2}{\varphi(p)} = \rho_g(p) \frac{f(p)^2}{p} + O\left(\frac{1}{p^2}\right)$$



for all  $p^\nu$  with  $p > Q_0$  and  $\nu \geq 2$ . In addition, the quantity

$$\left| \sum_{Q_0 < p_1, \dots, p_m \leq z} J_{p_1 \dots p_m}(x) \right|$$

is bounded above by

$$\sum_{\substack{q: \omega(q) \leq m \\ P^+(q) \leq z}} \mu(q)^2 \sum_{c \in \mathcal{Z}_g^*(q)} |\Delta_\alpha(x; q, c)| \sum_{a|q} \sum_{s \leq m} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \sum_{\substack{Q_0 < p_1 < \dots < p_s \leq z \\ q|p_1 \dots p_s}} \left| f_{p_1^{k_1} \dots p_s^{k_s}}(a) \right|.$$

The inner sum over  $s$  is

$$\begin{aligned} &\leq \sum_{s=\omega(q)}^m \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \frac{1}{(s - \omega(q))!} \left( \sum_{\substack{Q_0 < p \leq z \\ p|q}} |f_p(a)| \right)^{s - \omega(q)} \\ &\leq 2^{O(m)} \sum_{s=\omega(q)}^m \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \frac{1}{(s - \omega(q))!} (\log \log x)^{s - \omega(q)} \\ &\leq 2^{O(m)} T_m(\log \log x) \\ &\leq 2^{O(m)} (\log \log x)^m. \end{aligned}$$

It follows by (2.5) that

$$\begin{aligned} \left| \sum_{Q_0 < p_1, \dots, p_m \leq z} J_{p_1 \dots p_m}(x) \right| &\leq 2^{O(m)} (\log \log x)^m \sum_{\substack{q: \omega(q) \leq m \\ P^+(q) \leq z}} \mu(q)^2 \sum_{c \in \mathcal{Z}_g^*(q)} |\Delta_\alpha(x; q, c)| \\ &\leq 2^{O(m)} (\log \log x)^m S(x) \exp(-(\log \log x)^{1/3 + \epsilon_0}), \end{aligned}$$

which is  $o(S(x))$ . These observations allow us to conclude the proof of Theorem 2.3 by arguing as in Sections 5 and 6. It is also clear from the last inequality above that the bound  $S(x) \exp(-(\log \log x)^{1/3 + \epsilon_0})$  in (2.5) can be weakened to a complicated one involving  $\delta(x)$  and  $B_{f,g}(x)$ .

## 9. PROOFS OF THEOREM 2.4 AND COROLLARY 2.5 (SKETCH)

Now we outline the proof of Theorem 2.4, which borrows the ideas from the proofs of Theorem 2.1 and [9, Theorem 1] with proper modifications. Let  $0 < \epsilon < \min(1, K)$ , and take  $z := x^{1/v}$  and

$$w := \begin{cases} x^{1/\log(v+2)}, & \text{if } \beta = 1, \\ x^{1/(\epsilon \log(v+2))}, & \text{if } \beta \neq 1, \end{cases}$$

where we recall that  $v \asymp m$  when  $\beta = 1$  and  $v = (\log \log x)^{m(\vartheta_0+2)}$  when  $\beta \neq 1$  as chosen in Section 6. Having made these choices, we have  $\epsilon \log(v+2) \rightarrow \infty$  as  $x \rightarrow \infty$  in the case

$\beta \neq 1$ . Let

$$\begin{aligned}\mathcal{P}_\epsilon^-(x) &:= \left\{ p \leq x : |f(p)| \leq \epsilon \sqrt{B^*(x)} \right\}, \\ \mathcal{P}_\epsilon^+(x) &:= \left\{ p \leq x : \epsilon \sqrt{B^*(x)} < |f(p)| \leq K \sqrt{B^*(x)} \right\}, \\ \mathcal{P}_\infty(x) &:= \left\{ p \leq x : |f(p)| > K \sqrt{B^*(x)} \right\},\end{aligned}$$

and put  $\mathcal{P}_K(x) := \mathcal{P}_\epsilon^-(x) \cup \mathcal{P}_\epsilon^+(x)$ . We consider the strongly additive function

$$f_\epsilon(n; x) := \sum_{\substack{p|n \\ p \in \mathcal{P}_\epsilon^-(x)}} f(p) + \epsilon_{\beta,1} \sum_{\substack{p|n \\ p \in \mathcal{P}_\epsilon^+(x) \cap (z, x]}} f(p) + \sum_{\substack{p|n \\ p \in \mathcal{P}_\infty(x)}} f(p),$$

where we recall that  $\epsilon_{\beta,1}$  takes value 0 if  $\beta = 1$  and 1 otherwise, and define

$$\begin{aligned}A_\epsilon(x) &:= \sum_{p \in \mathcal{P}_\epsilon^-(x)} \alpha(p) \frac{f(p)}{p^{\sigma_0}}, \\ B_\epsilon(x) &:= \sum_{p \in \mathcal{P}_\epsilon^-(x)} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}}.\end{aligned}$$

By hypothesis,

$$B(x) - B_\epsilon(x) = \sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B^*(x)}}} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} = o(B^*(x)),$$

and so

$$|A_\epsilon(x) - A(x)| \leq \frac{1}{\epsilon \sqrt{B^*(x)}} \sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B^*(x)}}} \alpha(p) \frac{f(p)^2}{p^{\sigma_0}} = o\left(\epsilon^{-1} \sqrt{B^*(x)}\right).$$

We expect that the distribution of  $f_\epsilon(n; x)$  is close to being Gaussian with mean  $A(x)$  and variance  $B(x)$  when  $x$  gets sufficiently large. In what follows, we shall restrict our attention to the case  $\beta \neq 1$ , since the opposite case  $\beta = 1$  is not only similar but also easier. Looking back at the proof of Lemma 4.1, we find, for sufficiently large  $x$ , that

$$\sum_{p \in \mathcal{P}_\epsilon^-(x) \cap (Q_0, x]} f(p) F(\sigma_0, p) = A_\epsilon(x) + O\left(\epsilon \sqrt{B^*(x)}\right) = A(x) + O\left(\epsilon \sqrt{B^*(x)}\right),$$

so that

$$f_\epsilon(n; x) - A(x) = \sum_{p \in \mathcal{P}_\epsilon^-(x) \cap (Q_0, z]} f_p(n) + \sum_{\substack{p|n \\ p \in \mathcal{P}_K(x) \cap (z, w]}} f(p) + O\left(\epsilon \sqrt{B(x)}\right), \quad (9.1)$$

where we have used the hypothesis that  $f(n) = o(\sqrt{B(x)})$  for all  $n \leq x$  whose prime factors  $p$  satisfy  $|f(p)| > K \sqrt{B^*(x)}$ . This leads to an analogue of Lemma 4.1 in which the second sum above plays the same role as  $\omega(n; z, w)$ . To estimate the moments of  $f_\epsilon(n; x)$ , one only needs

to recycle the arguments used in the proof of Theorem 2.1 and make suitable modifications. For instance, the estimation of

$$\sum_{n \leq y} \alpha(n) \left( \sum_{p \in \mathcal{P}_\epsilon^-(x) \cap (Q_0, z]} f_p(n) \right)^m$$

is essentially the same as that of (4.1) given in Sections 4 and 5, except that we use the inequality  $|f(p)| \leq \epsilon \sqrt{B^*(x)}$  for  $p \in \mathcal{P}_\epsilon^-(x)$  in place of the bound  $f(p) = O(1)$  throughout the argument. This way, we obtain

$$\begin{aligned} \sum_{n \leq y} \alpha(n) \left( \sum_{p \in \mathcal{P}_\epsilon^-(x) \cap (Q_0, z]} f_p(n) \right)^m &= \lambda_\alpha \left( \mu_m + O \left( \frac{\epsilon \log v}{\log \log \log x} \right) \right) B(x)^{\frac{m}{2}} y^{\sigma_0} (\log y)^{\beta-1} \\ &= \lambda_\alpha (\mu_m + O(\epsilon)) B(x)^{\frac{m}{2}} y^{\sigma_0} (\log y)^{\beta-1} \end{aligned} \quad (9.2)$$

uniformly for  $y \in [x^{\eta_0}, x]$ , where  $\eta_0 \in (0, 1]$  is any given constant. On the other hand, the estimation of the error involving the second sum in (9.1) is essentially the same as that of  $E(y, z, w; m)$  in the case  $\beta \neq 1$  given in Section 6. The only difference is that we now make use of the estimates that  $|f(p)| \leq K \sqrt{B^*(x)}$  for all  $p \in \mathcal{P}_K(x)$  and that

$$\sum_{p \in \mathcal{P}_K(x) \cap (z, w]} \alpha(p) \frac{|f(p)|^\nu}{p^{\sigma_0}} \ll B^*(x)^{\frac{\nu-1}{2}} \sum_{p \in \mathcal{P}_K(x) \cap (z, w]} \alpha(p) \frac{|f(p)|}{p^{\sigma_0}} \ll \epsilon B^*(x)^{\frac{\nu}{2}} \log v \ll \epsilon B(x)^{\frac{\nu}{2}}$$

for all  $\nu \geq 1$ , which can be easily seen by considering  $p \in \mathcal{P}_\epsilon^-(x)$  and  $p \in \mathcal{P}_\epsilon^+(x)$  separately, in place of the estimates that  $f(p) = O(1)$  and that

$$\sum_{z < p \leq w} \alpha(p) \frac{|f(p)|^\nu}{p^{\sigma_0}} = O(\log v) = O(\log \log \log x),$$

respectively. One shows in this way that the error involving the second sum in (9.1) is  $O(\epsilon \lambda_\alpha B(x)^{\frac{m}{2}} y^{\sigma_0} (\log y)^{\beta-1})$ . Combining this estimate with (9.2) and taking  $y = x$  yields

$$S(x)^{-1} \sum_{n \leq x} \alpha(n) (f_\epsilon(n; x) - A(x))^m = (\mu_m + O(\epsilon)) B(x)^{\frac{m}{2}}$$

for every fixed  $m \in \mathbb{N}$  and all sufficiently large  $x$ , where the implied constant in the error term is independent of  $\epsilon$ .

To complete the proof of Theorem 2.4 for the case  $\beta \neq 1$ , it is sufficient to show

$$S(x)^{-1} \sum_{n \leq x} \alpha(n) |f(n) - f_\epsilon(n; x)|^m = O(\epsilon B^*(x)^{\frac{m}{2}}) \quad (9.3)$$

for every given  $\epsilon \in (0, 1)$  and  $m \in \mathbb{N}$ , where the implicit constant in the error term is independent of  $\epsilon$ . Since the case where  $m$  is odd follows from the case where  $m$  is even by Cauchy–Schwarz, we need only to consider the latter case. The proof of this case is largely the same as that of [9, Lemma 2], except for the slight complication in the possible case  $\beta \in (0, 1)$ . When  $m$  is even, we have

$$S(x)^{-1} \sum_{n \leq x} \alpha(n) |f(n) - f_\epsilon(n; x)|^m = S(x)^{-1} \sum_{n \leq x} \alpha(n) \sum_{\substack{p_1, \dots, p_m | n \\ p_1, \dots, p_m \in \mathcal{P}_\epsilon^+(x) \cap [2, z]}} f(p_1) \cdots f(p_m),$$

which, after grouping terms according to the distinct primes among  $p_1, \dots, p_m$ , becomes

$$S(x)^{-1} \sum_{s \leq m} \sum_{\substack{p_1 < \dots < p_s \leq z \\ p_1, \dots, p_s \in \mathcal{P}_\epsilon^+(x)}} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} f(p_1)^{k_1} \dots f(p_s)^{k_s} \sum_{\substack{n \leq x \\ p_1 \dots p_s | n}} \alpha(n). \quad (9.4)$$

By (3.4) we have

$$\sum_{\substack{n \leq x \\ p_1 \dots p_s | n}} \alpha(n) = \sum_{\substack{q \leq x \\ R_q = p_1 \dots p_s}} \alpha(q) \sum_{\substack{n' \leq x/q \\ \gcd(n', q) = 1}} \alpha(n') \ll \lambda_\alpha x^{\sigma_0} \sum_{\substack{q \leq x \\ R_q = p_1 \dots p_s}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3x}{q} \right)^{\beta-1}.$$

Appealing to (3.3) we derive

$$\begin{aligned} \sum_{\substack{q \leq x \\ R_q = p_1 \dots p_s}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3x}{q} \right)^{\beta-1} &\ll (\log x)^{\beta-1} \sum_{\substack{q \leq \sqrt{x} \\ R_q = p_1 \dots p_s}} \frac{\alpha(q)}{q^{\sigma_0}} + \sum_{\substack{\sqrt{x} < q \leq x \\ R_q = p_1 \dots p_s}} \frac{\alpha(q)}{q^{\sigma_0}} \left( \log \frac{3x}{q} \right)^{\beta-1} \\ &\ll (\log x)^{\beta-1} \prod_{i=1}^s \sum_{\nu=1}^{\infty} \frac{\alpha(p_i^\nu)}{p_i^{\sigma_0 \nu}} + \frac{(\log x)^{s+\beta-2}}{(\sqrt{x})^{1-\varrho_0}} \\ &= (\log x)^{\beta-1} \prod_{i=1}^s \left( \frac{\alpha(p_i)}{p_i^{\sigma_0}} + \psi_0(p_i) \right) + \frac{(\log x)^{s+\beta-2}}{(\sqrt{x})^{1-\varrho_0}}. \end{aligned}$$

These estimates together with (3.4) imply that (9.4) is  $\ll \Sigma_1 + \Sigma_2$ , where

$$\begin{aligned} \Sigma_1 &:= \sum_{s \leq m} \sum_{\substack{p_1 < \dots < p_s \leq z \\ p_1, \dots, p_s \in \mathcal{P}_\epsilon^+(x)}} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} |f(p_1)^{k_1} \dots f(p_s)^{k_s}| \prod_{i=1}^s \left( \frac{\alpha(p_i)}{p_i^{\sigma_0}} + \psi_0(p_i) \right), \\ \Sigma_2 &:= \frac{(\log x)^{m-1}}{(\sqrt{x})^{1-\varrho_0}} \sum_{s \leq m} \sum_{\substack{p_1 < \dots < p_s \leq z \\ p_1, \dots, p_s \in \mathcal{P}_\epsilon^+(x)}} \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} |f(p_1)^{k_1} \dots f(p_s)^{k_s}|. \end{aligned}$$

Since  $f(p) \leq K\sqrt{B^*(x)}$  for all  $p \in \mathcal{P}_\epsilon^+(x)$ , we have

$$\Sigma_2 \ll \frac{(\log x)^{m-1}}{(\sqrt{x})^{1-\varrho_0}} \pi(z)^m B^*(x)^{\frac{m}{2}} = o\left(B^*(x)^{\frac{m}{2}}\right) \ll \epsilon B^*(x)^{\frac{m}{2}}.$$

To bound  $\Sigma_1$ , we observe

$$|f(p_1)^{k_1} \dots f(p_s)^{k_s}| \ll B^*(x)^{\frac{m-s}{2}} |f(p_1) \dots f(p_s)|.$$

Thus, we have

$$\begin{aligned} \Sigma_1 &\leq \sum_{s \leq m} B^*(x)^{\frac{m-s}{2}} \frac{1}{s!} \left( \sum_{\substack{p \leq z \\ p \in \mathcal{P}_\epsilon^+(x)}} \left( \alpha(p) \frac{|f(p)|}{p^{\sigma_0}} + \psi_0(p) \right) \right)^s \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \\ &= \sum_{s \leq m} B^*(x)^{\frac{m-s}{2}} \frac{1}{s!} \left( o\left(\epsilon^{-1} \sqrt{B^*(x)}\right) \right)^s \sum_{\substack{k_1 + \dots + k_s = m \\ k_1, \dots, k_s \in \mathbb{N}}} \binom{m}{k_1, \dots, k_s} \ll \epsilon B^*(x)^{\frac{m}{2}}. \end{aligned}$$

Combining these estimates completes the proof of (9.3) in the case  $\beta \neq 1$ .

As we mentioned in Section 2, Corollary 2.5 is an immediate consequence of Theorem 2.4 when  $f$  is strongly additive. The transition to the general additive case is then accomplished by applying the following analogue of [30, Theorem B]. And this is the only place where we need to make use of characteristic functions.

**Lemma 9.1.** *Let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be an additive function, and let  $\alpha \in \mathcal{M}^*$  with parameters  $A_0, \beta, \sigma_0, \vartheta_0, \varrho_0, r$ . Denote by  $\tilde{f}$  the strongly additive contraction of  $f$ . Suppose that  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . Then  $X_N(n) := (f(n) - A(N))/\sqrt{B(N)}$  possesses a limiting distribution function with respect to the natural probability measure induced by  $\alpha$  if and only if  $\tilde{X}_N(n) := (\tilde{f}(n) - A(N))/\sqrt{B(N)}$  does, in which case they share the same limiting distribution function.*

*Proof.* As before, we shall assume  $A_0 \in (0, 1)$ . For each  $N \in \mathbb{N}$ , the distribution functions of  $X_N(n)$  and  $\tilde{X}_N(n)$  are given by

$$\begin{aligned}\Phi_N(V) &= S(N)^{-1} \sum_{\substack{n \leq N \\ X_N \leq V}} \alpha(n), \\ \tilde{\Phi}_N(V) &= S(N)^{-1} \sum_{\substack{n \leq N \\ \tilde{X}_N \leq V}} \alpha(n),\end{aligned}$$

respectively. We have to show that  $\Phi_N(V)$  converges weakly to a distribution function as  $N \rightarrow \infty$  if and only if  $\tilde{\Phi}_N(V)$  does, in which case they converge weakly to the same distribution function. Note that the characteristic functions of  $X_N(n)$  and  $\tilde{X}_N(n)$  are

$$\begin{aligned}\varphi_N(t) &= S(N)^{-1} \sum_{n \leq N} \alpha(n) e^{itX_N(n)}, \\ \tilde{\varphi}_N(t) &= S(N)^{-1} \sum_{n \leq N} \alpha(n) e^{it\tilde{X}_N(n)},\end{aligned}$$

respectively. By Lévy's continuity theorem [32, Theorem III.2.6], it suffices to show

$$\lim_{N \rightarrow \infty} (\varphi_N(t) - \tilde{\varphi}_N(t)) = 0 \tag{9.5}$$

for any given  $t \in \mathbb{R}$ . To prove this, let us fix  $t \in \mathbb{R}$  and let  $\epsilon \in (0, 1/(2|t|+1))$  be arbitrary. Denote by  $J_\epsilon(N)$  the greatest integer not exceeding  $\sqrt{N}$  such that the inequality  $|f(n)| \leq \epsilon\sqrt{B(N)}$  holds for all  $1 \leq n \leq J_\epsilon(N)$ . Since  $B(N) \nearrow \infty$  as  $N \rightarrow \infty$ , we have  $J_\epsilon(N) \nearrow \infty$

as  $N \rightarrow \infty$ . By (3.4) we have

$$\begin{aligned}
|\varphi_N(t) - \tilde{\varphi}_N(t)| &\leq S(N)^{-1} \sum_{n \leq N} \alpha(n) \left| \exp \left( it \frac{f(n) - \tilde{f}(n)}{\sqrt{B(N)}} \right) - 1 \right| \\
&= S(N)^{-1} \sum_{\substack{a \leq N \\ a \text{ squareful}}} \alpha(a) \left| \exp \left( it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right) - 1 \right| \sum_{\substack{b \leq N/a \\ b \text{ squarefree} \\ \gcd(b, a) = 1}} \alpha(b) \\
&\ll S(N)^{-1} \lambda_\alpha N^{\sigma_0} \sum_{\substack{a \leq N \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \left( \log \frac{3N}{a} \right)^{\beta-1} \left| \exp \left( it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right) - 1 \right|.
\end{aligned}$$

From (2.1) and (2.3) it follows that

$$\sum_{\substack{a=1 \\ a \text{ squareful}}}^{\infty} \frac{\alpha(a)}{a^s} = \prod_p \left( 1 + \sum_{\nu \geq 2} \frac{\alpha(p^\nu)}{p^{\nu s}} \right)$$

is absolutely convergent for  $s \in \mathbb{C}$  with  $\Re(s) > \max(\varrho_0, r) + \sigma_0 - 1$ . Thus

$$c(\delta) := \sum_{\substack{a=1 \\ a \text{ squareful}}}^{\infty} \frac{\alpha(a)}{a^{\sigma_0 - \delta}} = \prod_p \left( 1 + \sum_{\nu \geq 2} \frac{\alpha(p^\nu)}{p^{\nu(\sigma_0 - \delta)}} \right) < \infty$$

for any  $\delta < 1 - \max(\varrho_0, r)$ . Since

$$\left| it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right| \leq 2\epsilon |t| < 1$$

for all  $a \leq J_\epsilon(N)$ , this implies

$$\sum_{\substack{a \leq J_\epsilon(N) \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \left( \log \frac{3N}{a} \right)^{\beta-1} \left| \exp \left( it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right) - 1 \right| \ll \epsilon |t| (\log N)^{\beta-1}.$$

Now fix  $0 < \delta < 1 - \max(\varrho_0, r)$ . By partial summation we have

$$\sum_{\substack{a \leq x \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} = c(0) - \int_x^\infty \frac{1}{t^\delta} d \left( \sum_{\substack{a \leq t \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0 - \delta}} \right) = c(0) + o(x^{-\delta})$$

when  $x$  is sufficiently large. It follows that

$$\begin{aligned}
& \sum_{\substack{J_\epsilon(N) < a \leq N \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \left( \log \frac{3N}{a} \right)^{\beta-1} \left| \exp \left( it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right) - 1 \right| \\
& \leq 2 \sum_{\substack{J_\epsilon(N) < a \leq N \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \left( \log \frac{3N}{a} \right)^{\beta-1} \\
& = 2 \int_{J_\epsilon(N)}^N \left( \log \frac{3N}{t} \right)^{\beta-1} d \left( \sum_{\substack{a \leq t \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \right) \\
& = o(N^{-\delta}) + o((\log N)^{\beta-1} J_\epsilon(N)^{-\delta}) + o \left( \int_{J_\epsilon(N)}^N t^{-1-\delta} \left( \log \frac{3N}{t} \right)^{\beta-2} dt \right)
\end{aligned}$$

for sufficiently large  $N$ . By a change of variable we see that

$$\begin{aligned}
\int_{J_\epsilon(N)}^N t^{-1-\delta} \left( \log \frac{3N}{t} \right)^{\beta-2} dt &= (3N)^{-\delta} \int_{\log 3}^{\log(3N/J_\epsilon(N))} e^{\delta t} t^{\beta-2} dt \\
&\ll (3N)^{-\delta} \left( \frac{3N}{J_\epsilon(N)} \right)^\delta \left( \log \frac{3N}{J_\epsilon(N)} \right)^{\beta-2} \\
&\ll (\log N)^{\beta-2} J_\epsilon(N)^{-\delta}.
\end{aligned}$$

Hence, we have

$$\sum_{\substack{J_\epsilon(N) < a \leq N \\ a \text{ squareful}}} \frac{\alpha(a)}{a^{\sigma_0}} \left( \log \frac{3N}{a} \right)^{\beta-1} \left| \exp \left( it \frac{f(a) - f(R_a)}{\sqrt{B(N)}} \right) - 1 \right| = o((\log N)^{\beta-1} J_\epsilon(N)^{-\delta}).$$

for sufficiently large  $N$ . Gathering the estimates above, we obtain

$$\varphi_N(t) - \tilde{\varphi}_N(t) \ll \epsilon |t| + o(J_\epsilon(N)^{-\delta})$$

for sufficiently large  $N$ , where the implicit constants are independent of  $t$ ,  $\epsilon$  and  $N$ . From this estimate we infer that

$$\limsup_{N \rightarrow \infty} |\varphi_N(t) - \tilde{\varphi}_N(t)| = O(\epsilon |t|),$$

where the implicit constant is independent of  $t$  and  $\epsilon$ . Since  $\epsilon \in (0, 1/(2|t|+1))$  is arbitrary, we obtain (9.5) as desired.  $\square$

*Remark 9.1.* Let  $\alpha(n) = \tau(n)^2/n^{11}$ , where  $\tau$  is Ramanujan's  $\tau$ -function, and define the additive function  $f(n)$  by  $f(p^\nu) = \log \sqrt{\alpha(p^\nu)}$  if  $\alpha(p^\nu) \neq 0$  and  $f(p^\nu) = 0$  otherwise, where  $p^\nu$  is any prime power. Then  $\alpha(n)$  satisfies conditions (i)–(iv) with any fixed  $A_0 > 0$ ,  $\beta = 1$ ,  $\sigma_0 = 1$ ,  $\vartheta_0 = 0$ , and any fixed  $\varrho_0 \in (0, 1)$  and  $r \in (1/2, 1)$ . Moreover, we have  $\alpha(n) \leq d(n)^2$  by Deligne's bound [10]. As alluded to in Section 2, Elliott [13] showed, using ideas from probability theory, that the limiting distribution of  $(f(n) - A(x))/\sqrt{B(x)}$  with respect to the natural probability measure induced by  $\alpha$  is the standard Gaussian distribution. In fact,

we can derive his result from Corollary 2.5 in combination with Lemma 9.1 and [13, Lemma 7] without difficulty. In comparison to Elliott's probabilistic approach, our approach enables us to get around some of the complications resulting from the analysis of  $\tau(n)$ .

To illustrate this, let us consider the strongly additive function  $f_0(n)$  defined by  $f_0(p) = \log \sqrt{\alpha(p)}$  if  $p \notin E_0$  and  $f_0(p) = 0$  otherwise, where  $E_0 := \{p > 2: \alpha(p) \leq \exp(-2\sqrt[3]{\log \log p})\}$ . Denote by  $A_0(x)$  and  $B_0(x)$  the expected mean and variance of  $f_0(n)$  weighted by  $\alpha(n)$ , respectively. It can be shown [13, Lemma 7] that  $B(x) \asymp \log \log x$ . Since the inequality  $t|\log t| \leq \sqrt{t}$  holds for all  $t \in [0, 1]$ , we have

$$\sum_{\substack{p \leq x \\ p \in E_0}} \alpha(p) \frac{|f(p)|}{p} \leq \sum_{\substack{p \leq x \\ p \in E_0}} \frac{\sqrt{\alpha(p)}}{p} \leq \sum_{p > 2} \frac{1}{p} \exp\left(-\sqrt[3]{\log \log p}\right) < \infty.$$

It follows that  $A_0(x) = A(x) + O(1)$ . A similar argument shows that  $B_0(x) = B(x) + O(1) \asymp \log \log x$ . Thus,  $f_0(p) = O(B_0(p)^{1/3})$  for all  $p$ , which shows that  $f_0(n)$  satisfies the hypotheses in Corollary 2.5. Hence, the limiting distribution of  $(f_0(n) - A(x))/\sqrt{B(x)}$  with respect to the natural probability measure induced by  $\alpha$  is the standard Gaussian distribution.

To complete our argument, let  $\tilde{f}$  be the strongly additive contraction of  $f$ . Then  $f_0(n) \geq \tilde{f}(n)$  for all  $n \in \mathbb{N}$ . Moreover, Deligne's bound and the fact that  $\tau(n) \in \mathbb{Z}$  for all  $n \in \mathbb{N}$  imply that  $-(11 \log p)/2 \leq f(p) \leq \log 2$  whenever  $\alpha(p) \neq 0$ . Since

$$\sum_{\nu \geq 1} \frac{\alpha(p^\nu)}{p^\nu} \leq \frac{\alpha(p)}{p} + \sum_{\nu \geq 2} \frac{(\nu + 1)^2}{p^\nu} = \frac{\alpha(p)}{p} + O\left(\frac{1}{p^2}\right),$$

we have

$$\begin{aligned} S(x)^{-1} \sum_{n \leq x} \alpha(n) \left(f_0(n) - \tilde{f}(n)\right) &= S(x)^{-1} \sum_{\substack{p \leq x \\ p \in E_0}} |f(p)| \sum_{\substack{n \leq x \\ p|n}} \alpha(n) \\ &= S(x)^{-1} \sum_{\substack{p \leq x \\ p \in E_0}} |f(p)| \sum_{\nu \geq 1} \alpha(p^\nu) \sum_{\substack{n' \leq x/p^\nu \\ p \nmid n'}} \alpha(n') \\ &\ll S(x)^{-1} \lambda_\alpha x \sum_{\substack{p \leq x \\ p \in E_0}} |f(p)| \sum_{\nu \geq 1} \frac{\alpha(p^\nu)}{p^\nu} \\ &\ll \sum_{\substack{p \leq x \\ p \in E_0}} \alpha(p) \frac{|f(p)|}{p} + O\left(\sum_{p > 2} \frac{\log p}{p^2}\right) \ll 1. \end{aligned}$$

This estimate is sufficient for us to conclude that the limiting distribution of  $(\tilde{f}(n) - A(x))/\sqrt{B(x)}$  with respect to the natural probability measure induced by  $\alpha$  is also the standard Gaussian distribution. By Lemma 9.1, the same is true for  $(f(n) - A(x))/\sqrt{B(x)}$ .



## 10. CONCLUDING REMARKS

Although in the present paper we only focused on the subclass  $\mathcal{M}^*$  of multiplicative functions, it is also of interest to consider weight functions  $\alpha(n)$  which satisfy certain Landau–Selberg–Delange type conditions. Given more information about  $\alpha(n)$  and its associated Dirichlet series  $F(s) = \sum_{n=1}^{\infty} \alpha(n)n^{-s}$ , better results are obtainable in some circumstances. Below we give a brief description of the method in the special case where  $F(s)$  is close to an integral power of the Riemann zeta-function  $\zeta(s)$ .

For a complex number  $s \in \mathbb{C}$ , we write  $\sigma = \Re(s)$  and  $t = \Im(s)$ . Let  $\alpha: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$  be a multiplicative function whose Dirichlet series  $F(s) = \sum_{n=1}^{\infty} \alpha(n)n^{-s}$  is absolutely convergent for  $s \in \mathbb{C}$  with  $\sigma > \sigma_0$ , where  $\sigma_0 > 0$  is constant. Suppose that there exist constants  $\beta \in \mathbb{N}$ ,  $0 < \theta_0 < \sigma_0$ ,  $B > 0$ , and  $0 < \delta < 1$ , such that  $H_{\beta}(s) := F(s)\zeta(s - \sigma_0 + 1)^{-\beta}$  has an analytic continuation in the half plane  $\sigma \geq \theta_0$  with

$$\lim_{s \rightarrow \sigma_0} F(s)(s - \sigma_0)^{\beta} > 0,$$

and such that  $|H_{\beta}(s)| \leq B(1 + |t|)^{1-\delta}$  for all  $s \in \mathbb{C}$  with  $\sigma \geq \theta_0$ . It is clear that  $F(s)$  has (absolute) abscissa of convergence  $\sigma_0$ . Adapting the argument used in the proof of [25, Lemma 2.1] or [32, Theorem II.5.2], one can show that there exists some constant  $\epsilon_0 > 0$  such that

$$S(x) = \frac{1}{\sigma_0} \operatorname{Res}_{s=\sigma_0} \left( \frac{F(s)x^s}{s - \sigma_0 + 1} \right) - x^{\sigma_0} (\log x)^{\beta-1} \sum_{k=1}^{\beta-1} \sum_{j=0}^{k-1} c_{j,k} \frac{\mu_j(\beta)}{(\log x)^k} + O(Bx^{\theta}) \quad (10.1)$$

uniformly for all  $x \geq 3$  and  $\theta \in (\sigma_0 - \epsilon_0, \sigma_0)$ , where

$$\begin{aligned} \mu_k(\beta) &:= \frac{1}{k!} \cdot \frac{d^k}{ds^k} \left( \frac{F(s)(s - \sigma_0)^{\beta}}{s - \sigma_0 + 1} \right) \Big|_{s=\sigma_0}, \\ c_{j,k} &:= \frac{(-1)^{k-j}(\sigma_0 - 1)}{(\beta - k - 1)! \sigma_0^{k-j+1}}, \end{aligned}$$

and the implicit constant in the error term depends at most on  $\beta, \sigma_0, \theta_0, \delta, \epsilon_0$ . Notably, one gains an asymptotic for  $S(x)$  with a power-saving error term uniformly in  $B$ , in contrast to what is provided by (3.4). Furthermore, suppose that there exists a constant  $\lambda > 0$  such that  $\alpha(p^{\nu}) = O((\lambda p^{\sigma_0-1})^{\nu})$  for all prime powers  $p^{\nu}$ . Let

$$F(s, a) := \prod_{p|a} \left( 1 - \left( \sum_{\nu=0}^{\infty} \alpha(p^{\nu}) p^{-\nu s} \right)^{-1} \right)$$

for  $s \in \mathbb{C}$  with  $\sigma \geq \theta_0$  and squarefree  $a \in \mathbb{N}$ . When  $s = \sigma_0$ , this definition coincides with the one introduced in Lemma 3.3. As in the proof of Lemma 3.3, it is not hard to show that

$$F(s, p) = \frac{\alpha(p)}{p^s} + O \left( \frac{\alpha(p)^2}{p^{2\sigma}} + \frac{1}{p^{2(\sigma-\sigma_0+1)}} \right) \quad (10.2)$$

for all  $s \in \mathbb{C}$  with  $\sigma \geq \theta_0$  and all sufficiently large  $p$ . In addition, we observe that

$$\begin{aligned} \sum_{\substack{n=1 \\ a|n}}^{\infty} \frac{\alpha(n)}{n^s} &= \left( \prod_{p|a} \sum_{\nu=0}^{\infty} \alpha(p^\nu) p^{-\nu s} \right) \left( \prod_{p|a} \sum_{\nu=1}^{\infty} \alpha(p^\nu) p^{-\nu s} \right) \\ &= F(s) \left( \prod_{p|a} \sum_{\nu=0}^{\infty} \alpha(p^\nu) p^{-\nu s} \right)^{-1} \left( \prod_{p|a} \sum_{\nu=1}^{\infty} \alpha(p^\nu) p^{-\nu s} \right) = F(s) F(s, a) \end{aligned}$$

for  $s \in \mathbb{C}$  with  $\sigma > \sigma_0$  and squarefree  $a \in \mathbb{N}$ . Applying (10.1) to the above Dirichlet series expansion of  $F(s)F(s, a)$  and using (10.2) to obtain upper bounds for  $H_\beta(s)F(s, a)$  uniformly in  $\sigma \geq \theta_0$ , we see that there exist constants  $\epsilon \in (0, 1)$ ,  $Q_0 \geq 2$ , and  $d_{j,k} \in \mathbb{R}$ , where  $0 \leq j < k < \beta$ , such that

$$\begin{aligned} \sum_{\substack{n \leq x \\ a|n}} \alpha(n) &= \frac{\mu_0(\beta) F(\sigma_0, a)}{(\beta - 1)! \sigma_0} x^{\sigma_0} (\log x)^{\beta-1} + x^{\sigma_0} (\log x)^{\beta-1} \sum_{k=1}^{\beta-1} \sum_{j=0}^k d_{j,k} \frac{F^{(j)}(\sigma_0, a)}{(\log x)^k} \\ &\quad + O(B 2^{O(\omega(a))} a^{\sigma_0-1} (x/a)^\theta) \end{aligned} \tag{10.3}$$

uniformly for all  $x \geq 3$ ,  $\theta \in (\sigma_0 - \epsilon, \sigma_0)$  and square-free  $a \in \mathbb{N}$  with  $P^-(a) > Q_0$ , where  $F^{(j)}(\sigma_0, a)$  is the  $j$ th order derivative of  $F(s, a)$  with respect to  $s$  evaluated at  $s = \sigma_0$ . Again, one may compare this result with Lemma 3.3.

Now, if  $f: \mathbb{N} \rightarrow \mathbb{R}$  is a strongly additive function with  $|f(p)| \leq M$  for all primes  $p$ , where  $M > 0$  is constant, and if  $0 < h_0 < (3/2)^{2/3}$  is fixed but arbitrary, then we obtain, by using (10.3) as a substitute for Lemma 3.3 and arguing as before with the adoption of the technique used in [25, Section 4.2], that

$$M(x; m) = C_m B(x)^{\frac{m}{2}} \left( \chi_m + O\left(\frac{m^{\frac{3}{2}}}{\sqrt{B(x)}}\right) \right)$$

uniformly for all sufficiently large  $x$  and all  $1 \leq m \leq h_0(B(x)/M^2)^{1/3}$ , provided that  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . Analogously, let  $f: \mathbb{N} \rightarrow \mathbb{R}$  is strongly additive such that  $f(p) = O(\sqrt{B(p)})$  for all primes  $p$ ,  $B(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , and

$$\sum_{\substack{p \leq x \\ |f(p)| > \epsilon \sqrt{B(x)}}} \alpha(p) \frac{f(p)^2}{p} = o(B(x))$$

for any given  $\epsilon > 0$ . Then  $M(x; m) = (\mu_m + o(1)) B(x)^{\frac{m}{2}}$  for every fixed  $m \in \mathbb{N}$ . These results supplement Theorems 2.1 and 2.4. It may be worth pointing out that in the proofs of these results one can simply take  $z = x^{1/v}$  with  $v$  being a suitable constant multiple of  $m$ . We invite the reader to fill in the details.

One of the key ingredients in the proof of Theorem 2.1 is an asymptotic formula for

$$\sum_{\substack{n \leq x \\ d|n}} \alpha(n),$$

which is provided by Lemma 3.3. More generally, let  $\mathcal{A}(x) = \{a_n\}_{n \leq x}$  be a non-decreasing sequence of positive integers, and suppose that

$$\mathcal{A}_{d,\alpha}(x) := \sum_{\substack{n \leq x \\ d|a_n}} \alpha(n) = \rho(d)S(x) + r_d(x) \quad (10.4)$$

for square-free integers  $d \in \mathbb{N}$ , where  $\rho: \mathbb{N} \rightarrow [0, 1]$  is a multiplicative function, and  $r_d(x)$  is a remainder term which is expected to be small for all  $d$  or small on average over  $d$ . Here,  $\rho(d)$  can be viewed as the density of the set  $\{n \in \mathbb{N}: d \mid a_n\}$  with respect to the probability measure induced by  $\alpha$ . In this sieve-theoretic setting one can derive, without much difficulty, an analogue of [18, Proposition 4]. It may be of interest to determine if such an analogue can be used to obtain general weighted Erdős–Kac theorems for various interesting sequences  $\{a_n\}$  studied relatively recently, including  $g(p_n)$ ,  $\varphi(n)$ , the Carmichael function  $\lambda(n)$ , and the aliquot sum  $s(n) := \sigma(n) - n$ , where  $g \in \mathbb{Z}[x]$  is an irreducible polynomial,  $p_n$  is the  $n$ th prime, and  $\lambda(n)$  denotes the exponent of  $(\mathbb{Z}/n\mathbb{Z})^\times$  (see [21], [15, 17] and [28]). Besides, the same approach may also be adapted to prove results of weighted Erdős–Kac type for short intervals as well as in the function field setting. We hope to return to these and other related problems in the future.

**Acknowledgment.** The author thanks his advisor Carl Pomerance for stimulating discussions and helpful comments. He is also grateful to Prof. Paul Pollack for his cheerful encouragement and valuable feedback. Finally, he would like to express his gratitude to the anonymous referee for a careful reading of the manuscript and for detailed comments and suggestions which helped improve the paper considerably.

## REFERENCES

- [1] K. Alladi, *The distribution of  $\nu(n)$  in the sieve of Eratosthenes*, Quart. J. Math. Oxford Ser. (2) **33** (1982), 129–148. MR657120
- [2] ———, *An Erdős–Kac theorem for integers without large prime factors*, Acta Arith. **49** (1987), 81–105. MR913766
- [3] P. Billingsley, *On the central limit theorem for the prime divisor functions*, Amer. Math. Monthly **76** (1969), 132–139. MR0242222
- [4] ———, *Probability and measure*, 3rd. ed., Wiley Ser. Probab. Stat., Wiley-Intersci. Publ., John Wiley & Sons, Inc., New York, 1995. MR1324786
- [5] E. Bombieri, J. B. Friedlander, and H. Iwaniec, *Primes in arithmetic progressions to large moduli*, Acta Arith. **156** (1986), no. 3–4, 203–251. MR0834613
- [6] R. de la Bretèche and G. Tenenbaum, *Remarks on the Selberg–Delange method*, Acta Arith. **200** (2021), no. 4, 349–369. MR4331908
- [7] H. Delange, *Sur le nombre des diviseurs premiers de  $n$* , C. R. Acad. Sci. (Paris) **237** (1953), 542–544. MR0057282
- [8] ———, *Sur un théorème d’Erdős et Kac*, Acad. Roy. Belg. Bull. Cl. Sci. (5) **42** (1956), 130–144. MR0076800
- [9] H. Delange and H. Halberstam, *A note on additive functions*, Pacific J. Math. **7** (1957), 1551–1556. MR0092820
- [10] P. Deligne, *La conjecture de Weil. I.*, Inst. Hautes Études Sci. Publ. Math. **43** (1974), 273–307. MR0340258
- [11] R. Durrett, *Probability: theory and examples*, Camb. Ser. Stat. Probab. Math., vol. 49, Cambridge University Press, Cambridge, 2019. MR3930614
- [12] D. Elboim and O. Gorodetsky, *Multiplicative arithmetic functions and the generalized Ewens measure*, preprint (2022), to appear in Israel J. Math., arXiv:1909.00601.

- [13] P. D. T. A. Elliott, *A central limit theorem for Ramanujan's tau function*, Ramanujan J. **29** (2012), no. 1–3, 145–161. MR2994095
- [14] ———, *Corrigendum: central limit theorems for classical cusp forms*, Ramanujan J. **36** (2015), no. 1–2, 99–102. MR3296712
- [15] P. Erdős, A. Granville, C. Pomerance, and C. Spiro, *On the normal behavior of the iterates of some arithmetic functions*, Analytic Number Theory, Proc. Conf. in honor of Paul T. Bateman, B. C. Berndt et al. (eds.), Progr. Math., vol. 85, Birkhäuser, Boston, 1990. MR1084181
- [16] P. Erdős and M. Kac, *The Gaussian law of errors in the theory of additive number theoretic functions*, Amer. J. Math. **62** (1940), 738–742. MR0002374
- [17] P. Erdős and C. Pomerance, *On the normal number of prime factors of  $\varphi(n)$* , Rocky Mountain J. Math. **15** (1985), no. 2, 343–352. MR0823246
- [18] A. Granville and K. Soundararajan, *Sieving and the Erdős–Kac theorem*, Equidistribution in Number Theory, An Introduction, A. Granville and Z. Rudnick (eds.), NATO Sci. Ser. II Math. Phys. Chem., vol. 237, Springer, Dordrecht, 2007. MR2290492
- [19] H. Halberstam, *On the distribution of additive number-theoretic functions*, J. London Math. Soc. **30** (1955), 43–53. MR0066406
- [20] ———, *On the distribution of additive number-theoretic functions (II)*, J. London Math. Soc. **31** (1956), 1–14. MR0073626
- [21] ———, *On the distribution of additive number-theoretic functions (III)*, J. London Math. Soc. **31** (1956), 14–27. MR0073627
- [22] G. H. Hardy and S. Ramanujan, *The normal number of prime factors of a number  $n$* , Quart. J. Pure. Appl. Math. **48** (1917), 76–92. MR2280878
- [23] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 6th. ed., Revised by D. R. Heath-Brown and J. H. Silverman; With a forward by A. J. Wiles, Oxford Univ. Press, Oxford, 2008. MR2445243
- [24] M. Kac, *Probability methods in some problems of analysis and number theory*, Bull. Amer. Math. Soc. **55** (1949), 641–665. MR0031504
- [25] R. Khan, M. B. Milinovich, and U. Subedi, *A weighted version of the Erdős–Kac theorem*, J. Number Theory **239** (2022), 1–20. MR4434484
- [26] Wm. J. LeVeque, *On the size of certain number-theoretic functions*, Trans. Amer. Math. Soc. **66** (1949), 440–463. MR0030993
- [27] D. T. Nguyen, *Generalized divisor functions in arithmetic progressions: I*, J. Number Theory **227** (2021), 30–93. MR4254563
- [28] P. Pollack and L. Troupe, *Sums of proper divisors follow the Erdős–Kac law*, Proc. Amer. Math. Soc. **151** (2023), no. 3, 977–988. MR4531632
- [29] A. Rényi and P. Turán, *On a theorem of Erdős–Kac*, Acta Arith. **4** (1958), 71–84. MR0096629
- [30] H. N. Shapiro, *Distribution functions of additive arithmetic functions*, Proc. Nat. Acad. Sci. U.S.A. **42** (1956), 426–430. MR0079609
- [31] J. M. Song, *Sums of nonnegative multiplicative functions over integers without large prime factors I*, Acta Arith. **97** (2001), no. 4, 329–351. MR1823551
- [32] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, 3rd. ed., Grad. Stud. Math., vol. 163, Amer. Math. Soc., Providence, RI., 2015. MR3363366
- [33] ———, *Moyennes effectives de fonctions multiplicatives complexes*, Ramanujan J. **44** (2017), no. 3, 641–701. MR3723446
- [34] ———, *Correction to: moyennes effectives de fonctions multiplicatives complexes*, Ramanujan J. **53** (2020), no. 1, 243–244. MR4148467
- [35] P. Turán, *On a theorem of Hardy and Ramanujan*, Ramanujan J. **9** (1934), no. 4, 274–276. MR1574877

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA

Email address: `steve.fan.gr@dartmouth.edu`