

PHOENIX Evaluation Study

Phase 1 — Expert Generation Survey *Healthcare Professional (PRE) Instrument*

Study title	Evaluating the Clinical Quality of an Ontology-Based Multi-Agent Mental Health Support System (PHOENIX)
Institution	Ghent University — Faculty of Psychology and Educational Sciences
Researcher	Stijn Van Severen (M.Sc. candidate)
Supervisors	Prof. Geert Crombez; Dr. Annick De Paepe
Phase	Phase 1 — Generation (PRE)
Estimated time	Approximately 60 minutes (can be completed across multiple sessions)
Contact	stijn.vanseveren@ugent.be

Thank you for participating.

You are taking part in a master's thesis evaluation study at Ghent University. The study examines a novel multi-agent artificial intelligence system — called **PHOENIX** — designed to support personalized digital mental health interventions. The system processes free-text complaints submitted by individuals experiencing mental health difficulties and works through a structured five-stage clinical reasoning process.

This survey asks you to independently perform the **same five clinical reasoning tasks** on ten standardised case vignettes. Your responses will serve as the expert reference corpus in a subsequent double-blind evaluation (Phase 2) where independent healthcare professionals compare outputs from multiple sources without knowing their origin.

There are no right or wrong answers. We are interested in your genuine clinical expertise and reasoning. Please respond as you would in an actual clinical context.

This document is the survey blueprint for Qualtrics configuration.
It contains the exact question text, case vignettes, standardised context, and response format specifications for all five parts of the Phase 1 survey.

April 14, 2026

Contents

1	Introduction and Participant Information	2
1.1	Background: The PHOENIX System	2
1.2	Your Role in This Study	2
1.3	Important Instructions	3
1.4	Ethical Notice	3
1.5	Estimated Session Length	3
2	Case Assignment Protocol	4
3	Case Vignettes Reference (All Parts)	5
4	Part 1: Operationalization of Mental State	7
5	Part 2: Initial Observational Model Construction	12
6	Part 3: Treatment-Target Identification	18
7	Part 4: Updated Observation Model	24
8	Part 5: Tailored Intervention Message	28
9	Completion and Submission	32
10	Qualtrics Configuration Notes	32

1 Introduction and Participant Information

1.1 Background: The PHOENIX System

PHOENIX (**P**ersonalized **H**ierarchical **O**ptimization **E**ngine for **N**avigating **I**nsightful **eX**plorations) is a multi-agent AI pipeline developed as part of a master’s thesis in psychology at the Faculty of Psychology and Educational Sciences (FPPW) of Ghent University. The system is designed to support personalized longitudinal mental health interventions by processing a free-text complaint and producing a structured five-step clinical reasoning output.

The five stages of the PHOENIX pipeline mirror the core clinical reasoning tasks involved in formulation-driven digital intervention design:

Step	Clinical Task	System Output
01	Operationalization of Mental State	List of 2–6 operationalized criterion variables derived from the complaint
02	Initial Observational Model	Bipartite network linking criteria to modifiable predictor variables
03	Treatment-Target Identification	Ranked shortlist of 2–4 modifiable treatment targets
04	Updated Observational Model	Refined measurement plan for the next monitoring cycle
05	Tailored Intervention Message	HAPA phase classification + 2–3 sentence coaching message

1.2 Your Role in This Study

As a healthcare professional with relevant clinical expertise, you are asked to perform the same five clinical reasoning tasks independently and without reference to AI outputs. You will be assigned exactly two case vignettes from the pool of ten (see Section 2). All five parts of this survey are completed for your two assigned cases only. Specifically, for each of your two assigned case vignettes:

- **Part 1:** Identify and operationalize the key mental health criteria present in the complaint.
- **Part 2:** Propose a set of modifiable predictor variables that could be monitored over time.
- **Part 3:** Identify and prioritize the most appropriate treatment targets.
- **Part 4:** Refine the measurement plan for the next monitoring cycle.
- **Part 5:** Write a brief, personalized digital coaching message.

Your responses will be compared to PHOENIX outputs in a double-blind evaluation. The comparing evaluators will not know which outputs came from a human expert and which from the AI system.

1.3 Important Instructions

Please read before beginning

1. **Work independently.** Do not consult colleagues, clinical guidelines, or other resources while completing this survey. We are interested in your personal clinical knowledge and judgment.
2. **Use the standardised context.** Parts 2–5 provide you with context from previous steps. Please base your responses on this standardised context rather than generating your own prior-step outputs. This ensures all participants work from identical inputs.
3. **Be concise but specific.** There is no required minimum length. Short, precise responses are preferred over lengthy ones. Aim for clinical clarity.
4. **Use the case profiles.** Each vignette indicates the person's approximate age, occupation, and complaint duration. Use this background as you would in a brief referral letter.
5. **Work through the parts in order.** Parts are designed to build sequentially. You may complete the survey across multiple sessions but please complete each part in a single sitting if possible.
6. **If a case does not fit your primary discipline,** use your best clinical judgment based on your broader mental health knowledge.

1.4 Ethical Notice

Participation is voluntary. Your responses will be pseudonymized before analysis and will not be shared outside the research team. The data will be used exclusively for the purpose of this thesis evaluation study. You may withdraw at any time by contacting the researcher. By proceeding, you confirm that you have read this information sheet and consent to participate.

1.5 Estimated Session Length

Part	Task	Response format	Est. time
1	Operationalization	Structured text (label + description)	≈6 min
2	Initial Obs. Model	Structured text (predictor + edges)	≈6 min
3	Treatment Targets	Rank-order with bipartite visual	≈7 min
4	Updated Obs. Model	Checkbox selection + optional note	≈5 min
5	Intervention Message	HAPA phase + 2–3 sentence message	≈8 min
Total (2 cases)		10 structured items (2 cases × 5 parts)	≈30–40 min

2 Case Assignment Protocol

Important: You only complete your two assigned cases.

This survey uses a **distributed assignment design**. The ten case vignettes are divided non-overlappingly across five healthcare professionals. You are assigned exactly two cases. This ensures all ten cases receive genuine independent expert responses, while keeping your total time to approximately 30–40 minutes.

Each participant's two responses are later pooled with the other HCPs' responses to form a corpus of ten unique expert outputs — one per case — which are then used in Phase 2 (POST evaluation).

HCP-to-Case Assignment Table

Participant Code	Assigned Cases	Case Profiles
HCP-PRE-01	C01 + C02	28-yr software developer (sleep/stress) 34-yr admin (panic/avoidance)
HCP-PRE-02	C03 + C04	41-yr teacher (burnout/withdrawal) 63-yr retiree (grief/rumination)
HCP-PRE-03	C05 + C06	25-yr student (OCD-type) 32-yr marketing (performance anxiety)
HCP-PRE-04	C07 + C08	46-yr manager (depression) 38-yr project manager (ADHD)
HCP-PRE-05	C09 + C10	27-yr sales rep (BPD-type) 52-yr director (stress/alcohol coping)

Qualtrics Display Logic Implementation

In the Qualtrics survey, a screening question at the beginning asks participants to select their assigned participant code from a dropdown (single-choice: HCP-PRE-01 through HCP-PRE-05). Each case block throughout all five parts is then set to **Display Logic: Show this block if Q_ParticipantCode = [relevant code]**.

Qualtrics Block	Display Condition	Cases Shown
Block A	Q_ParticipantCode = HCP-PRE-01	C01 (all 5 parts) + C02 (all 5 parts)
Block B	Q_ParticipantCode = HCP-PRE-02	C03 (all 5 parts) + C04 (all 5 parts)
Block C	Q_ParticipantCode = HCP-PRE-03	C05 (all 5 parts) + C06 (all 5 parts)
Block D	Q_ParticipantCode = HCP-PRE-04	C07 (all 5 parts) + C08 (all 5 parts)
Block E	Q_ParticipantCode = HCP-PRE-05	C09 (all 5 parts) + C10 (all 5 parts)

Note for Qualtrics configurator

For each of the five survey parts (Sections 4–8 of this document), the case items are grouped in pairs by assignment. Within each part, use Qualtrics Display Logic on the *question block* level. The amber-bordered **Display Logic** annotation boxes throughout this document indicate the exact condition to apply to each case block.

3 Case Vignettes Reference (All Parts)

The same ten cases are used throughout all five parts. This section provides the complete vignettes for reference. Each part will reproduce the relevant vignette alongside any additional standardised context.

Summary Reference Table

Case	Primary Presentation	Profile
C01	Sleep onset insomnia + cognitive hyperarousal + social anhedonia + somatic tension	28-year-old software developer; 2 months
C02	Recurrent panic attacks + anticipatory anxiety + situational avoidance	34-year-old healthcare administrator; 3 months
C03	Professional burnout + emotional exhaustion + social withdrawal	41-year-old secondary school teacher; 7 months
C04	Complicated grief + insomnia + guilt-based rumination	63-year-old retired professional; 9 months after loss
C05	Intrusive harm-related thoughts + compulsive reviewing + avoidance	25-year-old postgraduate student; 20 months
C06	Professional performance anxiety + post-event rumination + advancement avoidance	32-year-old marketing professional; >3 years
C07	Persistent low mood + anger + social isolation + hopelessness	46-year-old accounts manager; 14 months
C08	ADHD-related distress + executive dysfunction + relationship conflict	38-year-old project manager; childhood onset
C09	Emotional dysregulation + impulsivity + interpersonal instability	27-year-old sales representative; 3 years
C10	Chronic work stress + sleep maintenance insomnia + somatic + alcohol coping	52-year-old operations director; 9 months

4 Part 1: Operationalization of Mental State

Part 1 Instructions

Task: For each of your **two assigned case vignettes**, identify and operationalize the key mental health criteria present in the complaint. (This document shows all ten cases for researcher reference; Qualtrics Display Logic will show only your assigned two.)

What we mean by “operationalized criterion”: A criterion is a distinct, currently present, changeable dimension of the person’s mental state or functioning that (a) reflects a specific complaint aspect, (b) can in principle be measured repeatedly over time (e.g., via daily self-report), and (c) is not a demographic trait or static background factor.

Response format: For each case, list **2–6 criteria**. For each criterion provide:

- A **short label** (2–5 words, e.g., “Sleep onset insomnia”)
- A **brief description** (1–2 sentences: what it is and how it manifests here)

Use the structured template: **Label | Description** on one line per criterion. This allows direct automated comparison with machine outputs.

Qualtrics setup: 2 open-text questions per HCP (Text Entry — Large), shown via Display Logic. Prompt shown in each response box.

Time: Approximately 3 minutes per case (≈ 6 minutes for your two cases).

Case C01

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Profile: 28-year-old software developer; complaint duration: 2 months.

“Over the past two months I have been finding it very hard to fall asleep at night. My mind keeps going over everything I still need to do — deadlines, unfinished tasks, conversations I need to have. Even when I am exhausted, I cannot seem to switch my thoughts off. Something else I have noticed is that I no longer enjoy social occasions the way I used to. I go to gatherings with friends and I feel disconnected, like I am watching everyone else have fun from a distance. My neck and shoulders are also constantly tense and sore, which my GP says is likely stress-related.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C02

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Profile: 34-year-old healthcare administrator; complaint duration: 3 months.

“Around three months ago I started having episodes where my heart would pound, I felt like I could not get enough air, and I was convinced something terrible was happening to me. These episodes are

frightening and can come out of nowhere, even when I am sitting quietly at home. Since they started, I have become scared of having one in public and have been avoiding supermarkets, public transport, and crowded places. I have called in sick to work several times because I was afraid of having an episode there. I know logically that these situations are not dangerous but I cannot stop the fear.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C03

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Profile: 41-year-old secondary school teacher; complaint duration: 7 months.

“I have been teaching for fifteen years and I used to love my work, but over the past several months I feel completely empty. I go to school, deliver my lessons, and come home, but there is nothing left in me. I sit in front of the television every evening and cannot motivate myself to do anything else. I have stopped going out with friends and rarely reply to messages anymore. I have also started making mistakes at work — forgetting things, losing track of where I am in a lesson — which has never happened before. I feel ashamed about it but I do not know how to change things.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C04

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Profile: 63-year-old retired professional; 9 months after bereavement.

“My partner died nine months ago and I am still not coping well. I think about them constantly. Random things — a song, a smell, a television programme we used to watch together — trigger intense waves of grief that I cannot control. I am not sleeping properly; I wake in the early hours and cannot get back to sleep. I also feel a lot of guilt. I keep going over things I said or should have said, and times I was not patient enough. I cannot imagine what my future is supposed to look like anymore. My family tells me I should be feeling better by now, but I am not.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C05

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Profile: 25-year-old postgraduate student; complaint duration: approximately 20 months.

“I have thoughts that I might accidentally hurt people, even though I have no desire to do so and I know these thoughts are irrational. They come suddenly and feel very realistic and disturbing. After having one of these thoughts I spend a lot of time mentally reviewing what happened to reassure myself I have not done anything wrong. I have also started avoiding situations where I might hold something sharp, which affects my daily life and some social situations. This reviewing takes up hours of my day and is affecting my studies and my relationships.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C06

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Profile: 32-year-old marketing professional; complaint persistent since early career (>3 years).

“I have a great deal of anxiety around professional situations. Whenever I have to give a presentation, speak in a meeting, or receive feedback from my manager, I become so self-conscious that my mind goes blank. I spend hours preparing what to say beforehand but then freeze in the moment anyway. Afterwards I replay everything that happened and analyse every word I said for a long time. I have turned down several promotions and interesting projects because I was afraid of the visibility they would bring. I know my anxiety about how others perceive me is disproportionate but I cannot seem to stop it.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C07

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Profile: 46-year-old accounts manager; complaint duration: 14 months.

“For over a year I have woken up most mornings feeling heavy and grey. I would not call it dramatic sadness — more like a persistent flatness where nothing feels worthwhile. Very simple tasks feel like enormous effort: getting dressed, replying to an email, cooking a meal. I have gradually lost touch with almost everyone I used to socialise with because I do not have the energy to maintain relationships. I sleep a lot but never feel rested. Privately I wonder whether I will ever feel like my old self again. I

am seeking help because my GP told me I should.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C08

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Profile: 38-year-old project manager; difficulties since childhood, significantly worsened over past 12 months.

“I have always struggled to concentrate on tasks that do not immediately interest me, but it has gotten much worse over the past year. I start projects and rarely finish them. I forget appointments even when I have written them down. My partner is frustrated with me because they say I do not listen and that I am unreliable. I am competent in many ways but I feel like a constant failure. I have started avoiding tasks I know I will struggle with, which makes the backlog worse. I feel ashamed that something that seems straightforward for other people is so hard for me.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C09

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Profile: 27-year-old sales representative; complaint duration: 3 years.

“My mood changes very rapidly throughout the day and it often does not seem connected to what is actually happening around me. I can feel reasonably okay in the morning and be in tears by early afternoon. When I feel criticized or rejected, even over small things, my reaction is very intense — I become angry or very distressed almost instantly. I also act impulsively, particularly with money or in relationships, and I regret it afterwards. My relationships tend to become intense very quickly but then fall apart. I am aware that this pattern repeats but I do not know how to break it.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

Case C10

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Profile: 52-year-old operations director; complaint duration: approximately 9 months.

“Work stress has been building for the better part of a year. I have constant tension headaches and my stomach is unsettled most days. I lie awake at night going over work problems and decisions that need to be made, and I rarely fall back to sleep after waking. I have been having two or three glasses of wine each evening to help me wind down, which I know is not ideal but it feels necessary. My exercise routine has completely stopped and I have put on weight. I snap at people at home and then feel guilty about it. I am holding everything together externally but I am running on empty.”

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please list 2–6 operationalized criteria for this complaint. For each criterion, provide: (1) a short label (2–5 words) and (2) a brief description (1–2 sentences).

5 Part 2: Initial Observational Model Construction

Part 2 Instructions

Task: For each case, propose a set of **modifiable predictor variables** that could be monitored over time (e.g., via daily self-report / EMA) to explain or predict the criteria identified in the standardised context below.

What is a modifiable predictor? A predictor is a behavioural, cognitive, physiological, or environmental variable that (a) can change from day to day, (b) plausibly influences one or more of the criteria, and (c) can be measured by the person themselves with a simple daily rating or count.

Response format: List **3–5 predictors** using this structured template (one per line):

Label | 1-sentence measurement definition | CR connections (e.g. CR-1, CR-3)

This format allows direct automated comparison with machine outputs. Abbreviate freely; clinical precision is more important than length.

Context provided: Standardised criteria (from Step 01 reference run) appear for each case.

Qualtrics setup: 10 Text Entry (Large) questions. Copy the exact prompt shown in each response box.

Time: Approximately 3 minutes per case (≈ 6 minutes for your two cases).

Case C01 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

28-year-old software developer, 2 months. Difficulty falling asleep; bedtime rumination about tasks and deadlines; social disconnection and reduced positive affect; constant neck and shoulder tension.

Standardised criteria (from Step 01 reference run):

- **CR-1** *Sleep onset insomnia* — subjective difficulty initiating sleep; estimated latency >30 min most nights.
- **CR-2** *Cognitive hyperarousal at bedtime* — repetitive uncontrollable task-anticipatory thoughts at sleep onset.
- **CR-3** *Social anhedonia* — reduced capacity to experience positive affect in social contexts.
- **CR-4** *Musculoskeletal stress tension* — chronic neck and shoulder tension of probable stress etiology.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor, provide: (1) short label, (2) brief measurement definition (1 sentence), (3) which criteria (CR-1 to CR-4) it connects to.

Case C02 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

34-year-old healthcare administrator, 3 months. Recurrent panic episodes (palpitations, breathlessness, impending-doom cognition); persistent fear of future episodes; avoidance of public settings; occupational impact.

Standardised criteria (from Step 01 reference run):

- **CR-1** *Recurrent panic episodes* — discrete episodes with autonomic arousal and sudden onset.
- **CR-2** *Anticipatory anxiety* — persistent fear of having a further episode; interoceptive vigilance.
- **CR-3** *Situational avoidance* — active avoidance of supermarkets, public transport, crowded venues.
- **CR-4** *Occupational interference* — sick leave, avoidance of work attendance, reduced function.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C03 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

41-year-old teacher, 7 months. Emotional exhaustion; depersonalization/professional detachment; social withdrawal; concentration difficulties and occupational errors.

Standardised criteria:

- **CR-1** *Emotional exhaustion* — depleted affective resources; absence of professional engagement or motivation.
- **CR-2** *Professional depersonalization* — detached, going-through-the-motions quality of occupational performance.
- **CR-3** *Social withdrawal* — discontinued social activities; non-response to social contact.
- **CR-4** *Occupational cognitive impairment* — concentration difficulties, memory failures, and increased error rate at work.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C04 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

63-year-old retired professional, 9 months post-bereavement. Prolonged grief with intrusive memories; sleep maintenance insomnia; guilt-based counterfactual rumination; inability to envision a future.

Standardised criteria:

- **CR-1** *Prolonged grief response* — intrusive memories, yearning, difficulty accepting the loss.
- **CR-2** *Sleep maintenance insomnia* — early-morning awakening; inability to return to sleep.
- **CR-3** *Guilt-based counterfactual rumination* — repetitive thoughts about past interactions; self-blame and regret.
- **CR-4** *Reduced future orientation* — inability to imagine meaningful activities or goals beyond the loss.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C05 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

25-year-old postgraduate student, ~20 months. Intrusive harm-related obsessional thoughts; compulsive mental reviewing; situational avoidance; significant daily time interference (>2 hours).

Standardised criteria:

- **CR-1** *Intrusive harm-related obsessions* — unwanted ego-dystonic thoughts of accidentally harming others.
- **CR-2** *Compulsive reviewing behavior* — mental replaying and checking sequences triggered by intrusions.
- **CR-3** *Harm-trigger situational avoidance* — avoidance of sharp objects, specific locations, and social settings.
- **CR-4** *OCD time interference* — rituals and reviewing consuming >2 hours of productive daily time.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C06 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

32-year-old marketing professional, >3 years. Performance anxiety in professional situations; post-event rumination; pre-event preparation burden; avoidance of advancement opportunities.

Standardised criteria:

- **CR-1** *Professional performance anxiety* — fear and physiological arousal in presentations, meetings, and evaluations.
- **CR-2** *Post-event rumination* — prolonged post-hoc analysis of social and professional performances.
- **CR-3** *Pre-event cognitive preparation burden* — extensive anticipatory rehearsal consuming significant time and energy.
- **CR-4** *Advancement avoidance* — declining opportunities due to anxiety about increased visibility.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C07 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

46-year-old accounts manager, 14 months. Persistent low mood and anhedonia; significant anergia; social isolation; hopelessness about recovery.

Standardised criteria:

- **CR-1** *Persistent depressed mood* — pervasive low mood and flatness most days for >12 months.
- **CR-2** *Anergia and psychomotor slowing* — fatigue and effortful quality of routine daily tasks.
- **CR-3** *Social isolation* — loss of social network through withdrawal and failure to maintain contact.
- **CR-4** *Hopelessness about recovery* — stable negative expectation about future mood improvement.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C08 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

38-year-old project manager, childhood onset (worsened past 12 months). Sustained attention deficit; planning/organization failures; interpersonal inattentiveness; chronic performance shame.

Standardised criteria:

- **CR-1** *Sustained attention deficit* — difficulty maintaining focus on non-preferred, effortful tasks.
- **CR-2** *Planning and organization failures* — missed appointments, incomplete tasks, failure to use organizational systems.
- **CR-3** *Interpersonal inattentiveness* — partner reports not being listened to; perceived unreliability in relationships.
- **CR-4** *Chronic performance shame* — pervasive self-judgment tied to executive functioning difficulties.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C09 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

27-year-old sales representative, 3 years. Rapid intra-day mood swings; intense emotional reactivity to perceived criticism/rejection; impulsive behaviour; recurrent relationship instability.

Standardised criteria:

- **CR-1** *Rapid intra-day mood cycling* — multiple significant mood shifts within a single day.
- **CR-2** *Emotional reactivity to rejection cues* — intense and rapid affective response to perceived criticism or abandonment.
- **CR-3** *Impulsive behaviour patterns* — poorly modulated spending, relationship decisions, or interpersonal actions.
- **CR-4** *Recurrent relationship instability* — pattern of rapid escalation followed by breakdown in close relationships.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

Case C10 — Part 2

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

52-year-old operations director, 9 months. Chronic occupational stress response; sleep maintenance insomnia; somatic stress symptoms; evening alcohol use as coping mechanism.

Standardised criteria:

- **CR-1** *Chronic occupational stress* — persistent physiological stress response driven by work demands.
- **CR-2** *Sleep maintenance insomnia* — early awakening with prolonged wakefulness; ruminative nocturnal cognition.
- **CR-3** *Somatic stress expression* — recurrent tension headaches and gastrointestinal symptoms.
- **CR-4** *Alcohol use as coping* — regular evening alcohol consumption as a relaxation/sleep aid.

Qualtrics item type: Text Entry (Large / Essay). *Prompt:* Please propose 3–6 modifiable predictor variables for this case. For each predictor: (1) short label, (2) measurement definition, (3) connected criteria.

6 Part 3: Treatment-Target Identification

Part 3 Instructions

Task: For each case, **rank the top 2–4 treatment targets** from the bipartite network diagram provided.

What is a treatment target? A modifiable predictor (P-node) from the initial model that, if changed, is most likely to reduce the person's complaint criteria. Select based on: (a) monitoring data evidence, (b) clinical modifiability, and (c) downstream impact on multiple criteria.

Visual key: Each diagram shows **criteria nodes (blue, left)** and **predictor nodes (teal, right)**. Edge weight reflects the monitoring-derived importance of the predictor–criterion relationship:

—— Strong — Moderate - - - - Weak

Fair comparison note: The PHOENIX system processes the same numerical edge weights shown here. The visual is a human-readable representation of identical data. No additional information is presented to you vs. the machine.

Response format: Qualtrics Rank Order item. Drag/rank the predictor labels from the network in order of treatment priority (1 = highest). Select only 2–4 items.


Time: Approximately 3.5 minutes per case (≈ 7 minutes for your two cases).

Case C01 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Sleep onset insomnia, bedtime cognitive hyperarousal, social anhedonia, musculoskeletal tension. Duration: 2 months.

21-day monitoring summary: Screen time past 10 PM: 16/21 evenings (avg 78 min). Task incompleteness: avg 6.2 items/day. Social events: 2.1/week (enjoyment 2.3/10). Pre-sleep offloading used: 3/21 evenings. Physical activity: avg 0.7 sessions/week.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C02 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Recurrent panic episodes, anticipatory anxiety, situational avoidance, occupational impact. Duration: 3 months.

21-day monitoring summary: Panic episodes: avg 2.3/week. Avoidance events: avg 4.1/day. Safety behaviours: avg 8.7/day. Planned exposures: avg 0.3/day. Work attendance: 60%. Anticipated anxiety in public settings: 7.8/10.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2-4 targets only.

Case C03 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Emotional exhaustion, professional depersonalization, social withdrawal, occupational cognitive difficulties. Duration: 7 months.

21-day monitoring summary: Evening work violations: avg 5.4/day. Recovery activities: avg 1.2 events/week. Social contacts initiated: avg 0.7/week. Professional mastery moments: avg 0.9/day. Sleep quality: avg 4.1/10.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C04 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Prolonged grief response, sleep maintenance insomnia, guilt-based rumination, absent future orientation. Duration: 9 months post-bereavement.

21-day monitoring summary: Grief intrusion events: avg 4.2/day. Sleep quality: avg 2.8/10. Social contacts: avg 3.1/week. Guilt thoughts: avg 5.8/day. Future-plan activities: avg 0.4/week. Meaning-making activity: 4/21 days.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C05 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Intrusive harm-related obsessions, compulsive reviewing, harm-trigger avoidance, OCD time interference (>2 hr/day). Duration: 20 months.

21-day monitoring summary: Compulsion/review time: avg 2.4 hrs/day. Avoidance events: avg 3.8/day. Resistance practice: avg 18 min/day. Distress at resisting: 7.2/10. Sleep: avg 5.6 hrs/night. Reassurance seeking: avg 4.1/day.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C06 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Professional performance anxiety, post-event rumination, pre-event preparation burden, advancement avoidance. Duration: >3 years.

21-day monitoring summary: Presentations entered: avg 0.9/week. Post-event processing time: avg 68 min/event. Pre-event rehearsal: avg 3.2 hrs/event. Perceived performance vs. actual: rated 4.1 vs. 6.8/10. Advancement opportunities declined: 3 in past 4 weeks.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C07 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Persistent low mood and anhedonia, anergia, social isolation, hopelessness. Duration: 14 months.

21-day monitoring summary: Activity completion rate: 28%. Social contacts/week: avg 0.8. Mood rated: avg 2.9/10. Sleep wake-time variance: avg ± 2.3 hrs. Pleasurable activities: avg 0.9/day. Hopelessness challenge used: avg 0.6/day.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C08 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Sustained attention deficit, planning failures, interpersonal inattentiveness, performance shame. Duration: childhood onset.

21-day monitoring summary: Tasks initiated vs. completed: 68% vs. 29%. Missed commitments: avg 2.1/week. Partner conflict events: avg 4.7/week. External tool use: avg 1.8 uses/day. Positive self-acknowledgment: avg 0.4/day.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C09 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Intra-day mood cycling, emotional reactivity to rejection, impulsive behaviour, relationship instability.
Duration: 3 years.

21-day monitoring summary: Daily mood range: avg 5.8/10-point scale. ER skill use: avg 1.2/day. Impulsive actions: avg 2.1/day. Interpersonal conflicts: avg 3.8/week. Rejection events perceived: avg 2.4/day. Reappraisal before reacting: avg 0.7/day.




Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

Case C10 — Part 3

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Chronic occupational stress, sleep maintenance insomnia, somatic stress expression, evening alcohol use as coping. Duration: 9 months.

21-day monitoring summary: Work-related thoughts after 8 PM: avg 3.4 hrs. Alcohol units: avg 3.1/evening. Headache days: 16/21. Sleep efficiency: avg 62%. Daytime stress regulation: avg 0.9 events/day. Physical activity: avg 0.4 sessions/week.



Qualtrics item type: Rank Order. *Prompt:* Rank the predictors above in order of treatment priority (1 = highest). Select 2–4 targets only.

7 Part 4: Updated Observation Model

Part 4 Instructions

Task: For each case, **select the 2–4 predictors to retain in the next monitoring cycle**, given the treatment targets identified in Part 3 and the monitoring data pattern.

Response format: Multiple Choice (select all that apply) — tick the predictors from the initial model that should be tracked next cycle. Optionally add a 1-sentence note per predictor if an operationalization change is needed.

This structured format is directly comparable to the machine output, which produces a binary retain/drop decision per predictor plus an optional rationale string.

Context provided: Standardised treatment targets from Step 03 reference run (shown per case).

Time: Approximately 3 minutes per case (≈ 6 minutes for your two cases).

Case C01 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Standardised top treatment targets (from Step 03 reference run):

1. Evening screen time duration (*highest impact on CR-1 and CR-2; data shows consistent late-night use*)
2. Pre-sleep cognitive offloading practice (*near-zero usage despite high task backlog; direct CR-2 pathway*)
3. Aerobic physical activity (*very low frequency; connects to both CR-1 and CR-4*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C02 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-01**.

Standardised top treatment targets:

1. Planned exposure attempts (*nearly absent in monitoring data; direct avoidance reduction pathway*)
2. Safety behaviour frequency (*high frequency sustaining anticipatory anxiety; key maintenance loop*)
3. Interoceptive focus duration (*elevated; fuels panic escalation cycle*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C03 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Standardised top treatment targets:

1. Work-life boundary adherence (*highest maintenance factor for exhaustion; avg 5.4 violations/day*)
2. Recovery activity engagement (*critical low at 1.2 events/week; addresses CR-1 and CR-3*)
3. Professional mastery moments (*addresses depersonalization; low but showing variability*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C04 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-02**.

Standardised top treatment targets:

1. Structured meaning-making activity (*used only 4/21 days; high potential for CR-1 and CR-4*)
2. Sleep routine consistency (*poor sleep quality; downstream effects on grief processing*)
3. Self-compassion practices (*guilt ruminative burden high; underused coping resource*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C05 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Standardised top treatment targets:

1. Compulsion delay and resistance practice (*used minimally; avg 18 min vs target 60+; core ERP component*)
2. Avoidance scope management (*high avoidance events maintain the obsessional cycle*)
3. Uncertainty tolerance behaviour (*low; drives both intrusions and reviewing*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C06 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-03**.

Standardised top treatment targets:

1. Graduated professional exposure frequency (*low frequency; key mechanism for anxiety reduction*)
2. Post-event processing duration (*avg 87 min/event; major maintenance factor for CR-2*)
3. Avoidance decision logging (*avg 1.4 opportunities declined/week; directly limits advancement*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C07 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Standardised top treatment targets:

1. Daily activity scheduling and completion (*completion rate 28%; behavioral activation core mechanism*)
2. Sleep schedule regularity (*±2.3 hr variance; disrupts mood and energy regulation*)
3. Social contact initiation (*avg 0.8/week; critical for CR-3 and indirect mood lifting*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C08 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-04**.

Standardised top treatment targets:

1. External task structuring tool use (*inconsistent at 1.8 uses/day; core scaffolding system*)
2. Intentional listening behaviour (*partner conflict avg 4.7/week; direct CR-3 pathway*)
3. Task completion reinforcement (*very low; shame cycle maintenance factor*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C09 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Standardised top treatment targets:

1. Emotion regulation skill application (*low use; mood range 5.8 pts/day; primary intervention lever*)
2. Impulse delay behaviour (*avg 2.1 impulsive actions/day; core target for CR-3 and CR-4*)
3. Rejection appraisal reframing (*avg 0.7/day despite 2.4 perceived rejection events/day*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

Case C10 — Part 4

Qualtrics Display Logic: Show this block only if participant code = **HCP-PRE-05**.

Standardised top treatment targets:

1. Pre-sleep work disengagement routine (*absent; nocturnal cognitive arousal drives CR-2*)
2. Evening alcohol units (*avg 3.1 units; perpetuates insomnia; priority reduction target*)
3. Daytime stress regulation behaviour (*avg 0.9 events/day; low but trainable*)

Qualtrics item type: Multiple Choice (select all that apply) + optional Text Entry. *Prompt:* Select 2–4 predictors to retain in the next monitoring cycle. Tick each from the list above. Optional: add a 1-sentence operationalization note per selected predictor if a measurement adjustment is needed.

8 Part 5: Tailored Intervention Message

Part 5 Instructions

Task: For each case, complete two items: (A) identify the **HAPA motivational phase** most relevant to this person, and (B) write a **brief personalized coaching message**.

(A) HAPA phase classification: Select the phase that best characterizes where this person currently is:

- **Pre-intentional** — not yet motivated to change; needs motivational/risk communication
- **Intentional** — motivated but not yet planning; needs goal-setting and planning support
- **Action/Maintenance** — already attempting change; needs coping planning and self-efficacy support

(B) Message format: 2–3 sentences, second-person, warm and actionable. Address the top treatment target and the key coping strategy. No jargon; no diagnostic labels.

Response format: Two Qualtrics items per case — (A) multiple choice for phase, (B) Text Entry (Medium) for message.

Context provided: Top treatment target, primary barrier type, and suggested coping strategy per case.

Time: Approximately 3.5 minutes per case (≈ 7 minutes for your two cases).

Case C01 — Part 5

Primary challenge: Difficulty falling asleep due to an overactive mind at bedtime; social disconnection.

Top treatment target: Evening screen time reduction and pre-sleep wind-down routine.

Main barrier: *Self-efficacy* — habitual screen use is experienced as a necessary decompression tool; low confidence that reducing it is possible without losing the relaxation benefit.

Coping strategy: Implementation intention with a substitution plan (replace screen time with a brief physical or written offloading activity in the 30 minutes before bed).

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.

Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C02 — Part 5

Primary challenge: Fear of panic episodes leading to increasing avoidance and reduced daily functioning.

Top treatment target: Planned exposure — gradually re-entering feared situations.

Main barrier: *Outcome expectancy* — strong belief that entering avoided situations will inevitably lead to a panic episode that cannot be managed.

Coping strategy: Planning a hierarchy of small, manageable exposure steps with a coping script prepared in advance.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C03 — Part 5

Primary challenge: Profound occupational exhaustion with loss of meaning and progressive social withdrawal.

Top treatment target: Work-life boundary enforcement — structured protection of non-work time.

Main barrier: *Action planning deficit* — no concrete plan for what a work boundary looks like or when it applies; boundary attempts collapse under work demands.

Coping strategy: Implementation intention with a specific boundary signal (e.g., device-off at a set time) paired with one simple recovery activity immediately after.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C04 — Part 5

Primary challenge: Prolonged grief with pervasive guilt and absence of forward-looking engagement.

Top treatment target: Structured meaning-making activity — engaging with memory, legacy, or creative activities that honour the loss and rebuild connection to the future.

Main barrier: *Motivational deficit* — absence of motivation or a clear sense of what is worth doing; grief-related anhedonia blocks initiation.

Coping strategy: Reducing the size of the first step to a minimal, time-limited activity (e.g., writing one sentence in a memory book once this week).

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C05 — Part 5

Primary challenge: Distressing intrusive thoughts with extensive compulsive reviewing consuming hours daily.

Top treatment target: Compulsion delay and resistance practice — extending the time between intrusion and reviewing response.

Main barrier: *Perceived risk* — strong overestimation that not reviewing will lead to actual harm; reviewing feels morally necessary even when recognized as irrational.

Coping strategy: Delay rule — committing to a 5-minute pause before any reviewing begins, with a brief mindful grounding exercise during the delay.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C06 — Part 5

Primary challenge: Intense anxiety in professional performance situations with extensive post-event rumination.

Top treatment target: Graduated professional exposure — entering and completing professional situations despite anxiety.

Main barrier: *Self-efficacy gap* — strong conviction that performance is objectively poor and that exposure will confirm this; distorted self-evaluation persists despite contradictory external feedback.

Coping strategy: Behavioural experiment approach — after the next professional exposure, record the predicted rating and then gather actual feedback to compare.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C07 — Part 5

Primary challenge: Persistent low mood and anergia with near-complete loss of activity and social connection.

Top treatment target: Daily activity scheduling and completion — behavioural activation through planned, minimal daily activities.

Main barrier: *Planning barrier* — no structured activity menu available; activities feel meaningless before attempting them; anergia blocks initiation.

Coping strategy: Activity menu strategy — pre-selecting three very small activities (5 minutes each) the night before, with a “done-is-enough” standard.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C08 — Part 5

Primary challenge: Executive functioning difficulties creating task failures, relationship strain, and chronic shame.

Top treatment target: External task structuring tool use — consistent daily use of an external system (written or digital) to scaffold task planning and completion.

Main barrier: *Habit resistance* — tools have been tried before but abandoned; low self-efficacy about sustaining system use; shame about needing external scaffolding.

Coping strategy: Minimum viable habit — one non-negotiable daily check-in with the tool (morning

review, 3 minutes) as an anchored routine rather than a complex system.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C09 — Part 5

Primary challenge: Rapid mood swings and intense emotional reactivity causing impulsive behaviour and relationship breakdown.

Top treatment target: Emotion regulation skill application — consistent use of a specific, rehearsed skill when emotional intensity rises.

Main barrier: *Coping self-efficacy* — strong doubts about ability to manage intense affect; past skill use attempts have felt ineffective in the moment of peak emotion.

Coping strategy: Pre-committed cue-action plan — identifying one specific physical cue of rising emotion and linking it to one specific skill (e.g., TIPP, grounding) rehearsed daily in low-affect moments.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

Case C10 — Part 5

Primary challenge: Chronic work stress with fragmented sleep, somatic symptoms, and alcohol as the primary coping mechanism.

Top treatment target: Pre-sleep work disengagement routine — a structured sequence that creates a clear boundary between work and sleep.

Main barrier: *Habit strength* — nightly alcohol use is a deeply established coping pattern perceived as essential for sleep; no alternative wind-down routine currently exists.

Coping strategy: Routine substitution — designing a specific 20-minute wind-down sequence that begins at the same time each evening and replaces (rather than prohibits) alcohol as a cue-to-sleep signal.

Qualtrics item type: (A) Multiple Choice — HAPA phase; (B) Text Entry (Medium) — message.
Prompt: (A) Select the HAPA motivational phase for this person. (B) Write a 2–3 sentence personalized coaching message addressing the top treatment target and coping strategy above.

9 Completion and Submission

Thank you for completing all five parts of this survey. Your responses across the ten cases will form the expert reference corpus for the PHOENIX evaluation study. Before submitting, please review the checklist below.

Pre-submission checklist

- ☐ I have provided responses for all 10 cases in Part 1 (operationalization).
- ☐ I have provided responses for all 10 cases in Part 2 (initial model).
- ☐ I have provided responses for all 10 cases in Part 3 (treatment targets).
- ☐ I have provided responses for all 10 cases in Part 4 (updated model).
- ☐ I have provided responses for all 10 cases in Part 5 (intervention message).
- ☐ My responses reflect my genuine clinical judgment without external reference.
- ☐ I understand that my responses will be anonymized and used solely for evaluation purposes.

Once submitted, your responses will be assigned a pseudonymous participant ID and will not be modifiable. If you have any questions or concerns, please contact the researcher at stijn.vanseveren@ugent.be before submitting.

Qualtrics: End of Survey block

Include: automatic acknowledgement message and unique response ID display.

Optional: demographic block (professional background, years of clinical experience, primary discipline) for covariate adjustment in the analysis.

10 Qualtrics Configuration Notes

Block Structure

Implement the survey as five sequential Qualtrics **blocks**, one per part. Configure the following:

1. **Block display logic:** Show blocks sequentially (Block 1 must be completed before Block 2 is unlocked). Use Qualtrics *Survey Flow* to enforce order.
2. **Item type:** All 50 response items are *Text Entry (Essay/Large)* questions. Do not add scale or multiple-choice items within the generation phase.
3. **Forced response:** Set all 50 items to *Request response* (allow but flag missing). Do not use forced response, as some HCPs may prefer to leave a case blank rather than respond speculatively.
4. **Progress bar:** Enable the progress bar. Set milestones at Part 1 end (20%), Part 2 end (40%), Part 3 end (60%), Part 4 end (80%), and submission (100%).
5. **Session saving:** Enable *Save and Continue* so participants can complete across multiple sittings.
6. **Timing:** Enable *Page Timer* on each block for session-length analysis.
7. **Anonymity:** Disable IP recording if institutional ethics requires anonymous participation; use anonymous survey links.
8. **Demographic block:** Add an optional final block collecting: (a) professional discipline (psychiatry / clinical psychology / nursing / general practice / other), (b) years of clinical experience, (c) primary patient population (adult, youth, older adult, mixed), (d) familiarity with digital mental health tools (1–5).

Participant ID Assignment

Assign each participant a random 4-character alphanumeric ID (e.g., HCP_PRE_A3F7) at survey start. Display it prominently on the completion page. Record it in the response metadata. This ID is used to link PRE-phase responses to the blinded POST-phase evaluation.

Data Export Format

Export responses as CSV with one column per item (Part#_Case#) and one row per participant. The column naming convention is: P1_C01, P1_C02, ..., P5_C10.