

MATH 173 PROBLEM SET 9

Stepan (Styopa) Zharkov

June 1, 2022

Problem 1.

◁

Solution.

Problem 2.

◁

Solution.

- (a) We need $\int_0^1 |x^\alpha|^2 = \int_0^1 x^{2\alpha}$ to converge. This converges for $\alpha > -1/2$ and diverges for $\alpha \leq -1/2$, so $\phi_\alpha \in L^2((0, 1))$ for $\alpha > -1/2$. \square
- (b) We need $\phi_\alpha \in L^2((0, 1))$, so $\alpha > -1/2$. But, since ϕ_α are smooth, we also need $\int_0^1 |\phi'_\alpha|^2$ to converge. We see $\phi'_\alpha = \alpha x^{\alpha-1}$. and $\int_0^1 |\alpha x^{\alpha-1}|^2 = |\alpha|^2 \int_0^1 x^{2(\alpha-1)}$ converges for $\alpha > 1/2$ and diverges for $\alpha \leq 1/2$. So, $\phi_\alpha \in H^1((0, 1))$ for $\alpha > 1/2$. \square

Problem 3.

◁

Solution.

Problem 4.

◁

Solution.

Problem 5.

◁

Solution.

Problem 6.

◁

Solution.

Problem 7.

◁

Solution.