

MATH 173 PROBLEM SET 9

Stepan (Styopa) Zharkov

June 1, 2022

Problem 1.

Solution.

◁

Problem 2.

◁

Solution.

- (a) We need $\int_0^1 |x^\alpha|^2 = \int_0^1 x^{2\alpha}$ to converge. This converges for $\alpha > -1/2$ and diverges for $\alpha \leq -1/2$, so $\phi_\alpha \in L^2((0, 1))$ for $\alpha > -1/2$. \square
- (b) We need $\phi_\alpha \in L^2((0, 1))$, so $\alpha > -1/2$. But, since ϕ_α are smooth, we also need $\int_0^1 |\phi'_\alpha|^2$ to converge. We see $\phi'_\alpha = \alpha x^{\alpha-1}$. and $\int_0^1 |\alpha x^{\alpha-1}|^2 = |\alpha|^2 \int_0^1 x^{2(\alpha-1)}$ converges for $\alpha > 1/2$ and diverges for $\alpha \leq 1/2$. So, $\phi_\alpha \in H^1((0, 1))$ for $\alpha > 1/2$. \square

Problem 3.

◁

Solution.

- (a) We know the statement is true for $f \in C^1((a, b))$ by FTC. Now, let $f_n \rightarrow f$ where $f_n \in C^1((a, b))$.
By the continuity of the trace operator,

$$f(x) - f(y) = \lim_{n \rightarrow \infty} (f_n(x) - f_n(y)) = \lim_{n \rightarrow \infty} \int_x^y f'_n(t) dt$$

TODO: finish this.

- (b) TODO

Problem 4.

◁

Solution.

Problem 5.

◁

Solution. Consider the dogbowl functions

$$f_n := 1 - \min(n \cdot d(x, \partial B), 1).$$



We see that $Tf_n = 1$ for all $n \geq 4$, so $Tf_n \rightarrow 1 \neq 0$. However,

$$\|f_n\|_{L^2}^2 = \int_B |f_n(x)|^2 dx = \int_{d(x, \partial B) < 1/n} |f_n(x)|^2 dx \leq \int_{d(x, \partial B) < 1/n} 1 dx = O(1/n) \rightarrow 0.$$

So, $f_n \rightarrow 0$ in L^2 .

□

Problem 6.

◁

Solution. Let $u = \lim_{n \rightarrow \infty} u_n$ where u_n are compactly supported continuous functions. Note that we are given that $u = \lim_{n \rightarrow \infty} -u_n(x^*)$. This means that

$$u = \frac{\lim_{n \rightarrow \infty} u_n(x) + \lim_{n \rightarrow \infty} -u_n(x^*)}{2} = \lim_{n \rightarrow \infty} \frac{u_n(x) - u_n(x^*)}{2}.$$

Note that $\frac{u_n(x) - u_n(x^*)}{2} = 0$ when $x_n = 0$, so

$$T_{B_+} \left(\frac{u_n(x) - u_n(x^*)}{2} \right) = \frac{u_n(x) - u_n(x^*)}{2} |_{\partial B_+} = 0.$$

We have shown in class that this is sufficient to say $T_{B_+}(u|_{B_+}) = 0$, so $u|_{B_+} \in H_0^1(B_+)$. □

Problem 7.

◁

Solution.

- (a) Let $V_k = \{x : |x| < 1/k\}$ and let $W_k = \{x : 1 - 1/k < |x| < 1\}$. Then, consider lemonsqueezer functions $f_k \in C_0^1(U)$ such that $f_k|_{B(V_k \cap W_k)} = 1$ and $f_k|_{V_{2k} \cup W_{2k}} = 0$.

TODO: image

Note that $u_k := uf_k \in C_0^1(U)$. We claim $u_k \rightarrow u$ in $H^1(B)$. Note that

$$\|u - u_k\|_{H^1}^2 = \int_B |u - u_k|^2 + \int_B |\nabla u - \nabla u_k|^2.$$

Now, since u is bounded

$$\begin{aligned} \int_B |u - u_k|^2 &= \int_B |u|^2 |1 - f_k|^2 \\ &= \int_B O(1) |1 - f_k|^2 \\ &= O(1) \int_B |1 - f_k|^2 \\ &= O(1) \int_{V_k \cup W_k} |1 - f_k|^2 \\ &= O(1) \int_{V_k \cup W_k} O(1) \\ &= O(1/k^2) = o(1) \end{aligned}$$

where the asymptotic notation is with respect to k . Also,

$$\int_B |\nabla u - \nabla u_k|^2 = \int_B |\nabla u - \nabla u f_k - u \nabla f_k|^2.$$

(b) TODO

(c) TODO