

# MATH 173 PROBLEM SET 9

Stepan (Styopa) Zharkov

June 1, 2022

**Problem 1.**

*Solution.*

◁

**Problem 2.**

◁

***Solution.***

- (a) We need  $\int_0^1 |x^\alpha|^2 = \int_0^1 x^{2\alpha}$  to converge. This converges for  $\alpha > -1/2$  and diverges for  $\alpha \leq -1/2$ , so  $\phi_\alpha \in L^2((0, 1))$  for  $\alpha > -1/2$ .  $\square$
- (b) We need  $\phi_\alpha \in L^2((0, 1))$ , so  $\alpha > -1/2$ . But, since  $\phi_\alpha$  are smooth, we also need  $\int_0^1 |\phi'_\alpha|^2$  to converge. We see  $\phi'_\alpha = \alpha x^{\alpha-1}$ . and  $\int_0^1 |\alpha x^{\alpha-1}|^2 = |\alpha|^2 \int_0^1 x^{2(\alpha-1)}$  converges for  $\alpha > 1/2$  and diverges for  $\alpha \leq 1/2$ . So,  $\phi_\alpha \in H^1((0, 1))$  for  $\alpha > 1/2$ .  $\square$

**Problem 3.**

◁

***Solution.***

- (a) We know the statement is true for  $f \in C^1((a, b))$  by FTC. Now, let  $f_n \rightarrow f$  where  $f_n \in C^1((a, b))$ . By the continuity of the trace operator,

$$f(x) - f(y) = \lim_{n \rightarrow \infty} (f_n(x) - f_n(y)) = \lim_{n \rightarrow \infty} \int_x^y f'_n(t) dt$$

TODO: finish this.

- (b) TODO

**Problem 4.**

◁

*Solution.*

**Problem 5.**

◁

**Solution.** Consider the dogbowl functions

$$f_n := 1 - \min(n \cdot d(x, \partial B), 1).$$



We see that  $Tf_n = 1$  for all  $n \geq 4$ , so  $Tf_n \rightarrow 1 \neq 0$ . However,

$$\|f_n\|_{L^2}^2 = \int_B |f_n(x)|^2 dx = \int_{d(x, \partial B) < 1/n} |f_n(x)|^2 dx \leq \int_{d(x, \partial B) < 1/n} 1 dx = O(1/n) \rightarrow 0.$$

So,  $f_n \rightarrow 0$  in  $L^2$ .

□

**Problem 6.**

◁

**Solution.** Let  $u = \lim_{n \rightarrow \infty} u_n$  where  $u_n$  are compactly supported continuous functions. Note that we are given that  $u = \lim_{n \rightarrow \infty} -u_n(x^*)$ . This means that

$$u = \frac{\lim_{n \rightarrow \infty} u_n(x) + \lim_{n \rightarrow \infty} -u_n(x^*)}{2} = \lim_{n \rightarrow \infty} \frac{u_n(x) - u_n(x^*)}{2}.$$

Note that  $\frac{u_n(x) - u_n(x^*)}{2} = 0$  when  $x_n = 0$ , so

$$T_{B_+} \left( \frac{u_n(x) - u_n(x^*)}{2} \right) = \frac{u_n(x) - u_n(x^*)}{2} |_{\partial B_+} = 0.$$

We have shown in class that this is sufficient to say  $T_{B_+}(u|_{B_+}) = 0$ , so  $u|_{B_+} \in H_0^1(B_+)$ . □

**Problem 7.**

◁

**Solution.**

- (a) Let  $V_n = \{x : |x| < 1/n\}$  and let  $W_n = \{x : 1 - 1/n < |x| < 1\}$ . Then, consider lemonsqueezer functions  $f_n \in C_0^1(U)$  such that  $f_n|_{B(V_n \cap W_n)} = 1$  and  $f_n|_{V_{2n} \cup W_{2n}} = 0$ .

TODO: image

Note that  $u_n := uf_n \in C_0^1(U)$ . We claim  $u_n \rightarrow u$  in  $H^1(B)$ . Note that

$$\|u - u_n\|_{H^1}^2 = \int_B |u - u_n|^2 + \int_B |\nabla u - \nabla u_n|^2.$$

Now, since  $u$  is bounded,

$$\begin{aligned} \int_B |u - u_n|^2 &= \int_B |u|^2 |1 - f_n|^2 \\ &= \int_B O(1) |1 - f_n|^2 \\ &= O(1) \int_B |1 - f_n|^2 \\ &= O(1) \int_{V_n \cup W_n} |1 - f_n|^2 \\ &= O(1) \int_{V_n \cup W_n} O(1) \\ &= O(1/n^2) \end{aligned}$$

(b) TODO

(c) TODO