

# MATH 173 PROBLEM SET 9

Stepan (Styopa) Zharkov

June 1, 2022

**Problem 1.**

*Solution.*

◁

**Problem 2.**

◁

***Solution.***

- (a) We need  $\int_0^1 |x^\alpha|^2 = \int_0^1 x^{2\alpha}$  to converge. This converges for  $\alpha > -1/2$  and diverges for  $\alpha \leq -1/2$ , so  $\phi_\alpha \in L^2((0, 1))$  for  $\alpha > -1/2$ .  $\square$
- (b) We need  $\phi_\alpha \in L^2((0, 1))$ , so  $\alpha > -1/2$ . But, since  $\phi_\alpha$  are smooth, we also need  $\int_0^1 |\phi'_\alpha|^2$  to converge. We see  $\phi'_\alpha = \alpha x^{\alpha-1}$ . and  $\int_0^1 |\alpha x^{\alpha-1}|^2 = |\alpha|^2 \int_0^1 x^{2(\alpha-1)}$  converges for  $\alpha > 1/2$  and diverges for  $\alpha \leq 1/2$ . So,  $\phi_\alpha \in H^1((0, 1))$  for  $\alpha > 1/2$ .  $\square$

**Problem 3.**

◁

***Solution.***

- (a) We know the statement is true for  $f \in C^1((a, b))$  by FTC.
- (b) TODO

**Problem 4.**

◁

*Solution.*

**Problem 5.**

◁

*Solution.*

**Problem 6.**

◁

*Solution.*

**Problem 7.**

◁

*Solution.*