

# MATH 173 PROBLEM SET 7

Stepan (Styopa) Zharkov

May 18, 2022

## Problem 1.

◁

**Solution.** This problem is straightforward. Since  $\overline{\Omega}$  is closed, and  $c > 0$ , there exists constants  $C_1, C_2 > 0$  such that  $C_1 < c(x) < C_2$ . Assume  $C_1 < 1$  and  $C_2 > 1$ . If not, we can always choose smaller  $C_1$  and larger  $C_2$ . So, for any  $u \in C^1(\overline{\Omega})$ ,

$$\begin{aligned} \|u\|_{H_c^1(\Omega)}^2 &= \int_{\Omega} |u(x)|^2 dx + \int_{\Omega} c(x) |\nabla u(x)|^2 dx \\ &\geq C_1^2 \int_{\Omega} |u(x)|^2 dx + \int_{\Omega} C_1^2 |\nabla u(x)|^2 dx \\ &= C_1^2 \|u\|_{H^1(\Omega)}^2. \end{aligned}$$

Similarly,

$$\begin{aligned} \|u\|_{H_c^1(\Omega)}^2 &= \int_{\Omega} |u(x)|^2 dx + \int_{\Omega} c(x) |\nabla u(x)|^2 dx \\ &\leq C_2^2 \int_{\Omega} |u(x)|^2 dx + \int_{\Omega} C_2^2 |\nabla u(x)|^2 dx \\ &= C_2^2 \|u\|_{H^1(\Omega)}^2. \end{aligned}$$

So, the statement follows for all  $u \in C^1(\overline{\Omega})$ . By continuity and density, it follows that the statement holds for  $u \in H^1(\Omega)$ .  $\square$

**Problem 2.**

◁

*Solution.* TODO

**Problem 3.**

◁

*Solution.* TODO

**Problem 4.**

◁

*Solution.* TODO

**Problem 5.**

◁

*Solution.* TODO

**Problem 6.**

◁

*Solution.* TODO