

UUB Charge and Peak histograms

Mauricio Suárez Durán and Ioana C. Mariş


IIHE-ULB

June 21, 2021





UUB Charge and Peak histograms

- ▶ Station studied: 863 1222 1219 1211 1740 1743 1221 1223 1217 1747 1741 1745 1818 1851 1729 1735 1746 1819 1791
- ▶ Data from CDAS.
- ▶ Software CDAS, pre-production version.




UUB Peak: Derivating histogram







- The algorithm:
1. Smooth/FFT Histogram
 2. Derivating of smooth/FFT Histogram
 3. Identifying Fit range (Slope changes)
 4. Fitting

Two function checked:

- Exp. + Log-normal
- 2nd order polinomial




UUB Peak: Fit and Residuals





The Exp.+Log-normal fit better than second order polinomium.





UUB Peak: Residuals distribution







UUB Peak: applying all histograms St. 863, Chi and Prob. distributions



UUB Charge: Derivating histogram




UUB Charge: Fit and Residuals





The Exp.+Log-normal fit better than second order polinomial.




UUB Charge: Residuals distribution





UUB Charge: applying all histograms St. 863, Chi and Prob. distributions



UUB AoP Station 863: Peak, Charge and AoP





UUB Peak Station 1740: Chi and Prob. distributions all histograms






For χ^2 / ndf plot, all histograms with χ^2 / ndf bigger than 6 are counted as 6.

UUB Charge Station 1740: Chi and Prob. distributions all histograms






For χ^2/ndf plot, all histograms with χ^2/ndf bigger than 6 are counted as 6.



UUB Station 1740: Failed fit




UUB AoP Station 1740: Peak, Charge and AoP







UB Peak: Derivating histogram



- The algorithm:
1. Smooth/FFT Histogram
 2. Derivating of smooth/FFT Histogram
 3. Identifying Fit range (Slope changes)
 4. Fitting
- Two function checked:
- Exp. + Log-normal
 - 2nd order polinomial




UB Peak: Fit and Residuals





The Exp.+Log-normal fit better than second order polinomial.


UB Peak: Residuals distribution







UB Peak: applying all histograms St. 863, Chi and Prob. distributions







UB Peak Station 863: Failed fit for PMT2



UB Charge: Derivating histogram




UB Charge: Fit and Residuals



The Exp.+Log-normal fit better than second order polinomial.

UB Charge: Residuals distribution



UB Charge: applying all histograms St. 863, Chi and Prob. distributions

