# SEDA: Spectral Energy Distribution Analyzer for Forward Modeling and Empirical Analysis of Ultracool Objects

11 February 2026

---


## Summary

Brown dwarfs and extrasolar gas giant planets exhibit complex, dynamic atmospheres that undergo diverse physical and chemical processes, including disequilibrium chemistry, cloud formation, auroral activity, and variability, which depend on fundamental properties and observational factors such as temperature, age, metallicity, rotation period, and viewing geometry (e.g., Marley and Robinson 2015; Biller 2017; Vos et al. 2017). These atmospheres also exhibit a rich diversity of chemical species, including water, methane, ammonia, carbon monoxide, carbon dioxide, and silicates (e.g., Kirkpatrick 2005; Cushing et al. 2006). Studying the chemistry and physics of these atmospheres provides insights not only into the atmospheric processes that govern their behavior, but also into the formation and evolution of these objects (e.g, Luhman 2012). In the long term, understanding these atmospheric conditions is essential for characterizing potential habitable worlds with next-generation observatories and for contextualizing the uniqueness of our own.

We introduce SEDA (Spectral Energy Distribution Analyzer), a versatile open-source Python package for forward modeling of observations of ultracool objects, such as brown dwarfs, directly imaged exoplanets, and low-mass stars. This workflow employs a Bayesian framework based on pre-computed, self-consistent atmospheric models. SEDA also incorporates a broad set of empirical techniques and auxiliary tools for spectral energy distribution (SED) analysis, as described in Functionality. Overall, SEDA enables the inference and analysis of atmospheric composition, cloud properties, and fundamental physical parameters of ultracool objects from observational data.

## Statement of Need

The James Webb Space Telescope (JWST) is revolutionizing our understanding of extrasolar atmospheres by providing high-quality observations of exoplanets and brown dwarfs. These data will synergize with observations from next-generation facilities such as the Nancy Grace Roman Space Telescope, the Habitable Worlds Observatory, the Vera C. Rubin Observatory, and 30-m class telescopes. Furthermore, a large volume of multiwavelength data is already available from wide-field photometric surveys (e.g., SDSS, York et al. 2000; and Pan-STARRS, Chambers et al. 2016), all-sky photometric surveys (e.g., Gaia, Gaia Collaboration et al. 2020; 2MASS, Skrutskie et al. 2006; and WISE, Cutri et al. 2013), targeted spectroscopic programs (e.g., SpeX Prism Library, Burgasser 2014; MOCAdb, Gagné et al. 2026), and both large (e.g., LAMOST, Luo et al. 2015; and APOGEE, Abdurro'uf et al. 2022) and all-sky (SPHEREx, Doré et al. 2014) spectroscopic surveys.

To fully exploit these extensive datasets, it is necessary to develop techniques and software capable of managing heterogeneous, multiwavelength observations (see State of the Field). `SEDA` contributes to this need by providing a framework for the analysis and forward modeling of photometric and spectroscopic multiwavelength data either separately or jointly, using spectra of varying resolution and/or overlapping wavelength coverage, as well as photometry from a broad range of filter systems compiled by the Spanish Virtual Observatory (SVO) Filter Profile Service[1]. In addition to its modeling capabilities, the package incorporates a growing set of empirical techniques commonly used in the literature, offering a unified and flexible framework for both model-based and empirical SED analysis (see Functionality).

## State of the Field

Several software packages enable forward modeling of ultracool dwarf data, including `VOSA` (Bayo et al. 2008), `Starfish` (Czekala et al. 2015), `SEDkit` (Filippazzo et al. 2015), `SPLAT` (Burgasser and Splat Development Team 2017), `species` (Stolker et al. 2020), `ForMoSA` (Petrus et al. 2023), and `PICASO` (JWST Transiting Exoplanet Community Early Release Science Team et al. 2023). Except for `VOSA`, these packages primarily focus on spectroscopic data, with limited ability to incorporate photometry in the model fitting. To our knowledge, only `species` can forward model SEDs constructed from photometry alone or combined with spectra, but its empirical analysis capabilities are relatively limited. In contrast, `SPLAT` provides more empirical analysis tools similar to those in `SEDA`, but it is restricted to modeling spectroscopic observations. `SEDA` complements these packages by supporting the modeling of spectrophotometric SEDs and offering extensive empirical analysis techniques, facilitating the study of brown dwarfs and directly imaged exoplanets.

[1] https://svo2.cab.inta-csic.es/theory/fps/

## Functionality

Figure 1 illustrates the core functionality and key tools of `SEDA`.

`SEDA` forward modeling employs two complementary approaches: a Bayesian framework with nested sampling to construct posterior distributions and a chi-square minimization method to find the best-fitting models (see Mathematics). The code supports both widely used and state-of-the-art atmospheric model grids, namely Lacy (2026, in preparation), Sonora Diamondback (Morley et al. 2024), Sonora Elf Owl (Mukherjee et al. 2024), Lacy and Burrows (2023), Sonora Cholla (Karalidi et al. 2021), Sonora Bobcat (Marley et al. 2021), ATMO 2020 (Phillips et al. 2020), BT-Settl (Allard et al. 2012), and Saumon and Marley (2008). Collectively, these models span a broad range of atmospheric and physical conditions, from cloud-free to cloudy atmospheres; equilibrium and non-equilibrium chemistry; temperatures from the coldest brown dwarfs through hot and temperate exoplanets to low-mass stars ($\sim$ 300–4200 K); and surface gravities ($\log g$=3.0–5.5) appropriate for both young and old ultracool objects. `SEDA` also allows users to easily incorporate additional atmospheric model grids. Fitting these model grids to observational data, the code infers chemical and physical parameters defined as free parameters in the grids, along with additional quantities such as bolometric luminosity and radius, when the distance is known (see Mathematics).

Empirical analyses of SEDs include measurements of the strengths of key spectral features produced by gas and grain species using literature-defined or user-defined spectral indices, i.e., flux ratios that quantify the depth of specific spectral signatures. These indices are applied to mid-infrared features (e.g., Cushing et al. 2006; Suárez and Metchev 2022, 2023), as well as to near-infrared diagnostics designed to identify potential variable objects (e.g., Ashraf et al. 2022; Oliveros-Gomez et al. 2022, 2024). Additional empirical analyses include the estimation of fundamental parameters such as bolometric luminosity and radius, and the application of published relationships between fundamental parameters, including temperature-spectral type, luminosity-spectral type, and inclination-cloudiness-color trends (e.g., Filippazzo et al. 2015; Vos et al. 2017; Kirkpatrick et al. 2021; Suárez et al. 2023).

In addition, `SEDA` provides several auxiliary tools that can be used independently, including: (*i*) the calculation of synthetic photometry from spectra for any filter available through SVO, (*ii*) a model interpolator that generates synthetic spectra for any parameter combinations within the grid coverage of available atmospheric models, (*iii*) a model examiner that enables inspection of atmospheric model properties, such as basic and free parameters, parameter coverage, resolving power, and spectral resolution, and (*iv*) access to 113 Spitzer IRS mid-infrared spectra of brown dwarfs from Suárez and Metchev (2022), together with associated target properties, observing logs, and measured spectral indices.




Figure 1: Overview of the SEDA workflow, illustrating key components such as forward modeling, empirical analysis, and additional useful tools.

## Mathematics

For model fitting, as described in Functionality, the code adopts two approaches:

### 1. Chi-square Minimization

The code uses the following weighted reduced chi-square ($\chi_r^2$) definition:

$$\chi_r^2 = \frac{1}{N - n_p} \sum_{i=1}^{N} w_i \left( \frac{F_{\mathrm{obs},i} - \alpha F_{\mathrm{mod},i}}{\sigma_i} \right)^2 \tag{1}$$

Here, $N$ is the number of fitted data points, $n_p$ is the number of free model parameters (so $N - n_p$ are the degrees of freedom), $w_i$ is the weight assigned to each data point to balance the relative contributions of datasets with different numbers of data points, $F_{\mathrm{obs},i}$ and $\sigma_i$ are the observed fluxes and their uncertainties, $F_{\mathrm{mod},i}$ are the model fluxes, and $\alpha$ is an unknown geometric dilution factor equal to $(R/d)^2$, where $R$ is the object radius and $d$ is its distance.

The code provides several options for assigning the weights $w_i$, which are designed to compensate for differences in the number of data points among datasets and to ensure no dataset has a negligible contribution to the fit (e.g., photometric data versus spectra, or low-resolution versus high-resolution spectra). These options include: ($i$) assigning each dataset a weight equal to the inverse of its total number of data points, so that all datasets contribute the same, even if they have very different numbers of points; and ($ii$) weighting spectroscopic data points by their wavelength resolution (i.e., wavelength step) and photometric data points by the filter effective width, such that broader filters or low-resolution spectra receive larger weights than high-resolution spectra.

To determine the value of $\alpha$ that minimizes $\chi_r^2$, the code can either use the non-linear least-squares minimization python package `LMFIT` (Newville et al. 2014) or compute the analytic solution obtained by setting $\frac{\partial \chi_r^2}{\partial \alpha} = 0$, which is given by:

$$\alpha = \frac{\sum_{i=1}^{N} w_i \frac{F_{\mathrm{obs},i} F_{\mathrm{mod},i}}{\sigma_i^2}}{\sum_{i=1}^{N} w_i \frac{F_{\mathrm{mod},i}^2}{\sigma_i^2}} \tag{2}$$

### 2. Nested Sampling

For Bayesian model fitting, the code adopts the weighted Gaussian log-likelihood ($\ln \mathcal{L}$):

$$\ln \mathcal{L} = -\frac{1}{2} \sum_{i=1}^{N} \left[ w_i \frac{(F_{\mathrm{obs},i} - \alpha F_{\mathrm{mod},i})^2}{\sigma_i^2} + \ln\left(2\pi\, \sigma_i^2\right) \right] \tag{3}$$

with all parameters defined as in Equation 1. The code wraps the model interpolator with the dynamic nested sampling package `DYNESTY` (Speagle 2020) to construct posteriors. During sampling, each generated synthetic spectrum is scaled using either $(R/d)^2$ if distance is known, or the factor that maximizes $\ln \mathcal{L}$, which is the same that minimizes $\chi_r^2$ (Equation 2).

The scaling factor $\alpha$ is used to derive the radius from the equation above, independently of evolutionary models, provided that the distance is known. The observed data are complemented with the best-fitting model to construct a full hybrid SED, which is then integrated to determine the bolometric luminosity, if the distance is available.

# Documentation

`SEDA` is hosted on GitHub[2] and the most up-to-date documentation is available on Read The Docs[3]. The documentation provides installation instructions, an overview of the code and its principal modules, a description of the code workflow, a list of useful tools, and details on the available atmospheric models. The repository also provides Jupyter notebook tutorials demonstrating typical scientific workflows.

The documentation is actively maintained and includes an API reference describing all functions and classes, with basic example provided where applicable. In addition, the documentation contains a frequently asked questions section and guidance on contributing to the code or submitting feedback.

# Software Design

`SEDA` v1.0 accompanies this publication and represents a stable, well-tested release following multiple pre-releases used for development and community feedback. The software is designed for flexible analysis of SEDs, centered on forward modeling and complemented by empirical analysis tools (see Functionality and Mathematics). The code includes automated tests to verify core functionality and has been successfully tested with Python 3.9 or later (latest tested: Python 3.14) on Linux, Windows, and macOS.

[2] https://github.com/suarezgenaro/seda
[3] https://seda.readthedocs.io/

## Research Impact

The foundation of `SEDA` was introduced in Suárez et al. (2021), which implemented only a chi-square minimization approach. In this work, we substantially expand upon that initial framework and release a significantly enhanced version of the software to the community. `SEDA` has already been used in at least 14 publications (including published, submitted, or in preparation works) (e.g., Suárez et al. 2025; Kiman et al. 2026; Rothermich et al. 2026; L'Heureux et al. 2026; Lam et al. 2026).

## Future Developments

Among several planned additions that will further increase the functionality of the code, future releases will (*i*) enable measurements of projected rotational velocity ($v \sin i$) and radial velocity (RV) by incorporating rotational broadening and Doppler shifting into atmospheric model spectra, as well as inference of viewing geometry based on $v \sin i$, rotation period, and radius constraints (e.g., Vos et al. 2017), (*ii*) derive physical parameters such as mass, age, surface gravity, and temperature by combining atmospheric model-fitting results with evolutionary models (e.g., Suárez et al. 2021), and (*iii*) implement cross-correlation analyses to identify molecular species in medium- to high-resolution spectroscopic observations (e.g., Petit dit de la Roche et al. 2018).

We welcome feedback, suggestions, and contributions from the community to help improve `SEDA` in future releases.

## AI Usage Disclosure

Generative AI tools (ChatGPT and Gemini) were used in a limited capacity to help polish the language of the documentation and to provide occasional coding suggestions or simplifications. All software design, implementation, and scientific decisions were developed by the authors. Any AI-generated suggestions were reviewed, verified, and modified by the authors before inclusion.

## Acknowledgements

GS and JKF acknowledge funding support from JWST-GO-02124.001-A as well as NASA XRP Award 80NSSC22K0142 and NSF Award 1909776. ML and JMV acknowledge support from a Royal Society - Research Ireland University Research Fellowship (URF/1/221932, RF/ERE/221108). ML acknowledges support from Trinity College Dublin via a Trinity Research Doctoral Award.

## References


Abdurro'uf, Katherine Accetta, Conny Aerts, et al. 2022. "The Seventeenth Data Release of the Sloan Digital Sky Surveys: Complete Release of MaNGA, MaStar, and APOGEE-2 Data." *Astrophysical Journal, Supplement* 259 (2): 35. https://doi.org/10.3847/1538-4365/ac4414.

Allard, F., D. Homeier, and B. Freytag. 2012. "Models of very-low-mass stars, brown dwarfs and exoplanets." *Philosophical Transactions of the Royal Society of London Series A* 370 (June): 2765–77. https://doi.org/10.1098/rsta.2011.0269.

Ashraf, Afra, Daniella C. Bardalez Gagliuffi, Elena Manjavacas, Johanna M. Vos, Claire Mechmann, and Jacqueline K. Faherty. 2022. "Disentangling the Signatures of Blended-light Atmospheres in L/T Transition Brown Dwarfs." *Astrophysical Journal* 934 (2): 178. https://doi.org/10.3847/1538-4357/ac7aab.

Bayo, A., C. Rodrigo, D. Barrado Y Navascués, et al. 2008. "VOSA: virtual observatory SED analyzer. An application to the Collinder 69 open cluster." *Astronomy & Astrophysics* 492 (December): 277–87. https://doi.org/10.1051/0004-6361:200810395.

Biller, Beth. 2017. "The time domain for brown dwarfs and directly imaged giant exoplanets: the power of variability monitoring." *The Astronomical Review* 13 (1): 1–27. https://doi.org/10.1080/21672857.2017.1303105.

Burgasser, A. J., and Splat Development Team. 2017. "The SpeX Prism Library Analysis Toolkit (SPLAT): A Data Curation Model." *Astronomical Society of India Conference Series*, Astronomical society of india conference series, vol. 14 (January): 7–12. https://arxiv.org/abs/1707.00062.

Burgasser, Adam J. 2014. "The SpeX Prism Library: 1000+ low-resolution, near-infrared spectra of ultracool M, L, T and Y dwarfs." *Astronomical Society of India Conference Series*, Astronomical society of india conference series, vol. 11 (January): 7–16. https://arxiv.org/abs/1406.4887.

Chambers, K. C., E. A. Magnier, N. Metcalfe, et al. 2016. "The Pan-STARRS1 Surveys." *ArXiv e-Prints*, December. https://arxiv.org/abs/1612.05560.

Cushing, M. C., T. L. Roellig, M. S. Marley, et al. 2006. "A Spitzer Infrared Spectrograph Spectral Sequence of M, L, and T Dwarfs." *Astrophysical Journal* 648 (September): 614–28. https://doi.org/10.1086/505637.

Cutri, R. M., E. L. Wright, T. Conrow, et al. 2013. "VizieR Online Data Catalog: AllWISE Data Release (Cutri+ 2013)." *yCat* 2328 (February): 0.

Czekala, Ian, Sean M. Andrews, Kaisey S. Mandel, David W. Hogg, and Gregory M. Green. 2015. "Constructing a Flexible Likelihood Function for Spectroscopic Inference." *Astrophysical Journal* 812 (2): 128. https://doi.org/10.1088/0004-637X/812/2/128.

Doré, Olivier, Jamie Bock, Matthew Ashby, et al. 2014. "Cosmology with the SPHEREX All-Sky Spectral Survey." *arXiv e-Prints*, December, arXiv:1412.4872. https://doi.org/10.48550/arXiv.1412.4872.

Filippazzo, Joseph C., Emily L. Rice, Jacqueline Faherty, Kelle L. Cruz, Mollie M. Van Gordon, and Dagny L. Looper. 2015. "Fundamental Parameters and Spectral Energy Distributions of Young and Field Age Objects with Masses Spanning the Stellar to Planetary Regime." *Astrophysical Journal* 810 (2): 158. https://doi.org/10.1088/0004-637X/810/2/158.

Gagné, Jonathan, Leslie Moranta, Jacqueline K. Faherty, et al. 2026. "The Montreal Open Clusters and Associations (MOCA) Database: A Census of Nearby Associations, Open Clusters, and Young Substellar Objects within 500 pc of the Sun." *arXiv e-Prints*, February, arXiv:2602.15695. https://arxiv.org/abs/2602.15695.

Gaia Collaboration, A. G. A. Brown, A. Vallenari, et al. 2020. "Gaia Early Data Release 3: Summary of the contents and survey properties." *arXiv e-Prints*, December, arXiv:2012.01533. https://arxiv.org/abs/2012.01533.

JWST Transiting Exoplanet Community Early Release Science Team, Eva-Maria Ahrer, Lili Alderson, et al. 2023. "Identification of carbon dioxide in an exoplanet atmosphere." *Nature* 614 (7949): 649–52. https://doi.org/10.1038/s41586-022-05269-w.

Karalidi, Theodora, Mark Marley, Jonathan J. Fortney, et al. 2021. "The Sonora Substellar Atmosphere Models. II. Cholla: A Grid of Cloud-free, Solar Metallicity Models in Chemical Disequilibrium for the JWST Era." *Astrophysical Journal* 923 (2): 269. https://doi.org/10.3847/1538-4357/ac3140.

Kiman, Rocio, Charles A. Beichman, Azul Ruiz Diaz, et al. 2026. "The Diversity of Cold Worlds: Age and Characterization of the Exoplanet COCONUTS-2 b." *Astronomical Journal* 171 (2): 60. https://doi.org/10.3847/1538-3881/ae230f.

Kirkpatrick, J. Davy. 2005. "New Spectral Types L and T." *Annual Review of Astronomy and Astrophysics* 43 (1): 195–245. https://doi.org/10.1146/annurev.astro.42.053102.134017.

Kirkpatrick, J. Davy, Christopher R. Gelino, Jacqueline K. Faherty, et al. 2021.

"The Field Substellar Mass Function Based on the Full-sky 20 pc Census of 525 L, T, and Y Dwarfs." *Astrophysical Journals* 253 (1): 7. https://doi.org/10.3847/1538-4365/abd107.

L'Heureux, J., G. Suárez, M. V. Vos, Metchev S., J. Faherty, and et al. 2026. "Inspecting Cloudy Substellar Atmospheres with JWST MIRI Synthetic Magnitudes from Spitzer Mid-infrared Spectra." *Astrophysical Journal* submitted.

Lacy, Brianna, and Adam Burrows. 2023. "Self-consistent Models of Y Dwarf Atmospheres with Water Clouds and Disequilibrium Chemistry." *Astrophysical Journal* 950 (1): 8. https://doi.org/10.3847/1538-4357/acc8cb.

Lam, M., J. M. Vos, G. Suárez, C-C. Bickle T. P. Hsu, and et al. 2026. "Clouds with a silicate lining: Using JWST spectra to probe atmospheric diversity in young AB Dor L dwarfs." *Astrophysical Journal* submitted.

Luhman, Kevin L. 2012. "The Formation and Early Evolution of Low-Mass Stars and Brown Dwarfs." *Annual Review of Astronomy and Astrophysics* 50 (September): 65–106. https://doi.org/10.1146/annurev-astro-081811-125528.

Luo, A.-Li, Yong-Heng Zhao, Gang Zhao, et al. 2015. "The first data release (DR1) of the LAMOST regular survey." *Research in Astronomy and Astrophysics* 15 (8): 1095. https://doi.org/10.1088/1674-4527/15/8/002.

Marley, M. S., and T. D. Robinson. 2015. "On the Cool Side: Modeling the Atmospheres of Brown Dwarfs and Giant Planets." *Annual Review of Astronomy and Astrophysics* 53 (August): 279–323. https://doi.org/10.1146/annurev-astro-082214-122522.

Marley, Mark S., Didier Saumon, Channon Visscher, et al. 2021. "The Sonora Brown Dwarf Atmosphere and Evolution Models. I. Model Description and Application to Cloudless Atmospheres in Rainout Chemical Equilibrium." *Astrophysical Journal* 920 (2): 85. https://doi.org/10.3847/1538-4357/ac141d.

Morley, Caroline V., Sagnick Mukherjee, Mark S. Marley, et al. 2024. "The Sonora Substellar Atmosphere Models. III. Diamondback: Atmospheric Properties, Spectra, and Evolution for Warm Cloudy Substellar Objects." *Astrophysical Journal* 975 (1): 59. https://doi.org/10.3847/1538-4357/ad71d5.

Mukherjee, Sagnick, Jonathan J. Fortney, Caroline V. Morley, et al. 2024. "The Sonora Substellar Atmosphere Models. IV. Elf Owl: Atmospheric Mixing and Chemical Disequilibrium with Varying Metallicity and C/O Ratios."

*arXiv e-Prints*, February, arXiv:2402.00756. https://doi.org/10.48550/arXiv.2402.00756.

Newville, Matthew, Till Stensitzki, Daniel B. Allen, and Antonino Ingargiola. 2014. *LMFIT: Non-Linear Least-Square Minimization and Curve-Fitting for Python*. V. 0.8.0. Zenodo, released September. https://doi.org/10.5281/zenodo.11813.

Oliveros-Gomez, Natalia, Elena Manjavacas, Afra Ashraf, et al. 2022. "Informed Systematic Method to Identify Variable Mid- and Late-T Dwarfs." *Astrophysical Journal* 939 (2): 72. https://doi.org/10.3847/1538-4357/ac96f2.

Oliveros-Gomez, Natalia, Elena Manjavacas, Daniella C. Bardalez Gagliuffi, Theodora Karalidi, Johanna M. Vos, and Jacqueline K. Faherty. 2024. "An Informed and Systematic Method to Identify Variable Mid-L Dwarfs." *Astrophysical Journal* 967 (2): 149. https://doi.org/10.3847/1538-4357/ad39e4.

Petit dit de la Roche, D. J. M., H. J. Hoeijmakers, and I. A. G. Snellen. 2018. "Molecule mapping of HR8799b using OSIRIS on Keck. Strong detection of water and carbon monoxide, but no methane." *Astronomy & Astrophysics* 616 (August): A146. https://doi.org/10.1051/0004-6361/201833384.

Petrus, S., G. Chauvin, M. Bonnefoy, et al. 2023. "X-SHYNE: X-shooter spectra of young exoplanet analogs. I. A medium-resolution 0.65-2.5 $\mu$m one-shot spectrum of VHS 1256−1257 b." *Astronomy & Astrophysics* 670 (February): L9. https://doi.org/10.1051/0004-6361/202244494.

Phillips, M. W., P. Tremblin, I. Baraffe, et al. 2020. "A new set of atmosphere and evolution models for cool T-Y brown dwarfs and giant exoplanets." *Astronomy & Astrophysics* 637 (May): A38. https://doi.org/10.1051/0004-6361/201937381.

Rothermich, Austin, Jacqueline K. Faherty, Ben Burningham, Catherine Manea, Emily Calamari, and et al. 2026. "The Buddy System: Characterizing the benchmark system containing the T0 brown dwarf CWISE J210640.16+250729.0 using JWST." *Astrophysical Journal* submitted.

Saumon, D., and Mark S. Marley. 2008. "The Evolution of L and T Dwarfs in Color-Magnitude Diagrams." *Astrophysical Journal* 689 (2): 1327–44. https://doi.org/10.1086/592734.

Skrutskie, M. F., R. M. Cutri, R. Stiening, et al. 2006. "The Two Micron All Sky Survey (2MASS)." *Astronomical Journal* 131 (February): 1163–83. https://doi.org/10.1086/498708.

Speagle, Joshua S. 2020. "DYNESTY: a dynamic nested sampling package for estimating Bayesian posteriors and evidences." *Monthly Notices of the Royal Astronomical Society* 493 (3): 3132–58. https://doi.org/10.1093/mnras/staa278.

Stolker, T., S. P. Quanz, K. O. Todorov, et al. 2020. "MIRACLES: atmospheric characterization of directly imaged planets and substellar companions at 4-5 $\mu$m. I. Photometric analysis of $\beta$ Pic b, HIP 65426 b, PZ Tel B, and HD 206893 B." *Astronomy & Astrophysics* 635 (March): A182. https://doi.org/10.1051/0004-6361/201937159.

Suárez, Genaro, Jacqueline K. Faherty, Ben Burningham, et al. 2025. "Diversity of Cold Worlds: Predicted Near-to-mid-infrared Spectral Signatures of a Cold Brown Dwarf with Potential Auroral Heating." *Astrophysical Journal* 993 (2): 165. https://doi.org/10.3847/1538-4357/ae0e6a.

Suárez, Genaro, and Stanimir Metchev. 2022. "Ultracool dwarfs observed with the Spitzer infrared spectrograph - II. Emergence and sedimentation of silicate clouds in L dwarfs, and analysis of the full M5-T9 field dwarf spectroscopic sample." *Monthly Notices of the Royal Astronomical Society* 513 (4): 5701–26. https://doi.org/10.1093/mnras/stac1205.

Suárez, Genaro, and Stanimir Metchev. 2023. "Ultracool dwarfs observed with the Spitzer Infrared Spectrograph - III. Dust grains in young L dwarf atmospheres are heavier." *Monthly Notices of the Royal Astronomical Society* 523 (3): 4739–47. https://doi.org/10.1093/mnras/stad1711.

Suárez, Genaro, Stanimir Metchev, Sandy K. Leggett, Didier Saumon, and Mark S. Marley. 2021. "Ultracool Dwarfs Observed with the Spitzer Infrared Spectrograph. I. An Accurate Look at the L-to-T Transition at 300 Myr from Optical Through Mid-infrared Spectrophotometry." *Astrophysical Journal* 920 (2): 99. https://doi.org/10.3847/1538-4357/ac1418.

Suárez, Genaro, Johanna M. Vos, Stanimir Metchev, Jacqueline K. Faherty, and Kelle Cruz. 2023. "Ultracool Dwarfs Observed with the Spitzer Infrared Spectrograph: Equatorial Latitudes in L Dwarf Atmospheres Are Cloudier." *Astrophysical Journal, Letters* 954 (1): L6. https://doi.org/10.3847/2041-8213/acec4b.

Vos, Johanna M., Katelyn N. Allers, and Beth A. Biller. 2017. "The Viewing Geometry of Brown Dwarfs Influences Their Observed Colors and Variability Amplitudes." *Astrophysical Journal* 842 (2): 78. https://doi.org/10.3847/1538-4357/aa73cf.

York, Donald G., J. Adelman, Jr. Anderson John E., et al. 2000. "The Sloan Digital Sky Survey: Technical Summary." *Astronomical Journal* 120

(September): 1579–87. https://doi.org/10.1086/301513.