

In [1]: *# Importing the necessary libraries*

```
import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
import seaborn as sns

from scipy.stats import ttest_ind, f_oneway, levene, kruskal, shapiro, chi2_
from statsmodels.graphics.gofplots import qqplot

import warnings
warnings.filterwarnings("ignore")
```

In [2]: *# converting data into dataframe*

```
yulu = pd.read_csv('bike_sharing.csv')
```

In [3]: *# making an copy of the dataset*

```
df = yulu.copy()
```

In [4]: *# Top 5 rows of the dataframe*

```
df.head()
```

Out[4]:

	datetime	season	holiday	workingday	weather	temp	atemp	humidity	windspeed	casu
0	2011-01-01 00:00:00	1	0	0	1	9.84	14.395	81	0.0	
1	2011-01-01 01:00:00	1	0	0	1	9.02	13.635	80	0.0	
2	2011-01-01 02:00:00	1	0	0	1	9.02	13.635	80	0.0	
3	2011-01-01 03:00:00	1	0	0	1	9.84	14.395	75	0.0	
4	2011-01-01 04:00:00	1	0	0	1	9.84	14.395	75	0.0	

In [5]: *# No of rows and columns*

```
df.shape
```

Out[5]: (10886, 12)

```
In [6]: # Checking of null values
```

```
df.isna().sum()
```

```
Out[6]: datetime    0
        season      0
        holiday     0
        workingday  0
        weather     0
        temp        0
        atemp       0
        humidity    0
        windspeed   0
        casual      0
        registered  0
        count       0
        dtype: int64
```

There are totally 10886 rows and 12 columns in the data

The data does not contain any nulls, thus no need of handling the missing data.

```
In [7]: # Duplicate values check
```

```
df.duplicated().sum()
```

```
Out[7]: 0
```

```
In [8]: # skewness of each column
```

```
df.skew(numeric_only = True)
```

```
Out[8]: season      -0.007076
        holiday     5.660517
        workingday  -0.776163
        weather     1.243484
        temp        0.003691
        atemp       -0.102560
        humidity    -0.086335
        windspeed   0.588767
        casual      2.495748
        registered  1.524805
        count       1.242066
        dtype: float64
```

Skewness Analysis of Variables

Symmetrical Majority:

- The majority of the variables, including 'season' and 'temp', exhibit skewness values close to zero, suggesting relatively symmetrical distributions.

Positive Skewness Insights:

- Variables such as 'holiday', 'weather', 'windspeed', 'casual', 'registered', and 'count' demonstrate positive skewness, pointing to a concentration of lower values and a rightward skew in their distributions.

Negative Skewness Observations:

- In contrast, 'workingday', 'atemp', and 'humidity' exhibit negative skewness, implying a concentration of higher values and a leftward skew in their distributions.

In [9]: *# Uniques values of each columns*

```
df.nunique()
```

```
Out[9]: datetime      10886
season              4
holiday            2
workingday         2
weather            4
temp              49
atemp             60
humidity          89
windspeed         28
casual            309
registered        731
count             822
dtype: int64
```

In [10]: *# data info*

```
df.info()
```

```
<class 'pandas.core.frame.DataFrame'>
RangeIndex: 10886 entries, 0 to 10885
Data columns (total 12 columns):
#   Column      Non-Null Count  Dtype
---  ---
0   datetime    10886 non-null  object
1   season      10886 non-null  int64
2   holiday     10886 non-null  int64
3   workingday  10886 non-null  int64
4   weather     10886 non-null  int64
5   temp        10886 non-null  float64
6   atemp       10886 non-null  float64
7   humidity    10886 non-null  int64
8   windspeed   10886 non-null  float64
9   casual      10886 non-null  int64
10  registered  10886 non-null  int64
11  count       10886 non-null  int64
dtypes: float64(3), int64(8), object(1)
memory usage: 1020.7+ KB
```

In [11]: *# count column is sum of casual and the registered users*

```
(df['casual'] + df['registered'] == df['count']).value_counts()
```

```
Out[11]: True      10886
Name: count, dtype: int64
```

```
In [12]: # converting the categorical columns into category

cat_col = ['season', 'holiday', 'workingday', 'weather']

for _ in cat_col:
    df[_] = df[_].astype('category')

df.info()
```

```
<class 'pandas.core.frame.DataFrame'>
RangeIndex: 10886 entries, 0 to 10885
Data columns (total 12 columns):
#   Column      Non-Null Count  Dtype
---  -
0   datetime    10886 non-null  object
1   season      10886 non-null  category
2   holiday     10886 non-null  category
3   workingday  10886 non-null  category
4   weather     10886 non-null  category
5   temp        10886 non-null  float64
6   atemp       10886 non-null  float64
7   humidity    10886 non-null  int64
8   windspeed   10886 non-null  float64
9   casual      10886 non-null  int64
10  registered  10886 non-null  int64
11  count       10886 non-null  int64
dtypes: category(4), float64(3), int64(4), object(1)
memory usage: 723.7+ KB
```

```
In [13]: # Converting datetime column into date time format

df['datetime'] = pd.to_datetime(df['datetime'])
df['datetime'].dtype
```

Out[13]: dtype('<M8[ns]')

```
In [14]: # Creating new columns from datetime and converting them to categories

df['year'] = df['datetime'].dt.year
df['month'] = df['datetime'].dt.month
df['day'] = df['datetime'].dt.day
df['hour'] = df['datetime'].dt.hour
```

```
In [15]: df.head(2)
```

Out[15]:

	datetime	season	holiday	workingday	weather	temp	atemp	humidity	windspeed	casu
0	2011-01-01 00:00:00	1	0	0	1	9.84	14.395	81	0.0	
1	2011-01-01 01:00:00	1	0	0	1	9.02	13.635	80	0.0	

```

In [16]: # replacing the number with category

# change of season
df['season'] = df['season'].replace({1: 'Spring', 2: 'Summer', 3: 'Fall', 4: 'Winter'})

# change of holiday
df['holiday'] = df['holiday'].replace({0: 'No', 1: 'Yes'})

# change of workingday
df['workingday'] = df['workingday'].replace({0: 'No', 1: 'Yes'})

# change of month
df['month'] = df['month'].replace({1: 'January',
                                   2: 'February',
                                   3: 'March',
                                   4: 'April',
                                   5: 'May',
                                   6: 'June',
                                   7: 'July',
                                   8: 'August',
                                   9: 'September',
                                   10: 'October',
                                   11: 'November',
                                   12: 'December'})

```

```

In [17]: df.describe().transpose()

```

Out[17]:

	count	mean	min	25%	50%	75%	max
datetime	10886	2011-12-27 05:56:22.399411968	2011-01-01 00:00:00	2011-07-02 07:15:00	2012-01-01 20:30:00	2012-07-01 12:45:00	2012-12-19 23:00:00
temp	10886.0	20.23086	0.82	13.94	20.5	26.24	41.0
atemp	10886.0	23.655084	0.76	16.665	24.24	31.06	45.455
humidity	10886.0	61.88646	0.0	47.0	62.0	77.0	100.0
windspeed	10886.0	12.799395	0.0	7.0015	12.998	16.9979	56.9969
casual	10886.0	36.021955	0.0	4.0	17.0	49.0	367.0
registered	10886.0	155.552177	0.0	36.0	118.0	222.0	886.0
count	10886.0	191.574132	1.0	42.0	145.0	284.0	977.0
year	10886.0	2011.501929	2011.0	2011.0	2012.0	2012.0	2012.0
day	10886.0	9.992559	1.0	5.0	10.0	15.0	19.0
hour	10886.0	11.541613	0.0	6.0	12.0	18.0	23.0

```
In [18]: df.describe(include = 'category').transpose()
```

```
Out[18]:
```

	count	unique	top	freq
season	10886	4	Winter	2734
holiday	10886	2	No	10575
workingday	10886	2	Yes	7412
weather	10886	4	1	7192

Overview and Feature Patterns

Temporal and Numerical Composition:

- The dataset encompasses both datetime information and various numerical features associated with bike rentals. The observations span from January 1, 2011, to December 19, 2012.

Diverse Numerical Feature Characteristics:

- Numerical features such as temperature, humidity, windspeed, and counts of casual and registered bike rentals show diverse ranges and distributions, highlighting the variability in rental patterns across different conditions.

Temporal Patterns and Concentrations:


- Observations on the year, day, and hour variables indicate temporal patterns, with a concentration in 2011 and 2012, a mean day value around 10, and an hourly distribution ranging from 0 to 23.

Outlier Detection

```
In [19]: plt.figure(figsize=(15, 12))
sns.set(style="white")

for i, column in enumerate(cat_col,1):
    plt.subplot(2, 2, i)
    sns.boxplot(x=column, y='count', data=df, color="#29B6F6")
    plt.title(f'Boxplot for {column}')

plt.tight_layout()
plt.show()
```



Outlier Analysis

Outliers in Different Seasons:

- In spring and winter, there are more unusual values in the data compared to other seasons.

Weather Outliers:

- Category 3 weather has a lot of unusual values, while category 4 weather doesn't have any.

Working Days vs. Holidays:

- On regular working days, there are more unusual values in the data than on holidays. This suggests some unexpected patterns during typical workdays that might need a closer look.

Univariate Analysis

```
In [20]: # Time span of data
time_span = df['datetime'].max() - df['datetime'].min()
time_span
```

Out[20]: Timedelta('718 days 23:00:00')

```
In [21]: df.columns
```

Out[21]: Index(['datetime', 'season', 'holiday', 'workingday', 'weather', 'temp', 'atemp', 'humidity', 'windspeed', 'casual', 'registered', 'count', 'year', 'month', 'day', 'hour'], dtype='object')

```
In [22]: # Season counts
df['season'].value_counts()
```

Out[22]: season
Winter 2734
Summer 2733
Fall 2733
Spring 2686
Name: count, dtype: int64

```
In [23]: # holiday counts
df['holiday'].value_counts()
```

Out[23]: holiday
No 10575
Yes 311
Name: count, dtype: int64

```
In [24]: # workingday counts
df['workingday'].value_counts()
```

Out[24]: workingday
Yes 7412
No 3474
Name: count, dtype: int64

```
In [25]: # weather counts
df['weather'].value_counts()
```

Out[25]: weather
1 7192
2 2834
3 859
4 1
Name: count, dtype: int64


```
In [26]: # year counts
df['year'].value_counts()
```

```
Out[26]: year
2012     5464
2011     5422
Name: count, dtype: int64
```

```
In [27]: # month counts
df['month'].value_counts()
```

```
Out[27]: month
May           912
June          912
July          912
August        912
December      912
October       911
November      911
April         909
September     909
February      901
March         901
January       884
Name: count, dtype: int64
```

```
In [28]: # day counts
df['day'].value_counts().sort_index()
```


```
Out[28]: day
1         575
2         573
3         573
4         574
5         575
6         572
7         574
8         574
9         575
10        572
11        568
12        573
13        574
14        574
15        574
16        574
17        575
18        563
19        574
Name: count, dtype: int64
```

```
In [29]: # countplot on categories

plt.figure(figsize=(15, 12))
sns.set(style="darkgrid")

for i, column in enumerate(cat_col, 1):
    plt.subplot(2, 2, i)
    sns.countplot(x=column, data=df, color="#29B6F6", width=0.4)
    plt.title(f'Countplot for {column}')

plt.tight_layout()
plt.show()
```





In [30]: *# Function for histogram & boxplot on numerical columns*

```
def hist_box(column):  
    f, axs = plt.subplots(1, 2, figsize=(10, 5))  
    sns.set(style="darkgrid")  
  
    # Histogram  
    plt.subplot(1, 2, 1)  
    sns.histplot(df[column], bins=20, kde=True)  
    plt.title(f'Histogram for {column}')  
  
    # Boxplot  
    plt.subplot(1, 2, 2)  
    sns.boxplot(df[column], color="#29B6F6")  
    plt.title(f'Boxplot for {column}')  
  
    tabular_data = df[column].describe().reset_index()  
    tabular_data.columns = ['Statistic', 'Value']  
    display(tabular_data)  
  
    plt.tight_layout()  
    plt.show()
```


```
In [31]: num_col = ['temp', 'atemp', 'humidity', 'windspeed', 'casual', 'registered']

for column in num_col:
    hist_box(column)
```


	Statistic	Value
0	count	10886.00000
1	mean	20.23086
2	std	7.79159
3	min	0.82000
4	25%	13.94000
5	50%	20.50000
6	75%	26.24000
7	max	41.00000





	Statistic	Value
0	count	10886.000000
1	mean	23.655084
2	std	8.474601
3	min	0.760000
4	25%	16.665000
5	50%	24.240000
6	75%	31.060000
7	max	45.455000





	Statistic	Value
0	count	10886.000000
1	mean	61.886460
2	std	19.245033
3	min	0.000000
4	25%	47.000000
5	50%	62.000000
6	75%	77.000000
7	max	100.000000





	Statistic	Value
0	count	10886.000000
1	mean	12.799395
2	std	8.164537
3	min	0.000000
4	25%	7.001500
5	50%	12.998000
6	75%	16.997900
7	max	56.996900





	Statistic	Value
0	count	10886.000000
1	mean	36.021955
2	std	49.960477
3	min	0.000000
4	25%	4.000000
5	50%	17.000000
6	75%	49.000000
7	max	367.000000



	Statistic	Value
0	count	10886.000000
1	mean	155.552177
2	std	151.039033
3	min	0.000000
4	25%	36.000000
5	50%	118.000000
6	75%	222.000000
7	max	886.000000



	Statistic	Value
0	count	10886.000000
1	mean	191.574132
2	std	181.144454
3	min	1.000000
4	25%	42.000000
5	50%	145.000000
6	75%	284.000000
7	max	977.000000



Numerical column analysis

Temp:

- The 'temp' column shows a diverse temperature range (0.82 to 41.0), with a median of 20.5 and moderate variability around the mean of approximately 20.23 degrees Celsius.

Atemp

- The 'atemp' column displays a wide range of apparent temperatures (0.76 to 45.455), with a mean of approximately 23.66 and moderate variability around the median of 24.24.

Humidity

- The 'humidity' column depicts a range of humidity values (0 to 100), with an average around 61.89. The distribution shows moderate variability, from 47 at the 25th percentile to 77 at the 75th percentile, indicating diverse humidity levels in the dataset.

WindSpeed

- The 'windspeed' column displays a range of wind speeds from 0 to 56.9979, with a mean of approximately 12.80.

Casual

- The 'casual' column demonstrates a broad range of casual bike rental counts, with values spanning from 0 to 367. The distribution is positively skewed, as indicated by the mean (36.02) being less than the median (17.0).

Registered

- The 'registered' column showcases a diverse range of registered bike rental counts, ranging from 0 to 886. The distribution is positively skewed, evidenced by the mean (155.55) being less than the median (118.0).

Count

- The 'count' column reveals a wide range of total bike rental counts, varying from 1 to 977. The distribution is positively skewed, with a mean (191.57) greater than the median (145.0), indicating a concentration of lower values

Bivariate Analysis

In [32]: cat_col


Out[32]: ['season', 'holiday', 'workingday', 'weather']

In [33]: *# barplot of categories*

```
plt.figure(figsize=(15, 12))
sns.set(style="darkgrid")

for i, column in enumerate(cat_col,1):
    plt.subplot(2, 2, i)
    sns.barplot(x=column, y='count', data=df, color="#29B6F8", width = 0.3)
    plt.title(f'{column} based distribution of rentals')

plt.tight_layout()
plt.show()
```



In [34]: *# correlation analysis*


```
correlation_matrix = df[["atemp", "temp", "humidity", "windspeed", "casual"]
correlation_df = pd.DataFrame(correlation_matrix)
correlation_df
```

Out[34]:

	atemp	temp	humidity	windspeed	casual	registered	count
atemp	1.000000	0.984948	-0.043536	-0.057473	0.462067	0.314635	0.389784
temp	0.984948	1.000000	-0.064949	-0.017852	0.467097	0.318571	0.394454
humidity	-0.043536	-0.064949	1.000000	-0.318607	-0.348187	-0.265458	-0.317371
windspeed	-0.057473	-0.017852	-0.318607	1.000000	0.092276	0.091052	0.101369
casual	0.462067	0.467097	-0.348187	0.092276	1.000000	0.497250	0.690414
registered	0.314635	0.318571	-0.265458	0.091052	0.497250	1.000000	0.970948
count	0.389784	0.394454	-0.317371	0.101369	0.690414	0.970948	1.000000

In [35]: `# correlation chart`

```
plt.figure(figsize = (16, 10))
sns.heatmap(correlation_matrix, annot = True)
plt.show()
```



Correlation Analysis

Atemp:

- Strong positive correlation with 'temp' (0.98), indicating a close relationship.
- Moderate positive correlation with 'casual' (0.46) and 'registered' (0.31).
- Positive correlation with 'count' (0.39), suggesting a relationship with overall bike rentals.

Temp (Temperature):

- Highly correlated with 'atemp' (0.98), indicating a strong connection.
- Moderate positive correlation with 'casual' (0.47) and 'registered' (0.32).
- Positive correlation with 'count' (0.39), showing a relationship with overall bike rentals.

Humidity:

- Weak negative correlation with 'atemp' (-0.04) and 'temp' (-0.06).
- Moderate negative correlation with 'casual' (-0.35), 'registered' (-0.27), and 'count' (-0.32).
- Indicates a tendency for fewer bike rentals during higher humidity.

Windspeed:

- Weak negative correlation with 'atemp' (-0.06) and 'temp' (-0.02).
- Weak positive correlation with 'casual' (0.09), 'registered' (0.09), and 'count' (0.10).
- Suggests a subtle influence on bike rentals with increasing wind speed.

Casual (Casual Bike Rentals):

- Strong positive correlation with 'atemp' (0.46) and 'temp' (0.47).
- Moderate negative correlation with 'humidity' (-0.35) and positive correlation with 'windspeed' (0.09).
- Highly correlated with 'registered' (0.50) and 'count' (0.69), indicating a significant impact on overall rentals.

Registered (Registered Bike Rentals):

- Positive correlation with 'atemp' (0.31) and 'temp' (0.32).
- Negative correlation with 'humidity' (-0.27) and positive correlation with 'windspeed' (0.09).
- Highly correlated with 'casual' (0.50) and 'count' (0.97), emphasizing a substantial impact on overall rentals.

Count (Total Bike Rentals):

- Positive correlation with 'atemp' (0.39), 'temp' (0.39), and 'casual' (0.69).
- Negative correlation with 'humidity' (-0.32).
- Highly correlated with 'registered' (0.97), emphasizing the joint impact of casual and registered rentals on the overall count.

```
In [36]: # counts based on months

monthly_count = df.groupby('month')['count'].sum().reset_index()

monthly_count = monthly_count.sort_values(by='count', ascending=False)

monthly_count
```


Out[36]:

	month	count
6	June	220733
5	July	214617
1	August	213516
11	September	212529
10	October	207434
8	May	200147
9	November	176440
0	April	167402
2	December	160160
7	March	133501
3	February	99113
4	January	79884

```
In [37]: # rentals on monthly counts

plt.figure(figsize=(10, 6))
sns.barplot(x='month', y='count', data=monthly_count, palette='flare', width=0.8)

plt.title('Total Count by Month')
plt.xlabel('Month')
plt.ylabel('Total Count')
plt.show()
```



Monthly analysis on rentals

Peak Rental Months:

- June stands out as the peak month for bike rentals, with the highest count of 220,733, followed closely by July and August.

Seasonal Trend:

- Summer months (June, July, August) show higher bike rental counts, consistent with favorable weather conditions.

Off-Peak Rental Months:

- January, February, and March have notably lower bike rental counts, indicating potential off-peak periods, possibly influenced by colder weather or fewer outdoor activities.

Hypothesis Testing

Demand of bicycles on rent is the same on Weekdays & Weekends

Since we have two independent samples, we can go with Two Sample Independent T-Test.

Assumptions of Two Sample Independent T-Test :

- The data should be normal distributed
- variances of the two groups are equal

Let the Confidence interval be 95%, so significance (alpha) is 0.05

To check if the data is normal, we will go with Wilkin-ShapiroTest.

The test hypothesis for the Wilkin-Shapiro test are:

- Ho: Data is normally distributed
- Ha: Data is not normally distributed.

```
In [38]: np.random.seed(41)

df_subset = df.sample(100)["count"]

test_stat, p_val = shapiro(df_subset)

p_val
```

```
Out[38]: 2.6341072612012795e-07
```


Hence the p_values is lesser than the significance level, Null hypothesis can be rejected.

Therefore, the Data is not normally distributed.

QQ Plot analysis

In [39]: `# QQ plot`

```
qqplot(df['count'], line = 's')  
plt.show()
```



To check if the variances of two groups are equal. We will perform Levene's test

The Test hypotheses for Levene's test are:


- **Ho: The variances are equal.**
- **Ha: The variances are not equal.**

```
In [40]: working_day = df[df['workingday'] == 'Yes']['count']  
         holiday = df[df['workingday'] == 'No']['count']  
         levene_stat, p_val = levene(working_day, holiday)  
         p_val
```

Out[40]: 0.9437823280916695


```
In [41]: sns.kdeplot(data = df, x = 'count', hue = 'workingday')
```

```
Out[41]: <Axes: xlabel='count', ylabel='Density'>
```



```
In [61]: sns.histplot(data = df, x = 'count', hue = 'workingday')
```

```
Out[61]: <Axes: xlabel='count', ylabel='Count'>
```



Hence the p_values is greater than the significance level, Null hypothesis can be accepted.

Therefore, the variances are approximately equal.

Despite the data is not normally distributed according to both the Wilkin-ShapiroTest and qq-plot

It is important to highlight that the variances between the two groups are equal**

So we can proceed with the Two Sample Independent T-Test.

The hypothesis for the t-test are:

- **Ho: There is no significant difference between working and non-working days.**
- **Ha: There is a significant difference between working and non-working days.**

```
In [43]: ttest_stat, p_val = ttest_ind(working_day, holiday)
p_val
```

```
Out[43]: 0.22644804226361348
```

Hence the p_values is greater than the significance level, Null hypothesis can be accepted.

Therefore, There is no significant difference on bike rentals between working and non-working days.

```
In [44]: kruskal_stat, p_val = kruskal(working_day, holiday)
p_val
```

```
Out[44]: 0.9679113872727798
```

Hence the p_values is greater than the significance level, Null hypothesis can be accepted.

Therefore, There is no significant difference on bike rentals between working and non-working days.

Demand of bicycles on rent is the same for different Weather conditions

Since we have more than two categories now, so will use ANOVA here.

Assumptions for ANOVA are:

1. The population data should be normally distributed- The data is not normal as verified by **Wilkin-Shapiro test and the qqplot.**
2. The data points must be independent- This condition is satisfied.

3. Approximately equal variance within groups- This will be verified using **Levene's test**.

```
In [45]: # skewness of weather  
  
df.groupby('weather')['count'].skew()
```


```
Out[45]: weather  
1      1.139857  
2      1.294444  
3      2.187137  
4         NaN  
Name: count, dtype: float64
```

```
In [46]: # kurtosis test of weather  
  
df.groupby('weather')['count'].apply(lambda x: x.kurtosis())
```

```
Out[46]: weather  
1      0.964720  
2      1.588430  
3      6.003054  
4         NaN  
Name: count, dtype: float64
```


```
In [47]: sns.kdeplot(data = df, x = 'count', hue = 'weather')
```

```
Out[47]: <Axes: xlabel='count', ylabel='Density'>
```



```
In [48]: sns.histplot(data = df, x = 'count', hue = 'weather')
```

```
Out[48]: <Axes: xlabel='count', ylabel='Count'>
```



The Test hypothesis for Levene's test are:

- **Ho: The variances are equal.**
- **Ha: The variances are not equal.**

```
In [49]: weather1 = df[df['weather'] == 1]['count']  
weather2 = df[df['weather'] == 2]['count']  
weather3 = df[df['weather'] == 3]['count']  
weather4 = df[df['weather'] == 4]['count']  
  
levene_stat, p_val = levene(weather1, weather2, weather3, weather4)  
  
p_val
```

```
Out[49]: 3.504937946833238e-35
```

Hence the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, the variances are not equal.

Two of the three conditions of ANOVA are not met, **We will still perform ANOVA.**

Then We will also perform **Kruskal's test and compare the results.**

In case of any discrepancies, Kruskal's test results will be considered, since data does not met conditions of ANOVA.

The hypothesis for ANOVA are:

- **Ho: There is no significant difference between demand of bicycles for different Weather conditions.**
- **Ha: There is a significant difference between demand of bicycles for different Weather conditions.**

```
In [50]: anova_stat, p_val = f_oneway(weather1, weather2, weather3,weather4)
p_val
```

```
Out[50]: 5.482069475935669e-42
```

Hence the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, There is a significant difference between demand of bicycles for different Weather conditions.

Kruskal Test on weather

```
In [51]: kruskal_stat, p_val = kruskal(weather1, weather2, weather3,weather4)
p_val
```

```
Out[51]: 3.501611300708679e-44
```

Again the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, we can conclude that there is a significant difference between demand of bicycles for different Weather conditions.

Demand of bicycles on rent is the same for different Seasons

Here also we have more than two categories now, so will use ANOVA here.

Assumptions for ANOVA are:

1. The population data should be normally distributed- The data is not normal as verified by **Wilkin-Shapiro test and the qqplot.**
2. The data points must be independent- This condition is satisfied.

In [52]: *# skewness of seasons*

```
df.groupby('season')['count'].skew()
```

Out[52]: season
Spring 1.888056
Summer 1.003264
Fall 0.991495
Winter 1.172117
Name: count, dtype: float64


In [53]: *# kurtosis test of seasons*

```
df.groupby('weather')['count'].apply(lambda x: x.kurtosis())
```

Out[53]: weather
1 0.964720
2 1.588430
3 6.003054
4 NaN
Name: count, dtype: float64


In [54]: `sns.kdeplot(data = df, x = 'count', hue = 'season')`

Out[54]: <Axes: xlabel='count', ylabel='Density'>



```
In [55]: sns.histplot(data = df, x = 'count', hue = 'season')
```

```
Out[55]: <Axes: xlabel='count', ylabel='Count'>
```



The Test hypothesis for Levene's test are:

- Ho: The variances are equal.
- Ha: The variances are not equal.

```
In [56]: spring = df[df['season'] == 'Spring']['count']
summer = df[df['season'] == 'Summer']['count']
fall = df[df['season'] == 'Fall']['count']
winter = df[df['season'] == 'Winter']['count']

levene_stat, p_val = levene(spring, summer, fall, winter)

p_val
```

```
Out[56]: 1.0147116860043298e-118
```

Hence the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, the variances are not equal.

As like before, we still use both ANOVA and Kruskal's test, comparing the results.

If discrepancies arise, we'll rely on **Kruskal's test**, Since data does not met the conditions for ANOVA.

The hypothesis for ANOVA are:

- **Ho: There is no significant difference between demand of bicycles for different Seasons.**
- **Ha: There is a significant difference between demand of bicycles for different Seasons.**

```
In [57]: anova_stat, p_val = f_oneway(spring ,summer, fall, winter)

p_val
```

```
Out[57]: 6.164843386499654e-149
```

Hence the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, There is a significant difference between demand of bicycles for different Seasons.

Kruskal Test on season

```
In [58]: kruskal_stat, p_val = kruskal(spring ,summer, fall, winter)

p_val
```

```
Out[58]: 2.479008372608633e-151
```

Again the p_values is smaller than the significance level, Null hypothesis can be rejected.

Therefore, we can conclude that there is a significant difference between demand of bicycles for different Seasons.

Analysis of Weather Conditions Across Seasons using Chi-square Test

The hypothesis for the chi-square test are:

Ho: Season and Weather are independent of each other.

Ha: Season and Weather are dependent on each other.

```
In [59]: contingency_table = pd.crosstab(df['weather'], df['season'])

contingency_table
```

```
Out[59]:
```

	season	Spring	Summer	Fall	Winter
weather					
1	1759	1801	1930	1702	
2	715	708	604	807	
3	211	224	199	225	
4	1	0	0	0	

```
In [60]: chi2_contingency(contingency_table)
```

```
Out[60]: Chi2ContingencyResult(statistic=49.15865559689363, pvalue=1.5499250736864862e-07, dof=9, expected_freq=array([[1.77454639e+03, 1.80559765e+03, 1.80559765e+03, 1.80625831e+03],
        [6.99258130e+02, 7.11493845e+02, 7.11493845e+02, 7.11754180e+02],
        [2.11948742e+02, 2.15657450e+02, 2.15657450e+02, 2.15736359e+02],
        [2.46738931e-01, 2.51056403e-01, 2.51056403e-01, 2.51148264e-01]]))
```

Hence the $p_values(1.5499250736864862e-07)$ is smaller than the significance level, Null hypothesis can be rejected.

Therefore, we can conclude that Season and Weather are dependent on each other.

Strategic Recommendations for Yulu's Profitable Growth

Optimize Bike Distribution in Peak Months:

- Concentrate bike deployment efforts during peak months, especially in June, July, and August, to meet increased demand and capitalize on favorable weather conditions.

Seasonal Marketing Strategies:

- Tailor marketing efforts to leverage the seasonal trend, promoting Yulu's services more aggressively during summer months to attract a larger user base.

Enhance User Engagement in Off-Peak Months:

- Implement targeted promotional campaigns or discounts during off-peak months (e.g., January to March) to encourage increased bike rentals and maintain consistent revenue flow.

Weather-Responsive Pricing:

- Consider implementing dynamic pricing strategies that respond to weather conditions. For example, adjusting rental rates during extreme weather days to optimize revenue.

Diversify Revenue Streams:

- Explore additional revenue streams, such as partnerships, sponsorships, or offering premium membership services with added benefits, to diversify income sources and boost overall profitability.

Enhance User Experience:

- Invest in technology and infrastructure to improve the overall user experience, including app features, bike maintenance, and customer support, fostering loyalty and repeat business.

Optimize Bike Deployment on Working Days:

- Given the lack of significant differences in bike rentals between working and non-working days, consider adjusting bike deployment strategies to ensure optimal resource allocation throughout the week.

Adapt to Different Weather Conditions:

- Change promotions or discounts based on the weather. If it's rainy, for example, offer special deals to encourage more people to use the bikes.

Promote Bikes Differently in Each Season:

- Advertise the bikes differently in each season. For example, highlight summer promotions in June, July, and August when more people want to ride bikes.

Combine Season and Weather Plans:

- Plan bike availability based on both the season and the weather to make sure people have the bikes they need when they want them. For example, have more bikes available on sunny days in the summer.