

Multivariable Calculus

MATH 212 LECTURE NOTES

Subhadip Chowdhury, PhD



Spring 2022

Contents



I Vectors and Three Dimensional Geometry	1
1 Three Dimensional Coordinate Geometry	3
1.1 Coordinate Axes and Points in 3-space	3
1.2 Distance between two points	4
1.2.1 Equation of a Sphere	5
1.3 Sets of points described using coordinates	5
2 Vectors in 3D	6
2.1 Definition	6
2.2 Graphical representation of vectors	6
2.3 Vector Arithmetic	6
2.3.1 Addition	6
2.3.2 Scalar Multiplication	6
2.3.3 Subtraction	7
2.4 Symbolic representation of vectors	7
2.4.1 Position vector - components vs coordinates	7
2.5 Magnitude	8
2.6 Calculating components in 2D	8
2.7 Word problems	8
3 Dot Product of Vectors	9
3.1 Definition	9
3.1.1 Algebraic	9
3.1.2 Geometric	9
3.2 Basic properties	9
3.3 Angle between two vectors	9
3.4 Vector projections	10
4 Cross Product of Vectors	11
4.1 Definition	11
4.1.1 Geometric	11
4.1.2 Algebraic	11
4.2 Basic Properties	11
4.3 Cross product as area	12
4.3.1 Volume of a parallelepiped - Coplanarity	12
5 Lines and Planes	13
5.1 Lines in space	13
5.2 Planes in space	13
5.3 Types of problems	14
5.3.1 Finding a Plane	14
5.3.2 Finding a Line	14
5.4 Practice problems	14

6 Polar Coordinate System and Polar Curves	16
6.1 Definition	16
6.1.1 When r is negative	16
6.2 Converting between Polar and Cartesian Coordinates	17
6.3 Polar Curves	17
6.3.1 Symmetry	18
7 Vector Valued Functions and Space Curves	19
7.1 Different ways of parametrizing the same curve	20
7.2 3D Curves	20
7.2.1 Helix	20
7.3 Parametrization from equation of curve	21
8 Calculus of Vector-valued Functions	23
8.1 Derivative of a Vector-valued function	23
8.1.1 Differentiation Rules	23
8.1.2 Interpretation as Velocity	24
8.2 Integral of a vector-valued function - arc length	24
8.2.1 Arc Length Parameterization	25
8.3 Curvature	26
8.4 The Osculating Plane	27
8.5 Acceleration	29
II Multivariable Functions and their Derivatives	31
9 Functions of Two Variables	33
9.1 Examples of functions of two variables	33
9.2 Representations of two-variable functions	33
9.2.1 Numerical	33
9.2.2 Algebraic	33
9.2.3 Visual - Using Graphs	33
9.3 Analyzing Graphs using Cross-sections	34
9.4 Group Problems	34
10 Contour Plots - Level Curves and Level Surfaces	36
10.1 Level Curves	36
10.1.1 Definition	36
10.1.2 Basic Facts	36
10.2 Functions of three variables - Level surface	39
10.2.1 Level surfaces of $f(x,y,z) = x^2 + y^2 + z^2$	39
11 Partial Derivatives	41
11.1 Motivation	41
11.2 Definition	41
11.3 Partial derivative function	42
11.4 Higher order partials	42
11.5 Interpreting second partial derivatives	43
12 Tangent Plane and Linear Approximation	44
12.1 Tangent Plane	44
12.2 Linear Approximation	45

13 Directional Derivative and Gradient	46
13.1 Significance of the gradient	47
13.2 Gradient and Level Curves	47
14 Three Dimensional Gradient and the Tangent Plane	48
14.1 3D Gradient as the Normal vector	48
15 Chain Rule	50
15.1 Chain Rule diagram	50
15.2 More complicated diagram	51
16 Limit, Continuity, and Differentiability	53
16.1 Limit and Continuity	53
16.2 Continuity	55
16.3 Differentiability	55
III Application of Derivatives - Optimization	57
17 Local Optimization	59
17.1 How to detect a local extremum from the Contour Plot?	59
17.2 Critical Points	60
17.3 Classifying Stationary Points - The Second-Derivative test	61
18 Constrained Optimization	63
18.1 Motivation	63
18.2 Graphical Reasoning	63
18.3 Algebraic Solution	64
18.3.1 Algorithm	65
18.3.2 Tips for Solving the System of Equations	65
18.4 Three-variable Functions	65
18.5 Interpretation of λ as a rate of change	66
19 Global Optimization	67
19.1 Definitions	67
19.2 How do we know whether a function has a global maximum or minimum?	67
19.3 Algorithm for Global Optimization	68
19.4 Practice Problems	68
IV Integration of Multivariable Functions	69
20 Definite Integration in two variables	71
20.1 Introduction	71
20.2 Construction	71
21 Iterated Integrals	74
21.1 Type I and Type II regions	74
21.2 Switching the Order of Integration	76
22 Double Integral in Polar Coordinates	77
22.1 When to Use it	77

22.2 What is dA in Polar Coordinates?	77
23 Change of Variables in Multiple Integrals	79
23.1 One-to-one Transformation	79
23.2 The Jacobian	80
23.2.1 Harder Problems	81
24 Triple Integrals	82
24.1 Volume and Mass as an Iterated Integral	82
24.2 Changing the Order of Integration - distinct viewpoints	84
24.3 Practice Problems with Polar Coordinates	87
V Vector Calculus	89
25 Vector Fields and Flow Lines	91
25.1 Motivation	91
25.2 Examples of Vector Fields	91
25.3 Flow Lines	93
26 Differential Operators	95
26.1 Curl	95
26.1.1 Understanding Curl in 2D	96
26.2 Divergence	98
26.2.1 Understanding Divergence	98
27 Line Integrals on Parameterized Curves	99
27.1 Motivation	99
27.2 Definition	99
27.2.1 Circulation	100
27.3 Evaluating a line integral	100
28 The Fundamental theorem of Line Integrals	102
28.1 Gradient Vector Field	102
28.1.1 Finding the potential function	103
28.2 The Fundamental theorem of Line Integrals	104
28.2.1 Formulation	104
28.3 Path Independence	105
28.4 Circulation Freedom	108
28.5 Conclusion	109
29 Parametric Surface	110
30 Irrotational Vector Fields	111
30.1 Green's Idea: Volume Integrate the Curl	111
30.1.1 Careful Formulation of Green's Theorem	111
30.2 The Venn Diagram	112
31 Green's Theorem - Applications and Generalizations	114
31.1 Applications of Green's Theorem	114
31.1.1 Evaluating Difficult Line Integrals	114
31.1.2 Calculating Area	115

31.2 Extended Versions of Green's Theorem	116
31.2.1 Finite union of simple regions	116
31.2.2 Finite intersection of simple regions	118
31.2.3 Not simply-connected regions	118
31.3 More Practice Problems	119
Projects	121
A Distances	122
A.1 Distance between two points	122
A.2 Distance from a point to a plane	122
A.3 Distance from a point to a line	122
A.4 Distance between two straight lines	123
A.5 Distance between two planes	123
B Conic Sections and Quadric Surfaces	124
B.1 Conic Sections	124
B.2 Quadric Surface	125
C Epicycloids	128

Part I


Vectors and Three Dimensional Geometry

Chapter 1 | Three Dimensional Coordinate Geometry


To locate a point in a plane, we need two numbers. Any point in the plane can be represented as an ordered pair (a, b) of real numbers, where a is the x -coordinate and b is the y -coordinate. For this reason, a plane is called two-dimensional. To locate a point in space, three numbers are required. We represent any point in space by an ordered triple (a, b, c) of real numbers. For this reason, we will refer to the space as the 3-space or \mathbb{R}^3 .

§1.1 Coordinate Axes and Points in 3-space

The three coordinate axes in 3-space are drawn using a right-hand-thumb rule as follows. It is important that you understand how to draw the axes in different orientations.




A point in three dimension has three coordinates (x, y, z) that denote respectively how far in/out, left/right, up/down a point is from the origin.



Example 1.1.1

Choose one of the corners of this classroom as the origin. Where are the points $P(-4, 3, -5)$ and $Q(3, -2, -6)$?




■ Question 1.

You awaken one morning to find that you have been transferred onto a grid which is set up like a standard right-hand coordinate system. You are at the point $(-1, 3, -3)$, standing upright, and facing the xz -plane. You walk 2 units forward, turn left, and walk for another 2 units. What is your final position? □

§1.2 Distance between two points

The distance between two points (a, b, c) and (x, y, z) in space is given by

$$\sqrt{(x-a)^2 + (y-b)^2 + (z-c)^2}.$$



■ Question 2.

What is the distance from $(1, 2, 1)$ to $(-3, 1, 2)$? □

■ Question 3.

Which of the following points lies closest to the xy -plane?

(i) $(3, 0, 3)$

(ii) $(0, 4, 2)$

(iii) $(2, 4, 1)$

(iv) $(2, 3, 4)$

1.2.1 Equation of a Sphere

We can define a sphere to be collection (locus) of points that are equidistant to a fixed point called the **center**. Suppose the center is at (a, b, c) and the radius is r . Then the distance formula tells us that the required points (x, y, z) satisfy

$$\sqrt{(x-a)^2 + (y-b)^2 + (z-c)^2} = r \iff (x-a)^2 + (y-b)^2 + (z-c)^2 = r^2$$

§1.3 Sets of points described using coordinates**■ Question 4.**

Assume the origin is at the bottom, left, front corner of the classroom. Describe the set of points (x, y, z) with

- $x > 0, y > 0$ and $z \leq 0$
- $z = 3$
- $x = 0, y = 0$

■ Question 5.

Describe/draw the set of points with $0 \leq x \leq 1, 0 \leq y \leq 1-x$, and $z \leq 0$.

■ Question 6.

Find any points where the sphere $(x-1)^2 + (y+3)^2 + (z-2)^2 = 4$ intersects the y -axis.

■ Question 7.

Consider the set of points P given by the equation $z = 5$. Also, let Q be the set of points that satisfy $x^2 + y^2 = 1$.

- Sketch P, Q, and their intersection.
- Find an equation for the intersection.

■ Question 8.

How would you describe the points making up the **solid** cube with sides of length 2 and centered at the origin?

■ Question 9.

An equilateral triangle is standing vertically in 3-space with a vertex above the xy -plane and its two other vertices at $(7, 0, 0)$ and $(9, 0, 0)$. What are the coordinates of the third vertex?

Chapter 2 | Vectors in 3D



§2.1 Definition

Vectors in two or three dimensions are quantities that have both **magnitude** and **direction**. Quantities that are not vectors are called **scalars**.

■ Question 10.

□

Which of the following are vectors?

- (a) The cost of a ticket to the zoo.
- (d) The weight of a giraffe.
- (b) The number of crocodiles in a pond.
- (e) The velocity of a cheetah.
- (c) The volume of an elephant.
- (f) The speed of a gazelle.

§2.2 Graphical representation of vectors

The vector $\vec{v} = \overrightarrow{PQ}$ can be drawn as an arrow with "tail" at the point P and "tip" at the point Q. It represents a vector whose magnitude is equal to the length of PQ and its direction is from Q towards P. This vector is defined entirely by its direction and length, and can be moved around the space.


§2.3 Vector Arithmetic

2.3.1 Addition

To add two vectors \vec{u} and \vec{v} , draw them one after the other, tip to tail. The vector from the base of the pair to the final tip is the sum of the vectors.

2.3.2 Scalar Multiplication

If λ is a scalar i.e. a real number, then $\lambda\vec{v}$ is a vector whose magnitude is λ times the magnitude of \vec{v} . It has the same direction as \vec{v} if $\lambda > 0$ and the opposite direction if $\lambda < 0$.



Warning: If $\lambda = 0$, we get the **Zero Vector**, a vector whose magnitude is zero and is **omnidirectional**!

Definition 2.3.2

Two vectors are called *parallel* if they point in the same direction. That means \vec{u} and \vec{v} are parallel if there exists a *positive* constant λ such that $\vec{u} = \lambda \vec{v}$. The vectors are called *antiparallel* if λ is negative.

2.3.3 Subtraction

$\vec{u} - \vec{v}$ is defined as $\vec{u} + (-1) \times \vec{v}$.

■ Question 11.

Suppose the three sides of a triangle $\triangle ABC$ are denoted by vectors as $\vec{c} = \overrightarrow{AB}$, $\vec{a} = \overrightarrow{BC}$, and $\vec{b} = \overrightarrow{CA}$. What is $\vec{a} + \vec{b} + \vec{c}$? □

■ Question 12.

Consider $\triangle ABC$ as above. Let D be the mid-point of BC and let $\vec{m} = \overrightarrow{AD}$ be one of the medians of the triangle. Find \vec{m} in terms of \vec{a} , \vec{b} and \vec{c} . □

Exploration Activity

It is in fact also possible to find the angle bisector vector in terms of the sides. However that requires the law of sines in a triangle. In case you are interested, here is the precise problem. Suppose D is a point on BC such that $\angle BAD = \angle CAD$. Find \overrightarrow{AD} in terms of \vec{a} , \vec{b} and \vec{c} .

§2.4 Symbolic representation of vectors

A vector of magnitude 1 is called a **unit vector**. The unit vectors along X-, Y- and Z-axes are called \hat{i} , \hat{j} and \hat{k} respectively.

Consider the vector $\vec{v} = \overrightarrow{PQ}$ where $P = (x_1, y_1, z_1)$ and $Q = (x_2, y_2, z_2)$. Then it can be also written as

$$\vec{v} = \underbrace{(x_2 - x_1)}_{\Delta x} \hat{i} + \underbrace{(y_2 - y_1)}_{\Delta y} \hat{j} + \underbrace{(z_2 - z_1)}_{\Delta z} \hat{k} = \langle \Delta x, \Delta y, \Delta z \rangle$$

The projection of a vector on to the axes are called its **components**. Thus the X-component of the vector \vec{v} above is $(\Delta x)\hat{i}$ etc. Clearly a vector is the sum of its components.

Addition or scalar multiplication of a vector can be done component-wise.

$$\langle a_1, a_2, a_3 \rangle + \langle b_1, b_2, b_3 \rangle = \langle a_1 + b_1, a_2 + b_2, a_3 + b_3 \rangle \quad c \langle a_1, a_2, a_3 \rangle = \langle ca_1, ca_2, ca_3 \rangle$$

2.4.1 Position vector - components vs coordinates

If we write $\vec{v} = \langle a_1, a_2, a_3 \rangle$ or equivalently $\vec{v} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$, we mean a vector with tail at $(0, 0, 0)$ and tip at $P(a_1, a_2, a_3)$. In this way every point (a_1, a_2, a_3) corresponds to the unique vector $\langle a_1, a_2, a_3 \rangle$ called its **position vector** (see fig. 2.1). The length of the components of the position vector are equal to the coordinates of the point.




Figure 2.1

§2.5 Magnitude

The magnitude of a vector $\vec{v} = \langle a, b, c \rangle$ is denoted by $\|\vec{v}\|$, pronounced “norm” or \vec{v} , and is equal to $\sqrt{a^2 + b^2 + c^2}$. This is an easy consequence of Pythagoras theorem!

■ Question 13.

How can you use $\|\vec{v}\|$ to create a unit vector parallel to \vec{v} ? □

§2.6 Calculating components in 2D

Suppose a vector \vec{v} makes an angle θ with the positive X-axis. Then we can express the magnitude $\|\vec{v}\|$ to express a vector \vec{v} in terms of trigonometric functions as follows:

$$\vec{v} = (\Delta x)\hat{i} + (\Delta y)\hat{j} = (\|\vec{v}\| \cos \theta)\hat{i} + (\|\vec{v}\| \sin \theta)\hat{j} = \|\vec{v}\|(\cos \theta\hat{i} + \sin \theta\hat{j})$$

■ Question 14.

Suppose a three dimensional vector \vec{v} makes angles α, β and γ with positive X-, Y- and Z-axis respectively. Then show that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$$

§2.7 Word problems

■ Question 15.

Bert and Ernie are trying to drag a large box on the ground. Bert pulls the box toward the north with a force of 30 N, while Ernie pulls the box toward the east with a force of 40 N. What is the resultant force on the box?

■ Question 16.

An airplane at altitude is flying NE with airspeed 700 km/hr, with wind from the West at 60 km/hr. Use vectors to determine the resulting direction and ground speed of the plane.

Chapter 3 | Dot Product of Vectors



§3.1 Definition

3.1.1 Algebraic

The **dot product** or scalar product of two vectors $\vec{u} = \langle u_1, u_2, u_3 \rangle$ and $\vec{v} = \langle v_1, v_2, v_3 \rangle$ is defined to be the sum of the product of the components.

$$\vec{u} \cdot \vec{v} = \langle u_1, u_2, u_3 \rangle \cdot \langle v_1, v_2, v_3 \rangle := u_1 v_1 + u_2 v_2 + u_3 v_3$$

■ Question 17.



What is $\langle 1, 2, 3 \rangle \cdot \langle 4, -5, 6 \rangle$? How about $\langle 1, 2 \rangle \cdot \langle 3, 4, 5 \rangle$?

3.1.2 Geometric

If the angle between \vec{u} and \vec{v} is θ then

$$\vec{u} \cdot \vec{v} := \|\vec{u}\| \|\vec{v}\| \cos \theta$$

■ Question 18.



Does it matter whether the angle θ is calculated from \vec{u} to \vec{v} or in the other order?

■ Question 19.



What is $\vec{u} \cdot \vec{v}$ if

- (a) $\vec{u} \perp \vec{v}$?
- (b) $\vec{u} = \vec{v}$?
- (c) $\vec{u} \parallel \vec{v}$?

Theorem 3.1.3

Two vectors \vec{u} and \vec{v} are orthogonal (i.e. perpendicular) if and only if $\vec{u} \cdot \vec{v} = 0$.

§3.2 Basic properties

From the definitions the following basic properties of the dot product are easy to prove. If \vec{u}, \vec{v} and \vec{w} are vectors of the same dimension and c is a scalar, then

- (a) $\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$
- (b) $\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$
- (c) $(c\vec{v}) \cdot \vec{w} = c(\vec{v} \cdot \vec{w}) = \vec{v} \cdot (c\vec{w})$

§3.3 Angle between two vectors

We can use both definitions of dot product together to calculate angle between two given vectors.

■ Question 20.

Find the angle between $\langle 1, 2, 1 \rangle$ and $\langle 1, -1, 1 \rangle$.

□

■ Question 21.

In a molecule of Methane (CH_4) you have four hydrogen atoms bonded to a carbon atom. The four hydrogen atoms form the corner of a regular tetrahedron and the carbon atom lies in the center. What is the angle between any two of the C – H bonds?

Hint: Think of the four hydrogen atoms lying on four corners of a cube.

§3.4 Vector projections

Take a vector \vec{u} and resolve it into two components, one along another given vector \vec{v} and another perpendicular to \vec{v} . We call these two components \vec{u}_{\parallel} and \vec{u}_{\perp} respectively. Thus

$$\vec{u} = \vec{u}_{\parallel} + \vec{u}_{\perp}$$

The component \vec{u}_{\parallel} is called the **projection** of \vec{u} on to \vec{v} , and is denoted $\text{Proj}_{\vec{v}} \vec{u}$.

Geometrically speaking, if we take a screen along \vec{v} and shine a light perpendicular to it from above, the shadow cast by \vec{u} would be $\text{Proj}_{\vec{v}} \vec{u}$.

Theorem 3.4.4: Projection Formula

$$\text{Proj}_{\vec{v}} \vec{u} = \vec{u}_{\parallel} = \left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

■ Question 22.

Prove above theorem.

HINT: Use the dot product formula for $\cos \theta$.

□

■ Question 23.

The vertices of a triangle $\triangle ABC$ are $A = (4, 3, 2)$, $B = (1, 3, 1)$, and $C = (-5, 5, -2)$. Let D be the foot of the perpendicular from A to the side \overline{BC} . Find the vector \overrightarrow{AD} .

HINT: Find \overrightarrow{BD} first.

□

Chapter 4 | Cross Product of Vectors



§4.1 Definition

4.1.1 Geometric

The cross product of two vectors \vec{u} and \vec{v} is the **vector** $\vec{u} \times \vec{v}$ whose

- magnitude is equal to $\|\vec{u}\| \|\vec{v}\| \sin \theta$, where θ is the angle from \vec{u} to \vec{v} and
- direction is perpendicular to the both \vec{u} and \vec{v} , determined by the right-hand-thumb rule.

■ Question 24.

Does it matter whether the angle θ is calculated from \vec{u} to \vec{v} or in the other order?

■ Question 25.

What is $\hat{i} \times \hat{j}$? What is $\hat{i} \times \hat{i}$? What is $\hat{j} \times \hat{i}$?

4.1.2 Algebraic

If $\vec{u} = \langle u_1, u_2, u_3 \rangle$ and $\vec{v} = \langle v_1, v_2, v_3 \rangle$, then we define $\vec{u} \times \vec{v}$ to be

$$\langle u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1 \rangle.$$

There is a handy way of remembering this definition: the cross product $\vec{u} \times \vec{v}$ is equal to the determinant

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ v_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} = \begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix} \hat{i} - \begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix} \hat{j} + \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix} \hat{k}$$



Warning: The cross product is only defined for three-dimensional vectors.

§4.2 Basic Properties

From the definitions the following basic properties of the cross product are easy to prove. If \vec{u} , \vec{v} and \vec{w} are vectors of the same dimension and c is a scalar, then

- $\vec{v} \times \vec{w} = -\vec{w} \times \vec{v}$
- $\vec{u} \times (\vec{v} + \vec{w}) = \vec{u} \times \vec{v} + \vec{u} \times \vec{w}$
- $(c\vec{v}) \times \vec{w} = c(\vec{v} \times \vec{w}) = \vec{v} \times (c\vec{w})$

Theorem 4.2.5

Two vectors \vec{u} and \vec{v} are parallel or antiparallel if and only if $\vec{u} \times \vec{v} = 0$.

§4.3 Cross product as area


From the geometric definition, it is easy to see that $\|\vec{u} \times \vec{v}\|$ is equal to the area of the parallelogram whose two adjacent sides are given by \vec{u} and \vec{v} .

■ Question 26.

Show this. Use the fact that the area of the parallelogram is twice the area of the triangle whose two sides are given by \vec{u} and \vec{v} . Then use the area formula for a triangle. □

4.3.1 Volume of a parallelepiped - Coplanarity

Extending this geometric idea, we can show that the volume of a parallelepiped whose three adjacent sides are given by \vec{a} , \vec{b} , and \vec{c} is equal to $|\vec{a} \cdot (\vec{b} \times \vec{c})|$.



Consequently, we have the following theorem

Theorem 4.3.6

Three vectors \vec{a} , \vec{b} , and \vec{c} are coplanar iff $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$.

Exploration Activity

$\vec{a} \cdot (\vec{b} \times \vec{c})$ is called the **scalar triple product** or **box product** of the three vectors, as also denoted by $[\vec{a}, \vec{b}, \vec{c}]$. An interesting observation here is that since the box product relates to the volume of the parallelepiped, we get

$$[\vec{a}, \vec{b}, \vec{c}] = [\vec{b}, \vec{c}, \vec{a}] = [\vec{c}, \vec{a}, \vec{b}]$$

So if our goal is to only calculate the volume, it doesn't matter which vectors we choose as \vec{a} , \vec{b} and \vec{c} as long they are the three adjacent sides.

■ Question 27.

Suppose λ and μ are real numbers such that

- the three vectors

$$\vec{u} = 2\hat{i} + 3\hat{j} + \hat{k}, \quad \vec{v} = \hat{i} + \lambda\hat{j} + \mu\hat{k}, \quad \vec{w} = 7\hat{i} + 3\hat{j} + 2\hat{k}$$

are coplanar, and

- The vector \vec{v} has magnitude $\sqrt{2}$.

Find all possible values of λ and μ .

3

Chapter 5 | Lines and Planes



§5.1 Lines in space

The equation of a line through a point (x_0, y_0, z_0) and parallel to the vector $\vec{u} = \langle a, b, c \rangle$ can be expressed in many ways:

- as parametric scalar equations:

$$x = x_0 + ta, \quad y = y_0 + tb, \quad z = z_0 + tc;$$

- as a parametric vector equation:

$$\vec{r}(t) = \vec{r}_0 + t\vec{u}, \quad \text{where } \vec{r} = \langle x, y, z \rangle \text{ and } \vec{r}_0 = \langle x_0, y_0, z_0 \rangle$$

- or by symmetric equations:

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$



Warning: Observe that the symmetric equation form doesn't make sense in case one of a, b or c is zero.

■ Question 28.

Let \mathcal{L} be the line which passes through the points $(1, -2, 3)$ and $(4, -4, 6)$. Find its equation in all three forms. □

§5.2 Planes in space

The equation of a plane through the point (x_0, y_0, z_0) and perpendicular (or normal or orthogonal) to the vector $\vec{n} = \langle a, b, c \rangle$ also has many (equivalent) equations:

- as a vector equation:

$$\vec{n} \cdot (\vec{r} - \vec{r}_0) = 0 \quad \text{or} \quad \vec{n} \cdot \vec{r} = \vec{n} \cdot \vec{r}_0$$

where again $\vec{r} = \langle x, y, z \rangle$ and $\vec{r}_0 = \langle x_0, y_0, z_0 \rangle$; or equivalently

- as a scalar equation:

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0 \quad \text{or} \quad ax + by + cz = d$$

where $d = ax_0 + by_0 + cz_0$ is a constant.

Note: Two planes $Ax + By + Cz = D$ and $ax + by + cz = d$ are parallel iff there exists a constant λ such that $\langle A, B, C \rangle = \lambda \langle a, b, c \rangle$. The planes are identical if they are parallel and $D = \lambda d$.

■ Question 29.

Find an equation describing the plane which goes through the point $(1, 3, 5)$ and is perpendicular to the vector $\langle 2, 1, -3 \rangle$. □

■ Question 30.

□

Find an equation describing the plane which passes through the points P(2, 2, 1), Q(3, 1, 0), and R(0, -2, 1).

§5.3 Types of problems

One of the fundamental topic in Multivariable Calculus is to learn how to find equations of straight lines and planes in three dimensions using ideas from vectors (dot and cross product). The following lists an incomplete but fairly diverse types of problems that you should be able to solve using these ideas.

5.3.1 Finding a Plane

You should know how to find equation of a plane from the following data.

- (a) Plane through a given point and perpendicular to a given vector.
- (b) Plane through a given point and parallel to a given plane.
- (c) Plane containing a given line and parallel to a given plane.
- (d) Plane passing through three given points.
- (e) Plane with specified x -, y - and z -intercepts.
- (f) Plane through a given point and containing a given straight line.
- (g) Plane through a given point and containing the line of intersection of two other given planes.
- (h) Plane through a given point and perpendicular to two other given planes.
- (i) Plane passing through two points and perpendicular to a given plane.
- (j) Plane containing the line of intersection of two other given planes and perpendicular to a given plane.

5.3.2 Finding a Line

You should know how to find equation of a straight line from the following data.

- (a) Line through two given points.
- (b) Line through one given point and in the direction of a given vector.
- (c) Line through one given point and parallel to a given straight line.
- (d) Line of intersection of two given planes.
- (e) Line through one given point and perpendicular to a given plane.
- (f) Line through a given point, that is perpendicular to a given straight line and intersects this second line.
- (g) Line through a given point, that is parallel to (i.e. lies in) a given plane and perpendicular to a given straight line.

§5.4 Practice problems**■ Question 31.**

□

Below is a list of vectors and a list of properties. Match the two sets in such a way that each entry in left column matches a different entry in right column.

A. $\langle 3, -2, 8 \rangle$	I. is parallel to the straight line $\frac{x-1}{2} = y - 3 = z$
B. $\langle 4, 2, 2 \rangle$	II. is perpendicular to the plane $z - 2y - x = 3$
C. $\langle 3, 1, -1 \rangle$	III. is perpendicular to both $\langle 2, 3, 0 \rangle$ and $\langle -2, 5, 2 \rangle$
D. $\langle 1, 2, -1 \rangle$	IV. is parallel to the plane $x - y + 2z = 3$

■ Question 32.



- (a) Find parametric equations for the line through the points $(6, 1, 1)$ and $(9, 1, 4)$. Call this line L_1 .
- (b) Find parametric equations for the line through the points $(-4, 4, 0)$ and $(-6, 5, 1)$. Call this line L_2 .
- (c) Find parametric equations for the line through the points $(6, -1, -5)$ and $(2, 1, -3)$. Call this line L_3 .
- (d) Verify that L_2 and L_3 are parallel. (Their direction vectors should be parallel.) Are they the same line? How could you tell?
- (e) Do lines L_1 and L_2 intersect? If so, where?
- (f) Find the intersection of L_1 with the plane given by the equation $2x + y + 3z = 7$.
- (g) (3 points) Find the point on the plane $2x + y + 3z = 7$ which is closest to the origin.
- (h) (3 points) Find the point on L_2 closest to the origin.

■ Question 33.



Find a vector parallel to the intersection of the two planes $2x - 3y + 5z = 2$ and $4x + y - 3z = 7$.

Find the equation of the line of intersection.

■ Question 34.



Find the distance of the point $P = (1, 0, 1)$ from the plane $x + y - z = 1$.

■ Question 35.



Let L_1 be the line with parametric vector equation $\vec{r}_1(t) = \langle 7, 1, 3 \rangle + t\langle 1, 0, -1 \rangle$ and L_2 be the line described parametrically by $x = 5, y = 1 + 3t, z = t$. How many planes are there that contain L_2 and are parallel to L_1 ? Find an equation describing one such plane.

■ Question 36.



Find an equation for the plane that contains the line in the XY-pane where $y = 1$, and the line in the XZ-pane where $z = 2$.


Chapter 6 | Polar Coordinate System and Polar Curves



§6.1 Definition

A coordinate system represents a point in the plane by an ordered pair of numbers called coordinates. Usually we use Cartesian coordinates, which are directed distances from two perpendicular axes. In this chapter, we look at a coordinate system introduced by Newton, called the **polar coordinate system**, which is more convenient for many purposes.

We choose a point in the plane that is called the pole (or origin) and is labeled O. Then we draw a ray (half-line) starting at O called the polar axis. This axis is usually drawn horizontally to the right and corresponds to the positive x -axis in Cartesian coordinates.



If P is any other point in the plane, let r be the distance from O to P and let θ be the angle (usually measured in radians) between the polar axis and the line OP as in the figure above. Then the point P is represented by the ordered pair (r, θ) and r, θ are called polar coordinates of P. We use the convention that an angle is positive if measured in the counterclockwise direction from the polar axis and negative in the clockwise direction. If $P = O$, then $r = 0$ and we agree that $(0, \theta)$ represents the pole for any value of θ .

6.1.1 When r is negative

The convention is to regard r as a ‘signed’ radius. That means if for some value of θ , we are provided a negative value of r , we go across origin and take the diametrically opposite point. In the second figure above, the points $(-r, \theta)$ and (r, θ) lie on the same line through O and at the same distance $|r|$ from O, but on opposite sides of O.

Notice that $(-r, \theta)$ represents the same point as $(r, \theta + \pi)$.

■ Question 37.

Plot the points $(2, -2\pi/3)$ and $(-3, 3\pi/4)$. Find the Cartesian coordinates of the points above.

■ Question 38.

In polar coordinates, what shapes are described by $r = k$ and $\theta = k$, where k is a constant?

(a) Draw $r = 0, r = \frac{2\pi}{3}, r = \frac{4\pi}{3}, r = 2\pi, \theta = 0, \theta = \frac{2\pi}{3}$, and $\theta = \frac{4\pi}{3}$.

(b) Find the equations of above curves in Cartesian coordinates.

§6.2 Converting between Polar and Cartesian Coordinates

Using simple trigonometry, we can find that the polar point (r, θ) corresponds to the Cartesian point $(r \cos \theta, r \sin \theta)$. Equivalently, starting from a Cartesian coordinate (x, y) we can make the change of variables

$$r = \sqrt{x^2 + y^2}, \quad \theta = \arctan \frac{y}{x}$$

to get (one possible) polar coordinate (r, θ) .

■ Question 39.

Express the point with polar coordinates $P(3, \frac{3\pi}{4})$ in Cartesian coordinates. Express the point with Cartesian coordinates $Q(3, 3)$ in Polar coordinates.

■ Question 40.

Find the polar equation of the straight line $x = 5$.

Note: Often you might find it easier to manipulate a given Cartesian equation **using trigonometry** to isolate $r \cos \theta$ and replace it with x and similarly replace $r \sin \theta$ with y .

■ Question 41.

Convert the following polar equations into Cartesian:

- | | |
|-------------------------|-----------------------------|
| (a) $r^2 = 5$. | (c) $r^2 \sin(2\theta) = 1$ |
| (b) $r \cos \theta = 5$ | (d) $r = \cos \theta$ |

§6.3 Polar Curves

The graph of a polar equation $r = f(\theta)$, or more generally $F(r, \theta) = 0$, consists of all points P that have at least one polar representation (r, θ) whose coordinates satisfy the equation.

So how do we plot a curve whose equation is of the form $r = f(\theta)$?

One possible method is to change it into Cartesian coordinates and plot that instead.

■ Question 42.

Sketch the curve with polar equation $r = 2 \cos \theta$.

This method is not always easy if the equation is complicated. In such cases, the following procedure is advised.

- **Step 1.** Instead of plotting points on the XY-plane, we first sketch the graph of $r = f(\theta)$ in Cartesian coordinates where the horizontal axis is θ and the vertical axis is r .
- **Step 2.** Use the Cartesian graph in Step 1 as a guide to sketch the values of r that correspond to increasing values of θ . Remember that θ can only go up to 2π .



We will demonstrate using an example.

Example 6.3.7

Suppose we wish to sketch the curve $r = \cos(2\theta)$.

We first sketch $r = \cos 2\theta$ for $0 \leq \theta \leq 2\pi$, in Cartesian coordinates. As θ increases from 0 to $\pi/4$, the value of r decreases from 1 to 0 and so we draw the corresponding portion of the polar curve (indicated by ①).

As θ increases from $\pi/4$ to $\pi/2$, the value of r decreases from 0 to -1 , this means that the distance from O increases from 0 to 1, but instead of being in the first quadrant this portion of the polar curve (indicated by ②) lies on the opposite side of the pole in the third quadrant. The remainder of the curve is drawn in a similar fashion, with the arrows and numbers indicating the order in which the portions are traced out. The resulting curve has four loops and is called a **four-leaved rose**.






■ Question 43.

Which part of the four-leaved rose correspond to $0 \leq \theta \leq \pi/2$? □

6.3.1 Symmetry

When we sketch polar curves it is sometimes helpful to take advantage of symmetry.

- If a polar equation is unchanged when θ is replaced by $-\theta$, the curve is symmetric about the polar axis.
- If the equation is unchanged when r is replaced by $-r$, or when θ is replaced by $\theta + \pi$, the curve is symmetric about the pole. (This means that the curve remains unchanged if we rotate it through 180° about the origin.)
- If the equation is unchanged when θ is replaced by $\pi - \theta$, the curve is symmetric about the vertical line $\theta = \pi/2$.



Chapter 7 | Vector Valued Functions and Space Curves



Any curve in 3-space can be described as either a collection of points

$$(x(t), y(t), z(t))$$

for t in some interval (possibly infinite) or as the trace of the heads of the position vectors (which start at the origin)

$$\vec{r}(t) = \langle x(t), y(t), z(t) \rangle = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$$



Definition 7.0.8

$\vec{r}(t)$ is an example of a **vector valued function** of a real variable t . In general a vector valued function $\vec{F}(t)$ can be written as

$$\vec{F}(t) = \langle f(t), g(t), h(t) \rangle$$

We will come back to general vector-valued functions in the next chapter.

Example 7.0.9: Circle

A circle in the xy -plane can be described as the set of pairs $(x(t), y(t))$ generated by the **parameterization**

$$x(t) = \cos(t) \text{ and } y(t) = \sin(t) \quad \text{for } 0 \leq t \leq 2\pi$$

associated with the position vectors

$$\vec{r}(t) = \cos(t)\hat{i} + \sin(t)\hat{j} \quad \text{for } 0 \leq t \leq 2\pi$$

where the parameter t is the angle from the x -axis.

When curves are generated in this manner, they are called **parametrized curves**. Notice that **parameterization** carries more information than simply the final curve shape in the xy -plane. In particular, the example above indicates that the curve shape (circle) in the xy -plane is traced exactly once around, and that this curve shape is traced counterclockwise as t increases through its range.

§7.1 Different ways of parametrizing the same curve

We can parametrize the same curve in different ways and interpret each parametrization as the motion of a particle with the parameter t being time.

■ Question 44.

Explain why all of the parametrized curves below should look like a circle. In each of the parametrization, determine how many times around, and in which direction (clockwise or counterclockwise) the curve is traced.

- $x(t) = \cos(2t)$ and $y(t) = \sin(2t)$ for $0 \leq t \leq 2\pi$
- $x(t) = \cos(t^2)$ and $y(t) = \sin(t^2)$ for $0 \leq t \leq 2\pi$
- $x(t) = \cos(t)$ and $y(t) = -\sin(t)$ for $0 \leq t \leq 2\pi$
- $x(t) = -\cos(t)$ and $y(t) = \sin(t)$ for $0 \leq t \leq 2\pi$
- $x(t) = -\cos(t)$ and $y(t) = -\sin(t)$ for $0 \leq t \leq 2\pi$

■ Question 45.

Recall that $\vec{r}(t) = \vec{r}_0 + t\vec{v}$ (for $t \in \mathbb{R}$) gives the parametrization of a straight line passing through \vec{r}_0 and parallel to \vec{v} . What kind of curve are the following:

- $\vec{r}(t) = \vec{r}_0 + t^2\vec{v}$ for $t \in \mathbb{R}$
- $\vec{r}(t) = \vec{r}_0 + t\vec{v}$ for $a \leq t \leq b$

■ Question 46.

Create a graph of each of the following vector-valued functions:

- The plane curve represented by $\vec{r}(t) = (5 \cos t)\hat{i} + (2 \sin t)\hat{j}$.
- The plane curve represented by $\vec{r}(t) = \langle e^t - 1, e^{2t} \rangle$.

§7.2 3D Curves

■ Question 47.


The vector function $\vec{r}(t) = \langle 5 + 2 \cos t, 1 + 2 \sin t, 3 \rangle$ (for $t \in \mathbb{R}$) traces a circle. Find the center, radius, and plane containing the circle.

7.2.1 Helix

The parametric equation of a helix is given by

$$\vec{r}(t) = \langle \cos t, \sin t, t \rangle$$

When looked on from above (from the positive z direction), this curve is simply a circle in the xy -plane. The $z = t$ component lifts the circle into the helix spinning above the circle in the plane. If we visualize this as a particle at the tip of the position vector $\vec{r}(t)$, then from above it looks like the particle is simply spinning in a circle. But we also know that $z = t$, so the particle is rising at a constant rate. Hence we get the picture below



■ **Question 48.**

□

Match each vector-valued function to the curve it parametrizes.

(a) $\vec{r}(t) = \langle t \cos t, t \sin t, t \rangle$







(d) $\vec{r}(u) = \langle \cos u, \sin u, 1 + \sin 4u \rangle$

(b) $\vec{r}(t) = \langle \cos t, \sin t, t^3 \rangle$

(e) $\vec{r}(u) = \langle \cos u, \sin u, 1 + 4 \sin u \rangle$

(c) $\vec{r}(t) = \langle \cos(t^3), \sin(t^3), t^3 \rangle$

(f) $\vec{r}(t) = \langle 2 \cos t, 1 + 4 \cos t, 3 \cos t \rangle$



§7.3 Parametrization from equation of curve

Example 7.3.10

A parametrization of the parabola $x = 1 - y^2$ in xy - plane can be given by


$$y(t) = t, \quad x(t) = 1 - t^2, \quad t \in \mathbb{R}$$

■ Question 49.

The surfaces $z = \sin(x - y)$ and $y = 2x$ intersect in a curve. Find a parameterization of the curve. □

■ Question 50.

The cylinder $x^2 + y^2 = 1$ and the plane $y + z = 2$ intersect in a curve. Find a parameterization of the curve. □



Chapter 8 | Calculus of Vector-valued Functions



§8.1 Derivative of a Vector-valued function

The derivative \vec{r}' of a vector function \vec{r} is defined in much the same way as for real-valued functions. The idea is to consider the difference vector $\vec{r}(t + \Delta t) - \vec{r}(t)$ between the position vector $\vec{r}(t + \Delta t)$, a little time Δt beyond t , and the position vector $\vec{r}(t)$ at t . By taking $\Delta t \rightarrow 0$, the instantaneous **velocity vector** at time t is given by

$$\begin{aligned}\vec{v}(t) = \vec{r}'(t) &= \lim_{\Delta t \rightarrow 0} \frac{\vec{r}(t + \Delta t) - \vec{r}(t)}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \left(\frac{x(t + \Delta t) - x(t)}{\Delta t} \right) \hat{i} + \lim_{\Delta t \rightarrow 0} \left(\frac{y(t + \Delta t) - y(t)}{\Delta t} \right) \hat{j} + \lim_{\Delta t \rightarrow 0} \left(\frac{z(t + \Delta t) - z(t)}{\Delta t} \right) \hat{k} \\ &= x'(t) \hat{i} + y'(t) \hat{j} + z'(t) \hat{k}\end{aligned}$$

Definition 8.1.11

A parametrization $\vec{r}(t)$ is called **regular** if it is differentiable and $\vec{r}'(t) \neq \vec{0}$ for all t in the domain of \vec{r} .

Geometrically, we observe that \vec{r}' is a vector that lies on the tangent line to the curve $\vec{r}(t)$. For this reason, the vector $\vec{r}'(t)$ is called the **tangent vector** to the curve defined by \vec{r} , provided that it is regular. So intuitively we can think of regular curves as curves which always have a tangent vector (i.e. no corners or cusps; and when the tangent vector turns, it turns smoothly).

Definition 8.1.12

Let C be a regular curve defined by the vector function $\vec{r}(t)$. The **principal unit tangent vector** to the curve $\vec{T}(t)$ at t is defined to be

$$\vec{T}(t) = \frac{\vec{r}'(t)}{\|\vec{r}'(t)\|}.$$

8.1.1 Differentiation Rules

Consider three vector-valued functions $\vec{r}(t)$, $\vec{r}_1(t)$, and $\vec{r}_2(t)$. In the last section, we defined $\vec{r}'(t)$. What do you think are the formula for the following? Here $p(t)$ is a one-variable differentiable scalar-valued (real) function of t . .

(a) $\frac{d}{dt}[c \vec{r}(t)]$

(b) $\frac{d}{dt}[p(t) \vec{r}(t)]$ (scalar product rule)

(c) $\frac{d}{dt}[\vec{r}_1(t) \cdot \vec{r}_2(t)]$ (dot product rule)

- (d) $\frac{d}{dt}[\vec{r}_1(t) \times \vec{r}_1(t)]$ (cross product rule)
- (e) $\frac{d}{dt}[\vec{r}(p(t))]$ (chain rule)

The following result will be important later as we discuss motion in 3-space!

■ Question 51.

If $\|\vec{r}(t)\| = c$, then show that $\vec{r}(t)$ is perpendicular to $\vec{r}'(t)$.

□

8.1.2 Interpretation as Velocity

Think of a parametrized curve as the trajectory of a point that is moving on the curve. At time t , its position vector is given by $\vec{r}(t)$. Then its velocity vector is the rate of change of position vector with respect to time, i.e. the tangent vector derived in last section!

■ Question 52.

A particle travels along the line $x = 1 + t, y = 5 + 2t, z = -7 + t$. When and where does the particle hit the plane $x + y + z = 1$?

□

■ Question 53.

Suppose the trajectories of two particles are given by these vector functions:

$$\vec{r}_1(t) = \langle t^2, 7t - 12, t^2 \rangle \quad \text{and} \quad \vec{r}_2(t) = \langle 4t - 3, t^2, 5t - 6 \rangle.$$

What is the difference between asking if the particles collide and asking if their paths intersect?

□

■ Question 54.

Let $f(x, y) = 3x^2 + 4y^2$.

- (a) Find the parametric equation $\vec{r}(t)$ of the level curve of f that passes through the point $(1, 1)$.
- (b) Find the velocity $\vec{r}'(t)$ at $(1, 1)$.
- (c) Verify that $\vec{r}'(t)$ at $(1, 1)$ is perpendicular to $\nabla f(1, 1)$.

□

■ Question 55.

A fly is sitting on the wall at the point $(0, 1, 3)$. At time $t = 0$, he starts flying; his velocity at time t is given by $\vec{v}(t) = \langle \cos 2t, e^t, \sin t \rangle$. Find the fly's location at time t .

□

■ Question 56.

Find the angle of intersection between the curve given by its parametric equation $\vec{r}(t) = \langle t, 2t^2 \rangle$, and the parabola $y = x^2 + 4$.

□

§8.2 Integral of a vector-valued function - arc length


The magnitude $\|\vec{r}'(t)\|$ of the velocity vector is the speed. Hence the value $\|\vec{r}'(t)\|$ approximates the distance traveled from t to $t + \Delta t$ along the curve parameterized by $\vec{r}(t)$. Adding such approximations from $t = a$ to $t = b$ approximates the length of the curve

$$\sum_{t=a}^{t=b} \|\vec{r}'(t)\| \Delta t \approx \text{Length of curve } \vec{r}(t) \text{ from } t = a \text{ to } t = b$$

Taking $\Delta t \rightarrow 0$ gives a perfect approximation in terms of the following integral:

$$s(t) = \int_{t=a}^{t=b} \|\vec{r}'(t)\| dt = \text{Length of curve } \vec{r}(t) \text{ from } t = a \text{ to } t = b$$

$s(t)$ is then the arc length of the curve - physically, the distance travelled along the curve.



■ Question 57.

□

- (a) What is the speed of an object on the circle parameterized by $\vec{r}(t) = \cos(t)\hat{i} + \sin(t)\hat{j}$?
- (b) Compute the length of this $\vec{r}(t)$ from $0 \leq t \leq 2\pi$.
- (c) Use a dot product to find the orientation of the circle's velocity vectors $\vec{r}'(t)$ relative to its position vectors $\vec{r}(t) = \cos(t)\hat{i} + \sin(t)\hat{j}$.
- (d) Use a dot product to find the orientation of the circle's acceleration vectors $\vec{r}''(t)$ relative to its velocity vectors.

8.2.1 Arc Length Parameterization

It is sometimes useful to parametrize a curve with respect to arc length because arc length arises naturally from the shape of the curve and does not depend on a particular coordinate system or a particular parametrization. If a curve $\vec{r}(t)$ is already given in terms of a parameter t and $s(t)$ is the arc length function, then we may be able to solve for t as a function of s i.e. $t = t(s)$. Then the curve can be reparametrized in terms of s by substituting for t , i.e. $\vec{r} = \vec{r}(t(s))$. Now if $s = 3$ for instance, then we know $\vec{r}(t(3))$ is the position vector of the point 3 units of length along the curve from its starting point.

■ Question 58.

□

Find an arc length parametrization for the following curve: $\vec{r}(t) = \langle 2t - 1, t + 3, 5 + 2t \rangle$ for $t \geq 0$.

■ **Question 59.**

□

Reparametrize the helix $\vec{r}(t) = \langle 4\cos t, 4\sin t, 3t \rangle$ with respect to arc length measured from $t = 0$ in the direction of increasing t .

§8.3 Curvature

Next, we would like to define the geometric idea of *curvature* of a curve \mathcal{C} at a point. Here are some of the properties we would like to have:

- (a) Curvature should be some kind of measure of how sharply \mathcal{C} is turning, measured point by point.
- (b) The curvature of a straight line should be everywhere.
- (c) The curvature of a circle should be everywhere, and the larger the radius of the circle, the the curvature.
- (d) The curvature should have something to do with the second derivative.

Definition 8.3.13

The curvature of \mathcal{C} at a given point κ is a measure of how quickly the curve changes direction at that point. Specifically, we define it to be the magnitude of the rate of change of the unit tangent vector with respect to arc length.

$$\kappa = \left\| \frac{d\vec{T}}{ds} \right\|$$

If the curve is not given using arc length parametrization, this definition is not very helpful. So we are going to do some manipulation using chain rule to redefine curvature in terms of t . Recall that the arc length of \mathcal{C} is given by

$$s(t) = \int \|\vec{r}'(t)\| dt$$

So if we differentiate both sides of the arc length formula, then by using the Fundamental Theorem of Calculus, we obtain

$$\frac{ds}{dt} = \|\vec{r}'(t)\|.$$

Hence, by chain rule,

$$\frac{d\vec{T}}{ds} = \frac{d\vec{T}}{dt} \frac{dt}{ds} = \frac{\frac{d\vec{T}}{dt}}{\frac{ds}{dt}} \implies \kappa = \frac{\|\vec{T}'(t)\|}{\|\vec{r}'(t)\|}.$$

We can further manipulate the above equation by using the definition of \vec{T} to get

Theorem 8.3.14

The curvature of the curve given by the vector function $\vec{r}(t)$ is

$$\kappa = \frac{\|\vec{r}'(t) \times \vec{r}''(t)\|}{\|\vec{r}'(t)\|^3}.$$

■ Question 60.

Find the curvature $\kappa(t)$, of the curve given by $\vec{r}(t) = \langle -2t, t^2, \frac{1}{3}t^3 \rangle$ and then find the curvature of the curve at the point $(0, 0, 0)$.

Note: When $\vec{r}(t)$ is a planar curve $\langle f(t), g(t) \rangle$, we can simplify above formula to

$$\kappa = \frac{\|f'g'' - g'f''\|}{\|(f')^2 + (g')^2\|^{3/2}}.$$

■ Question 61.

Find the curvature $\kappa(x)$ of the graph of $y = f(x)$, parametrized as $\vec{r}(x) = \langle x, f(x) \rangle$.

Use the formula to find the curvature of the graph of $y = x^2$ at the points $(0, 0)$, $(1, 1)$, and $(10, 100)$.

■ Question 62.

(a) Show that the curvature of a straight line is 0.

(b) Show that the curvature of a circle of radius a in \mathbb{R}^2 is $\frac{1}{a}$.

§8.4 The Osculating Plane

Having established a geometric interpretation of the curvature of a circle (it's the reciprocal of the radius), we might guess that the curvature of a general space curve is equal to the curvature of a circle that best approximates the curve at a given point. However, a circle is plane curve; and so we need to first establish which plane the 'best approximation circle' lies in.

Now we would definitely like the circle to be also tangent to the tangent vector of the curve. So the question is to determine another vector, that together with $\vec{T}(t)$ will span the plane for the circle to lie in. Calculations would be easier if we could also choose this second vector to be perpendicular to $\vec{T}(t)$.

Note that at a given point on a smooth space curve $\vec{r}(t)$, there are many vectors that are perpendicular to the unit tangent vector $\vec{T}(t)$. We will single out one as follows. Since $\|\vec{T}(t)\| = 1$ for all t , we have $\vec{T}(t) \cdot \vec{T}'(t) = 0$. So $\vec{T}'(t)$ is perpendicular to $\vec{T}(t)$. This $\vec{T}'(t)$ is not necessarily a unit vector. But, we can define the following:

Definition 8.4.15

At any point where $\kappa \neq 0$ the principal **unit normal vector** $\vec{N}(t)$ is defined as

$$\vec{N}(t) = \frac{\vec{T}'(t)}{\|\vec{T}'(t)\|}$$

We can think of the unit normal vector as indicating the direction in which the curve is turning at each point.

Definition 8.4.16

The **osculating plane** at a point P on the curve traced out by \vec{r} is the plane through P determined by the vectors \vec{T} and \vec{N} at P.

It is the plane that comes closest to containing the part of the curve near P.


Note: For a two-dimensional curve, the osculating plane is identical to the plane of the curve.

Definition 8.4.17

The **osculating circle** (or **circle of curvature**) of a space curve C at a point P on the curve is the unique circle having these properties:

- it lies in the osculating plane at P.
- it lies on the concave side of C at P (its center lies in the direction of \vec{N}).
- it has the same tangent at P as C does.
- it has the same curvature κ at P as C does (its radius is $1/\kappa$).

If \mathcal{O} is the osculating circle of a space curve at the point P, then the radius of \mathcal{O} is called the **radius of curvature at P** and the center of \mathcal{O} is called the **center of curvature at P**.

**Exploration Activity** →

Watch this video for a visual demonstration: <https://www.youtube.com/watch?v=s16ka6HeUdY>

■ Question 63.

Find the equation of the osculating plane of the helix $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$ at the point P = (0, 1, $\pi/2$). □

§8.5 Acceleration

In the previous section we studied curvature, a property of the underlying curve, independent of the vector parametrization \vec{r} . Now we study the parametrization itself and how it determines change in the speed and direction of a particle moving according to \vec{r} .

The **acceleration vector** is given by

$$\vec{a}(t) = \frac{d}{dt} \vec{v}(t) = \vec{r}''(t) = x''(t)\hat{i} + y''(t)\hat{j} + z''(t)\hat{k}$$

So we have an algebraic formula, now let's try to think of a geometric interpretation of the acceleration vector. The velocity vector is in the direction of the tangent. What does the acceleration vector point point toward?

■ Question 64.

□

Recall that $\vec{v}(t) = \|\vec{v}(t)\| \vec{T}(t)$, a product of a scalar-valued and a vector-valued function. Use the product rule to show that $\vec{a}(t)$ can be written as a combination of $\vec{T}(t)$ and $\vec{N}(t)$.

What do each of the components physically signify? A moving particle may be *changing speed* as well as *changing direction*; that is, $\vec{v}(t)$ may change in magnitude and in direction. Information on both of types of change in motion is conveyed in the acceleration vector $\vec{a}(t)$.

Definition 8.5.18

The **tangential and normal components of acceleration**, a_T and a_N respectively, are the real-valued functions satisfying the equation:

$$\vec{a}(t) = a_T(t) \vec{T}(t) + a_N(t) \vec{N}(t)$$

Note:

- The “tangential component” a_T is the rate at which the speed $v(t) = \|\vec{v}(t)\|$ changes.
 - ▶ If a_T is positive, then the particle’s speed is increasing.
 - ▶ If a_T is negative then the particle’s speed is decreasing.
- The “normal component” a_N describes how the velocity changes due to a change *in direction*.

■ Question 65.

□

What is a_N for a particle moving in a straight line?

■ Question 66.

□

Show that

$$a_T = \vec{a} \cdot \vec{T} = \frac{\|\vec{a} \cdot \vec{v}\|}{\|\vec{v}\|}$$

and

$$a_N = \frac{\|\vec{a} \times \vec{v}\|}{\|\vec{v}\|} = \kappa(t) \cdot \|\vec{v}(t)\|^2$$

■ Question 67.

□

Let $\vec{r}(t) = \langle 2t, t^2, \frac{1}{3}t^3 \rangle$.

- (a) Find the tangential and normal components of the acceleration vector.
- (b) Is the speed of the particle increasing or decreasing at $t = 1$?
- (c) Find the decomposition $\vec{a}(t) = a_T(t)\vec{T}(t) + a_N(t)\vec{N}(t)$ at the point where $t = 1$.

Part II

Multivariable Functions and their Derivatives

Chapter 9 | Functions of Two Variables



§9.1 Examples of functions of two variables

- Household lobster consumption is a function of income and the price of lobster.
- The density of cars along a highway is a function of position and time.
- The daily temperature across the United States is a function of latitude and longitude.
- Volume of a cylinder is a function of its radius and height.

§9.2 Representations of two-variable functions

9.2.1 Numerical

The body mass index (BMI) is a value that attempts to quantify a person's body fat based on their height h and weight w .

		Weight w (lbs)				
		120	140	160	180	200
Height h (inches)	60	23.4	27.3	31.2	35.2	39.1
	63	21.3	24.8	28.3	31.9	35.4
	66	19.4	22.6	25.8	29.0	32.3
	69	17.7	20.7	23.6	26.6	29.5
	72	16.3	19.0	21.7	24.4	27.1
	75	15.0	17.5	20.0	22.5	25.0

■ Question 68.

What is the BMI of a person who is 72 inches tall and weighs 160 lb?



9.2.2 Algebraic

A solid cylinder with closed ends has radius r and height h . Its volume V is given by

$$V = f(r, h)$$

■ Question 69.

Write a formula for f . What does $V(10, h)$ mean? What does $V(r, 10)$ mean?



■ Question 70.

Can you give a formula for the function $f(r, h)$? What about the surface area $A = g(r, h)$?



9.2.3 Visual - Using Graphs

Before we give the visual description, let's answer the following question first:

■ Question 71.

What is the domain of the function $f(x, y) = \frac{\sqrt{y^2 - x}}{\ln(x - 2)}$?

The **graph** of a function of two variables, f , is the set of all points (x, y, z) such that $z = f(x, y)$ for all (x, y) in the domain of f . In general, the graph of a function of two variables is a surface in 3-space.

■ Question 72.

Sketch the graph of the function $f(x, y) = 6 - 3x - 2y$.

■ Question 73.

Sketch the graph of $g(x, y) = \sqrt{9 - x^2 - y^2}$.

§9.3 Analyzing Graphs using Cross-sections

One way to visualize surfaces that are three-dimensional graphs is to view pieces of them as two-dimensional graphs. If we intersect the graph of $z = f(x, y)$ with a plane (such as $x = k$ or $y = k$), we get a graph in a two-dimensional plane (the kind we're used to). This is called a **cross-section** (or a trace).

■ Question 74.

Describe the cross-sections of the function $g(x, y) = x^2 - y^2$ with y fixed and then with x fixed. Use these cross-sections to describe the shape of the graph of g .





§9.4 Group Problems**■ Question 75.**

Here are six functions whose graphs we ought to be able to visualize in space:

- | | |
|---------------------------------|---|
| (a) $z = f(x, y) = 6 - 3x - 2y$ | (d) $z = f(x, y) = x^2 + y + 1$ |
| (b) $z = f(x, y) = x^2 + y^2$ | (e) $z = f(x, y) = (x - y)^2$ |
| (c) $z = f(x, y) = x^2 - y^2$ | (f) $z = f(x, y) = \frac{1}{1+x^2+y^2}$ |

Here are cross-sections for two of the surfaces above. Your job is to:


- identify which graph these cross-sections belong with,
- graph cross-sections of the remaining graphs of surfaces, and
- try to visualize the original surface.



■ Question 76.

□

Now that you have a good idea of what each of these graphs look like, you should have no problem identifying which of the following (axes-less) graphs go with each equation for the previous page. Your reasoning should involve the cross-sections you drew as well.

**■ Question 77.**

□

Match each function with its graph.



$$(a) \quad f(x, y) = \frac{x^2}{x^2+y^2}$$

$$(c) \quad f(x, y) = \sin(x + y)$$

$$(b) \quad f(x, y) = y \sin x$$

$$(d) \quad f(x, y) = \sin x \cos y$$

$$(e) \quad f(x, y) = xe^{-xy}$$



Chapter 10 | Contour Plots - Level Curves and Level Surfaces




§10.1 Level Curves

10.1.1 Definition


The **level curves** of a function f of two variables are the curves with equations $f(x, y) = c$, where c is a constant (in the range of f).

These are essentially the z -cross-sections of the graph of $f(x, y)$. The collection of all the level curves is called a **contour plot**.



10.1.2 Basic Facts

- The level curve for fixed output c is called the c -level of $f(x, y)$.
- unless otherwise indicated, contours are drawn at regular z -increments.
- We can build contour plots by hand following the same procedure we used for x and y cross-sections, but now fixing the z -values and plotting in the xy -plane.
- Two different level curves cannot cross.



Sea Surface Temperature (SST) Contour Chart from NOAA

Definition 10.1.19

A function of two variables of the form

$$f(x, y) = mx + ny + d$$

where m, n and d are fixed constants is called a linear functions.


Recall that the equation $z = mx + ny + d$ represents a plane in three-dimensions. Thus clearly, the graph of a linear function looks like a plane. The constant m and n respectively represent the slope of the graph in x -direction and y -direction.

What does the contour plot of a linear function look like? If we set $f(x, y) = c$, we can rewrite the equation as

$$y = -\frac{m}{n}x + \frac{c-d}{n}$$

which is a line with slope $-\frac{m}{n}$. So no matter what level c we choose, the lines remain parallel. Thus the contour plot of a linear function is a set of evenly-spaced parallel lines (assuming regular c -increments).

Note: Consider two functions $f(x, y) = x - y$ and $g(x, y) = (x - y)^3$. Observe the difference in the contour plots for each of the graphs below.





■ Question 78.

Consider the function $z = f(x, y) = 8x - 4y + 2$.

- Sketch the contour plot for the graph with z -increment value of 1.
- (i) Starting at any point (x, y) , what is the slope of the surface in the x -direction?
- (ii) What is the slope in y -direction?
- (iii) What is the slope along the line $x = y$?


■ Question 79.

Find the linear functions whose contour plots are shown below.



■ Question 80.

Assuming regular z -increments, describe what the corresponding graphs might look like.




§10.2 Functions of three variables - Level surface

It's very difficult to visualize a function f of three variables by its graph, since that would lie in a four-dimensional space. However, we do gain some insight into f by examining its **level surfaces**, which are the surfaces with equations $f(x, y, z) = c$, where c is a constant. If the point (x, y, z) moves along a level surface, the value of $f(x, y, z)$ remains fixed.

10.2.1 Level surfaces of $f(x, y, z) = x^2 + y^2 + z^2$

The level surfaces are $x^2 + y^2 + z^2 = k$, where $k \geq 0$. These form a family of concentric spheres with radius \sqrt{k} . Thus, as (x, y, z) varies over any sphere with center O the value of $f(x, y, z)$ remains fixed.



■ Question 81.

3

What do the level surfaces of $f(x, y, z) = x^2 + y^2 - z^2$ look like?

Exploration Activity 

Functions of any number of variables can be considered. A function of n variables is a rule that assigns a number $z = f(x_1, x_2, \dots, x_n)$ to an n -tuple (x_1, x_2, \dots, x_n) of real numbers. We denote by \mathbb{R}^n the set of all such n -tuples.

Recall that there is a one-to-one correspondence between points (x_1, x_2, \dots, x_n) in \mathbb{R}^n and their position vectors $\mathbf{x} = \langle x_1, x_2, \dots, x_n \rangle$. So we have three ways of looking at a function f defined on a subset of \mathbb{R}^n :

- (a) As a function of n real variables x_1, x_2, \dots, x_n
- (b) As a function of a single point variable (x_1, x_2, \dots, x_n)
- (c) As a function of a single vector variable $\vec{x} = \langle x_1, x_2, \dots, x_n \rangle$

We will see that all three points of view are useful.

Chapter II | Partial Derivatives



§II.1 Motivation

Consider the contour plot for the function $f(x, y) = x^2 + y$.

- Sketch the cross-section of the graph with the plane $x = 4$.
- Compute the rate of change of z with respect to y as (x, y) moves towards increasing y -value, along the line $x = 4$.
- What happens to the rate of change of z with respect to x as you move from $(4, 5)$ towards increasing x -value along the line $y = 5$.

§II.2 Definition

For a two-variable function $f(x, y)$ we define the **partial derivative** with respect to x by

$$f_x(a, b) = \frac{\partial f}{\partial x} = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$$

and similarly the **partial derivative** with respect to y by


$$f_y(a, b) = \frac{\partial f}{\partial y} = \lim_{h \rightarrow 0} \frac{f(a, b+h) - f(a, b)}{h}$$

In other words, $f_x(a, b)$ is the derivative at a of the single variable function $x \mapsto f(x, b)$ with $y = b$ fixed.

■ Question 82.

□

What are the values $f_x(0, 0)$ and $f_y(0, 0)$ for the linear function $f(x, y)$ whose contour plot looks like



§II.3 Partial derivative function

We can think of partial derivatives as functions of the base point.

- $f_x(x, y)$ is the function giving f 's rate of change in the x -direction from any point (x, y) .
- $f_y(x, y)$ is the function giving f 's rate of change in the y -direction from any point (x, y) .

Example 11.3.20

If $f(x, t)$ is the vertical height of a guitar string at time t and position x units from one end of the guitar,

- $f_x(x, t)$ measures the rate of change of string height as position changes, or the slope of the string "snapshot" at time t .
- $f_t(x, t)$ measures the rate of change of string height as time passes, or the velocity of the string at position x .

■ Question 83.

For each of the following functions, compute both first partial derivatives f_x and f_y (or f_t):

- $f(x, y) = e^x \cos y$
- $f(x, y) = x^3 - 3xy^2$
- $f(x, t) = e^{-(x+t)^2}$
- $f(x, t) = \sin(x - t) + \sin(x + t)$

■ Question 84.

Let $F(M, r) = \frac{GM}{r^2}$ denote the gravitational force experienced by a unit mass at distance r from the center of a planet with mass M . Calculate $\frac{\partial F}{\partial M}$ and $\frac{\partial F}{\partial r}$.

What does the partial derivative $\frac{\partial F}{\partial M}$ say about how the gravitational force changes when the mass of the planet increases? What does the partial derivative $\frac{\partial F}{\partial r}$ say about how the gravitational force changes when the distance from the planet increases?

§II.4 Higher order partials

We can compute higher order derivatives by simply repeating the process. For example,

$$f_{xy} = (f_x)_y = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial y \partial x}$$

$$f_{xx} = (f_x)_x = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2}$$

Theorem 11.4.21: Clairaut's Theorem

If f_{xy} and f_{yx} are both continuous, then $f_{xy} = f_{yx}$.

■ Question 85.


Use Clairaut's Theorem to compute the requested derivatives of the following functions:

(a) f_{xyxyxy} if $f(x, y) = x^2 \cos(e^y + y^2)$

(b) f_{xxxxyy} if $f(x, y) = x^3 y^2 - \frac{y}{x + \ln(x)}$

■ Question 86.

Below is the contour plot of a function $f(x, y)$. The value of f on each level set is labeled. Assume that the other level curves are of similar shapes.



Based on the diagram, decide whether each of the statements should be true or false. (For which can you be totally sure, and for which would you need more information to be totally sure?)

(a) $f_x(a_1, b_1) \geq 0$

(b) $f_y(a_2, b_2) \geq 0$

(c) $f_x(a_1, b_1) \geq f_x(a_2, b_2)$

(d) $f_{xx}(a_2, b_2) \geq 0$

(e) $f_{xy}(a_2, b_2) \geq 0$

§III.5 Interpreting second partial derivatives

■ Question 87.

Go back to our example of a guitar string above. Let $f(x, t)$ be the vertical height of a guitar string at time t and position x units from one end of the guitar.


- What does f_{xx} represent?
- What does f_{yy} represent?
- What do f_{xt} and f_{tx} represent?

Chapter 12 | Tangent Plane and Linear Approximation




§12.1 Tangent Plane

We have seen earlier that we can view the graph of a function $z = f(x, y)$ as a surface in 3D. If we zoom in on the graph, at most points it seems to flatten out and become planar. Similarly, if we zoom in at the contour plot of $f(x, y)$, the contours look more like equally spaced parallel lines, which are the contours of a linear function. (As we zoom in, we have to add more contours.)



Seeing a plane when we zoom in at a point tells us (provided the plane is not vertical) that $f(x, y)$ is closely approximated near that point by a **linear** function $L(x, y)$. Graphically, the graph of this linear function is a plane, which we call the **tangent plane** at that point.



■ Question 88.

We are going to find the formula of the tangent plane at the point $(a, b, f(a, b))$ on the graph $z = f(x, y)$.

- Write this graph as a level curve of the function of three variables $G(x, y, z) = f(x, y) - z$. What is the 3D gradient of G ?
- Use this to write down the formula for the tangent plane. This is the plane through $(a, b, f(a, b))$ and perpendicular to ∇G .

■ **Question 89.**

Find the equation of the tangent plane to the elliptic paraboloid $z = x^2 + 2y^2$ at the point $(x, y, z) = (1, 1, 3)$.

§12.2 Linear Approximation


Since the tangent plane function matches f (and its partial derivatives) at (a, b) and is a linear function, it is called a **local linearization** of f near (a, b) . We can use the local linearization to approximate a differentiable function at a point.

Theorem 12.2.22

The local linearization of a differentiable function $f(x, y)$ at a point (a, b) is given by

$$L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

Here is picture that explains above theorem graphically.



■ **Question 90.**

- Approximate the values of $z = f(x, y) = x^2 + 2y^2$ at the points $(x, y) = (0.9, 1.1)$ and $(x, y) = (0.95, 0.95)$ using the linearization from the previous exercise.
- Calculate the actual values of $f(0.9, 1.1)$ and $f(0.95, 0.95)$. Compare them to your answer from last part. Could you have predicted whether your estimate was an over-estimate or an under-estimate before calculating the actual values?

■ **Question 91.**

Approximate the value of $f(x, y) = ye^{xy}$ at the point $(x, y) = (0.90, 0.01)$. Which point should you chose to find the local linearization?

Chapter 13 | Directional Derivative and Gradient



Definition 13.0.23

The **directional derivative** of a function $f(x, y)$ in the direction of a unit vector \vec{u} is the rate of change of the value of f in the direction of \vec{u} and is denoted by $D_{\vec{u}}f$.

Suppose we start at (a, b) on the XY-plane and move in the direction of the unit vector \vec{u} in the XY-plane. We want to compare the slope of the graph of f above points along the path. To do this, we consider different contractions h of the unit vector $\vec{u} = u_1\hat{i} + u_2\hat{j} = \langle u_1, u_2 \rangle$ and find the function value at the end of such a contraction $f(a + hu_1, b + hu_2)$.

The average rate of change of $f(x, y)$ between the two points (a, b) and $(a + hu_1, b + hu_2)$ is given by the difference quotient

$$\frac{\Delta f}{\text{displacement}} = \frac{f(a + hu_1, b + hu_2) - f(a, b)}{\|h\vec{u}\|} = \frac{f(a + hu_1, b + hu_2) - f(a, b)}{h}$$

and so to get the instantaneous rate of change we calculate the limit as $h \rightarrow 0$. Thus,

$$D_{\vec{u}}f(a, b) = \lim_{h \rightarrow 0} \frac{f(a + hu_1, b + hu_2) - f(a, b)}{h}$$

Now recall the differential dz or $df = f(x, y) - f(a, b)$ is in fact equal to $f_x(a, b)(x - a) + f_y(a, b)(y - b)$. This is because the instantaneous rate of change in f is the same as the instantaneous rate of change of the linearization $L(x, y)$ of f at the point P (why?). So

$$\begin{aligned} D_{\vec{u}}f(a, b) &= \lim_{h \rightarrow 0} \frac{f_x(a, b)hu_1 + f_y(a, b)hu_2}{h} \\ &= f_x(a, b)u_1 + f_y(a, b)u_2 \\ &= \langle f_x(a, b), f_y(a, b) \rangle \cdot \vec{u} \end{aligned}$$

The **two-dimensional** vector $\langle f_x(a, b), f_y(a, b) \rangle$ is special for its own reasons (as we will see next), and deserves its own name: we call it the **gradient** of f at (a, b) . In general ∇f is also denoted by ∇f and is equal to

$$\nabla f = \langle f_x, f_y \rangle$$

Theorem 13.0.24

The directional derivative of f at (a, b) in the direction of unit vector \vec{u} is given by

$$D_{\vec{u}}f(a, b) = \nabla f(a, b) \cdot \vec{u}$$

■ Question 92.

Find the directional derivative of the function $f(x, y) = x^2 + 2y^2$ in the direction of the vector $\langle 1, 1 \rangle$.



§13.1 Significance of the gradient

We observe that

$$D_{\vec{u}} f(a, b) = \nabla f(a, b) \cdot \vec{u} = \|\nabla f(a, b)\| \cos(\theta)$$

where θ is the angle between ∇f and \vec{u} , since $\|\vec{u}\| = 1$. Hence starting at the point (a, b) , the rate of change of $f(x, y)$ i.e. $D_{\vec{u}} f(a, b)$ is maximized when $\theta = 0$ i.e. when \vec{u} and ∇f are parallel.

In other words,

Theorem 13.1.25

$D_{\vec{u}} f(a, b)$ is maximized in the direction of ∇f . Moreover, the maximum value of $D_{\vec{u}} f(a, b)$ is equal to $\|\nabla f\|$.

■ Question 93.

Suppose we drop some water on the surface $z = x^3y + y^2x$ at the point $(1, 1, 2)$. Which way does the water flow?

■ Question 94.

True or False: If we know the value of the directional derivative $D_{\vec{u}} f(a, b)$ in the direction of any two nonparallel different unit vectors, then we can determine $\nabla f(a, b)$.

§13.2 Gradient and Level Curves

What happens to $f(x, y)$ along a level curve of f ? If \vec{u} is the direction of the tangent at the point $P = (a, b)$ to the level curve passing through P , then the rate of change of f in the direction of \vec{u} is zero.

Hence for this choice of \vec{u} , we have

$$D_{\vec{u}} f(a, b) = 0 \implies \theta = \frac{\pi}{2}$$

where θ is the angle between \vec{u} and $\nabla f(a, b)$. In other words,

Theorem 13.2.26

The gradient vector at a point is perpendicular to the level curve through that point.

Chapter 14 | Three Dimensional Gradient and the Tangent Plane



The directional derivative formula and the gradient have the obvious extensions for functions of three-variables $f(x, y, z)$:

$$\begin{aligned} D_{\vec{u}} f(a, b, c) &= \nabla f(a, b, c) \cdot \vec{u} \\ &= \langle f_x(a, b, c), f_y(a, b, c), f_z(a, b, c) \rangle \cdot \vec{u} \\ &= \|\nabla f(a, b, c)\| \cos(\theta) \end{aligned}$$

so the gradient again points in the direction of greatest increase of f from (a, b, c) and has magnitude equal to that rate of change.

■ Question 95.

A fly is flying around a room in which the temperature is given by $T(x, y, z) = x^2 + y^4 + 2z^2$. The fly is at the point $(1, 1, 1)$ and realizes that he's cold. In what direction should he fly to warm up most quickly?

■ Question 96.

How should the gradient $\nabla f(a, b, c)$ be related to the level-surface of f through (a, b, c) ?

■ Question 97.

For each of the following functions $F(x, y, z)$, do the following

- Compute the gradient ∇F .
- Identify the level surface $F(x, y, z) = \text{constant}$ through the point $(x, y, z) = (1, 1, 1)$.
- Find the tangent plane to the level surface from part (b) at $(1, 1, 1)$.

$$(I) F(x, y, z) = 3x + 2y + z \quad (II) F(x, y, z) = x^2 + y^2 - z^2$$

§14.1 3D Gradient as the Normal vector

Since the gradient $\nabla f(a, b, c)$ is perpendicular to the level surface of f containing (a, b, c) , it is the normal vector to the tangent plane of the level surface at (a, b, c) . So even if the surface is not the graph of a function, we can use the three dimensional gradient to find tangent planes.

■ Question 98.

Find the equation of tangent plane to the surface $xy + yz + zx = 5$ at the point $(1, 1, 2)$.

■ Question 99.

□

Alice the “A” student is debating Chuck the “C” student. Alice says that the direction of greatest ascent on the graph $z = f(x, y)$ is in the direction ∇f . Chuck says that instead we should look at the level surface $F(x, y, z) = z - f(x, y) = 0$ and go in direction ∇F . Who is right? How would you explain this to the student that is wrong?

■ Question 100.

□

You’re hiking a mountain which is the graph of $f(x, y) = 15 - x^2 - 2xy - 3y^2$. You’re standing at $(1, 1, 9)$. You wish to head in a direction which will maintain your elevation (so you want the instantaneous change in your elevation to be 0). How many possible directions are there for you to head? What are they?

■ Question 101.

□

Show that every plane that is tangent to the cone $x^2 + y^2 = z^2$ passes through the origin.

■ Question 102.

□

Where does the normal line to the paraboloid $z = x^2 + y^2$ at the point $(1, 1, 2)$ intersect the paraboloid a second time?

■ Question 103.

□

Find the tangent plane to the surface of the solid described by inequalities $0 \leq x \leq 6$ and $0 \leq y^2 + z^2 \leq 4$ at the point $(6, 1, 1)$.

■ Question 104.

□

The ellipsoid $x^2 + 4y^2 + 9z^2 = 36$ and the surface $z = \sin[\pi(x - y)]$ intersect in a curve, call it \mathcal{C} . Find the line tangent to \mathcal{C} at the point $(6, 0, 0)$. (Please give a parametric vector equation for the line.)

■ Question 105.

□

Consider the surface

$$\sqrt{x} + \sqrt{y} + \sqrt{z} = \sqrt{c}.$$

- (a) Find the equation of the tangent plane to this surface at a point (p, q, r) .
- (b) Show that the sum of X-intercept, Y-intercept and Z-intercept of the above tangent plane does not depend on p, q , and r .

Chapter 15 | Chain Rule



Recall from one-variable calculus the chain rule

$$\frac{d}{dt} f(g(t)) = f'(g(t)) \cdot g'(t)$$

which is stated as “the derivative of the outside function (evaluated at inside function), times the derivative of the inside function”.

Now think of a particle is moving in a parameterized curve $\vec{r}(t) = \langle x(t), y(t) \rangle$ and we want to understand how some function $f(x, y)$ changes along the curve. For example, we might ask, what is the rate of change of height along a particular racetrack. For any composite function of the form $f(x(t), y(t)) = f(\vec{r}(t))$, the multivariable chain rule is


$$\frac{dz}{dt} = \frac{d}{dt} f(\vec{r}(t)) = \nabla f(\vec{r}(t)) \cdot \vec{r}'(t) = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle \cdot \langle x'(t), y'(t) \rangle$$

where the gradient ∇f acts like the derivative of the outside function f and the velocity $\vec{r}'(t)$ acts like the derivative of the inside vector-function $\vec{r}(t) = \langle x(t), y(t) \rangle$.

Warning: Notice that this just like the formula for the directional derivative, but now with $\vec{r}'(t)$ playing the role of the direction \vec{u} . Thus we can think of $\frac{df}{dt}$ as (a multiple of) the rate of change of f in the tangent direction of the parameterized motion.

§15.1 Chain Rule diagram

A chain of dependence diagram is a convenient way to represent the chain rule for $z = f(x(t), y(t))$.




Then, the derivative $\frac{dz}{dt}$ is obtained by adding the products of the derivatives on each path of the diagram leading from z to t :

$$\begin{aligned}\frac{dz}{dt} &= \frac{\partial z}{\partial x}x'(t) + \frac{\partial z}{\partial y}y'(t) \\ &= \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt}\end{aligned}$$

where the terms $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ are both evaluated at $(x(t), y(t))$.

§15.2 More complicated diagram

Curves in the xy -plane can also sometimes be described by even more parameters, like $x(u, v)$ and $y(u, v)$. Then the chain rule for differentiating $z = f(x(u, v), y(u, v))$ is even more complicated; but a dependence diagram can help.



Now the partial derivative $\frac{\partial z}{\partial u}$ is obtained by adding the products of the derivatives on each path from z to u in the diagram:

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial u}$$

and the partial derivative $\frac{\partial z}{\partial v}$ is obtained by adding the products of the derivatives on each path from z to v in the diagram:

$$\frac{\partial z}{\partial v} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial v} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial v}$$

■ Question 106.

Let $w = x^2 e^y$, $x = 4u$, and $y = 3u^2 - 2v$. Compute $\frac{\partial w}{\partial u}$ and $\frac{\partial w}{\partial v}$ using the chain rule.

■ Question 107.

Let $z = f(x, y) = x^2 + y^3$, and $x = x(s, t)$ and $y = y(s, t)$; i.e., x and y are functions of s and t . Suppose that when $(s, t) = (0, 1)$, we have:

$$x(0, 1) = -1, \quad x_s(0, 1) = -4, \quad x_t(0, 1) = -7, \quad y(0, 1) = 2, \quad y_s(0, 1) = 10, \quad y_t(0, 1) = 5.$$

Compute $\frac{\partial z}{\partial t}$ at $(s, t) = (0, 1)$.

■ Question 108.

The length of a side of a triangle is increasing at a rate of 3 in/s, the length of another side is decreasing at a rate of 2 in/s, and the contained angle Θ is increasing at a rate of 0.05 radian/s. How fast is the area of the triangle changing when $x = 40$ in, $y = 50$ in, and $\Theta = \pi/6$?

■ Question 109.

Let $p = g(u, v)$ be a differentiable function of two variables. Let $u = \frac{x}{y}$ and $v = \frac{y}{z}$. Show that

$$x \frac{\partial p}{\partial x} + y \frac{\partial p}{\partial y} + z \frac{\partial p}{\partial z} = 0$$

■ Question 110.

(a) Use the chain rule to compute $\frac{\partial z}{\partial r}$ and $\frac{\partial z}{\partial \theta}$ for $z = f(x(r, \theta), y(r, \theta))$ where

$$x(r, \theta) = r \cos(\theta) \quad y(r, \theta) = r \sin(\theta)$$

(b) Show that

$$\left(\frac{\partial z}{\partial x} \right)^2 + \left(\frac{\partial z}{\partial y} \right)^2 = \left(\frac{\partial z}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta} \right)^2$$

Chapter 16 | Limit, Continuity, and Differentiability



§16.1 Limit and Continuity

The definition of a limit for a function of two variables is pretty much the same as for a single variable function.

Definition 16.1.27

We write

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L$$

if $f(x,y)$ approaches L as (x,y) approaches (a,b) .

The key question here is figuring out what “approaches” means. In one-dimension, x can approach a number c either from left or right. But in two-dimension we can have lots of different paths to “approach” a point (a,b) .




Figure 16.1: Different paths approaching (a,b)

In particular, if $f(x,y) \rightarrow L_1$ as $(x,y) \rightarrow (a,b)$ along a path C_1 and $f(x,y) \rightarrow L_2$ as $(x,y) \rightarrow (a,b)$ along a path C_2 , where $L_1 \neq L_2$, then $\lim_{(x,y) \rightarrow (a,b)} f(x,y)$ does not exist.



Warning: We can only follow paths that are in the domain of our function and that go to the limit point!

In general, it's hard to show algebraically that a limit exists since we must consider infinitely many possible paths to approach (a,b) . If the function is of certain specific types, we can simplify the limit into a one-variable limit and calculate it that way.

Sometimes you can separate the variables to turn the multivariable limit into a single variable limit which means we can use L'Hopital.

Example 16.1.28

Consider $L = \lim_{(x,y) \rightarrow (0,3)} \frac{1 - \cos x}{xy}$.

We can write the limit as the product of two different limits _____.

Now calculating each separately, we get $L =$ _____.

Sometimes limits at a point like $(0,0)$ may be easier to evaluate by converting to polar coordinates. Remember that the same limit must be obtained as $r \rightarrow 0$ regardless of θ .

Example 16.1.29

Consider $L = \lim_{(x,y) \rightarrow (0,0)} \frac{\sin(x^2 + y^2)}{5x^2 + 5y^2}$.

We can rewrite the limit in polar coordinates as _____.

This is a one-variable limit. Hence, $L =$ _____.

For a general function, we will need to use the $\varepsilon - \delta$ definition. It is comparatively easier to show that a limit does not exist.

Example 16.1.30

We will show that $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$ does not exist.

First let's approach $(0,0)$ along the x -axis. Along this path $y =$ _____.

So the function becomes $f(x,y) =$ _____, and hence the limit is _____.

Next, approach $(0,0)$ along the y -axis. Along this path $x =$ _____.

So the function becomes $f(x,y) =$ _____, and hence the limit is _____.

Since f has two different limits as (x,y) approaches $(0,0)$ along two different curves, the given limit does not exist.

■ Question 111.

Show that $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$ does not exist by consider approaches along $x = 0$ and $x = y$. □

■ Question 112.

Show that $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$ does not exist by consider approaches along the line $y = x$ and the parabola $x = y^2$. □

§16.2 Continuity

We define continuity for functions of two variables in the same way as for functions of one variable:

Definition 16.2.31

We say that a function f is **continuous at the point** (a, b) if

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

A function is **continuous on a region** R in the xy -plane if it is continuous at each point in R .

Theorem 16.2.32

If $f(x, y)$ is a polynomial or a rational function $\frac{p(x, y)}{q(x, y)}$ where $q(x, y) \neq 0$, then f is continuous.

Example 16.2.33

The function $f(x, y) = \frac{x^2 - y^2}{x^2 + y^2}$ is clearly discontinuous at the origin (since it's not even defined there). But as a rational function, f is continuous everywhere else.

Just as for functions of one variable, composition of two continuous functions is continuous.

■ Question 113.

Where is the function $h(x, y) = e^{-(x^2+y^2)}$ continuous?

§16.3 Differentiability

First of all, recall that in the case of one-variable functions, the derivative is equal to slope of the tangent line. Analogously, we say a function of two-variables is differentiable at a point if the function is “well-approximated” by the tangent plane at the point.

From theorem 22, the tangent plane is given by the local linearization of $f(x, y)$ at (a, b)

$$L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b).$$

So we will focus on the precise meaning of the phrase “well approximated.” Let’s define the error function $E(x, y) = f(x, y) - L(x, y)$. We will define differentiability at a point in terms of this error and the distance from nearby points.

Consider a nearby point $(a + h, b + k)$. The distance between this point and (a, b) is $\sqrt{h^2 + k^2}$.

Definition 16.3.34

We say that the function $f(x, y)$ is differentiable at the point (a, b) if

$$\lim_{(h,k) \rightarrow (0,0)} \frac{E(a+h, b+k)}{\sqrt{h^2 + k^2}} = 0.$$

The function f is differentiable on a region R if it is differentiable at each point of R .

■ Question 114.

□

Consider the function $f(x, y) = \sqrt{x^2 + y^2}$. Show that f is not differentiable at the origin by showing that above limit does not exist.

HINT: What happens if we approach along the path $y = x$.

Theorem 16.3.35

If $f_x(x, y)$ and $f_y(x, y)$ both exist and are both continuous on a small disk R centered around the point (a, b) , then $f(x, y)$ is differentiable at (a, b) .

Note: Just the existence of both partial derivatives at a point does not guarantee differentiability of the function there. For example, consider the function

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$$

We have already seen that the function is not continuous at the origin. However, we can verify that both partial derivatives exist at $(0, 0)$.

■ Question 115.

□

Show that $f(x, y) = xe^{xy}$ is differentiable at $(1, 0)$.

Part III

Application of Derivatives - Optimization

Chapter 17 | Local Optimization





Definition 17.0.36

Suppose f is a continuous function on its domain D . Let P_0 be a point in D . Then we say f has

- a local maximum at the point P_0 if $f(P_0) \geq f(P)$ for all points P near P_0 .
- a local minimum at the point P_0 if $f(P_0) \leq f(P)$ for all points P near P_0 .
- a global (absolute) maximum at the point P_0 if $f(P_0) \geq f(P)$ for all points P in the domain of f .
- a global (absolute) minimum at the point P_0 if $f(P_0) \leq f(P)$ for all points P in the domain of f .

§17.1 How to detect a local extremum from the Contour Plot?

Consider the graph and contour plot of a function $z = f(x, y)$ as follows:




■ Question 116.

What does the contour plot look like around a local extremum? What can you say about the partial derivatives and the gradient at the local extremum? □

Use similar logic to answer the following question.

■ Question 117.

A function $f(x, y)$ of two variables has level curves as shown in the picture. Some of the gradient vectors are also given for your convenience. Fill in the table below with all appropriate choices of points. □



Enter all possible choices from A-I	is/are point(s) where
	$f_x(x, y) = 0 \text{ and } f_y(x, y) \neq 0$
	$f_y(x, y) = 0 \text{ and } f_x(x, y) \neq 0$
	$f_y(x, y) = 0 \text{ and } f_x(x, y) = 0$
	$f(x, y)$ has either a local maxima or a local minima.

§17.2 Critical Points

Definition 17.2.37

A point (a, b) is called a **critical point** for the function $z = f(x, y)$ if it is one of the following two types.

- **Stationary Point:** an interior point at which f is differentiable and $f_x(a, b) = 0$ and $f_y(a, b) = 0$.
- **Singular Point:** an interior point at which f is not differentiable.

A point (a, b) is called a **stationary point** for the function $z = f(x, y)$ iff That is,

From the table above, it should be clear that a local maxima or local minima is a critical point, but not all critical points are local extremum.


Note: Stationary points are the points where $\nabla f = \langle f_x, f_y \rangle = 0$. If f is differentiable everywhere, critical points and stationary points are the same things.

■ Question 118.

Find the critical points of the function $f(x, y) = 8y^3 + 12x^2 - 24xy$. Are all of them local extrema?

Here is a contour plot to help you.

□


Definition 17.2.38

A critical point which is neither a local maximum nor a local minimum is called a saddle point.

§17.3 Classifying Stationary Points - The Second-Derivative test

The **Hessian** of a function $f(x, y)$ is defined to be the matrix $H = \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix}$. Suppose (a, b) is a point where $\nabla f(a, b) = \vec{0}$ and let

$$D = \det(H(a, b)) = f_{xx}(a, b)f_{yy}(a, b) - (f_{xy}(a, b))^2$$

The second-derivative test says

Theorem 17.3.39: The Second Derivative test

Suppose the second partial derivatives of f are continuous for all points in an open disk around (a, b) where $\nabla f(a, b) = \vec{0}$.

- If $D > 0$ and $f_{xx}(a, b) > 0$, then f has a local minimum at (a, b) .
- If $D > 0$ and $f_{xx}(a, b) < 0$, then f has a local maximum at (a, b) .
- If $D < 0$, then f has a saddle point at (a, b) .
- If $D = 0$, anything can happen: f can have a local maximum, or a local minimum, or a saddle point, or none of these, at (a, b) .

■ Question 119.

How does the contour plot look like near a saddle point? Identify the saddle points in the plots above.



■ Question 120.

Find and classify all stationary points of the following functions:

- (a) $f(x, y) = x^2 + 2xy + 2y^2 - 8y + 12$
 (b) $f(x, y) = x^2 - 2y^2 + xy - 4$
 (c) $f(x, y) = 8 - x^2 - xy - y^2$
 (d) $f(x, y) = x^4 + y^4 + 4xy + 3$
 (e) $f(x, y) = y^2 + 2xy - x^2 + 2x - 2y + 36$
 (f) $f(x, y) = 5 - x^3 + y^3 + 3xy$

■ **Question 121.**

Two of the above functions are shown as contour plots below. For each plot, predict the location of the stationary points and classify them as maxima, minima, or saddle points. Can you identify which functions are plotted here?





Chapter 18 | Constrained Optimization



§18.1 Motivation

A racehorse lives in a valley which happens to be the graph of $f(x, y) = 3x^2 + y^2$. He is doomed to wander his racetrack, which is the set of points in the valley where $x^2 + y^2 = 1$. The racehorse secretly wishes to be a mountain climber, and his fantasy is to escape from his racetrack and take the steepest path up the mountain. At what point should he make his escape, and in what direction should he run?

Below we have drawn 3-D depiction of the valley and the racetrack in the first picture, and the contour plot of the function $f(x, y)$ along with the projection of the racetrack in the second picture.




It makes sense that the horse will make his escape when $f(x, y)$ is largest along the racetrack. Can you see how to locate those points on the contour plot?

§18.2 Graphical Reasoning

■ Question 122.


Consider in general, the contours of a differentiable function $f(x, y)$ overlaid with a curve $g(x, y) = c$.





- (a) For each plot, locate the local and global maximizer(s) and minimizer(s) of $f(x, y)$ on the curve $g(x, y) = c$.
- (b) How are the tangents to the level curve of f and the curve $g(x, y) = c$ related at the extreme points?
- (c) The curve $g(x, y) = c$ is evidently a level curve for the function $g(x, y)$. What does this imply about the relationship between ∇f and ∇g at the extreme points of f on the curve $g(x, y) = c$?
- [HINT: How is the gradient of any function related to the level curves of that function?]
- (d) Find the other points on the curve $g(x, y) = c$ which satisfy the gradient relationship. What is the behavior of f relative to the curve $g(x, y) = c$ at these points?

§18.3 Algebraic Solution



From the last exercise, we observe that ∇f must be parallel to ∇g at the extreme points (a, b) of f on the curve $g(x, y) = c$. So we should be able to find some scalar λ such that

$$\nabla f(a, b) = \lambda \nabla g(a, b)$$

This is called the **Lagrange multiplier rule** in honor of Joseph-Louis Lagrange, and the particular scalar λ that lines up the two vectors exactly is called the **Lagrange multiplier**.

In the picture above ∇f and ∇g lie on the same line at the extreme points.

18.3.1 Algorithm

Step 1. To find critical points of a function $f(x, y)$ on a constraint curve $g(x, y) = c$, first solve the following system of simultaneous equations for x, y , and λ :

$$\begin{aligned}f_x(x, y) &= \lambda g_x(x, y) \\f_y(x, y) &= \lambda g_y(x, y) \\g(x, y) &= c\end{aligned}$$

Step 2. Once you have found all the critical points (a, b) , evaluate $f(a, b)$ at each of them. The critical points where f is greatest are maxima and the critical points where f is smallest are minima.



Warning: In case of functions of three variables, Lagrange Multiplier problems with one constraint equation means solving a system of four simultaneous equations.

18.3.2 Tips for Solving the System of Equations

Solving the system of equations can be hard! Here are some possible approaches for solving these systems of equations:

- Solve for x, y, z in terms of λ . Then you can plug these back into the constraint, find the value of λ , which will give you x, y, z .
- Or, eliminate λ to solve for one variable in terms of the others. Substitute them in to $g(x, y) = c$ to solve for x and y .
- Remember that whenever you take a square root, you must consider both the positive and the negative square roots.
- Remember that whenever you divide an equation by an expression, you must be sure that the expression is not 0. It may help to split the problem into two cases:
first solve the equations assuming that a variable is 0, and then solve the equations assuming that it is not 0.

■ Question 123.

For each of the following, find the maxima and minima of f , on the constraint curve g .

- $f(x, y) = xy$, $g(x, y) = 3x^2 + y^2 = 6$.
- $f(x, y) = \frac{1}{x} + \frac{1}{y}$, $g(x, y) = \frac{1}{x^2} + \frac{1}{y^2}$
- $f(x, y) = x^2 + 4xy + y^2$, $g(x, y) = x - y - 6$

§18.4 Three-variable Functions

■ Question 124.

Let $T(x, y, z) = x - 2y + 5z + 64$ give the temperature, in degrees Celsius, at the point (x, y, z) , where x, y , and z are measured in meters. Find the maximum and minimum temperature on the sphere $x^2 + y^2 + z^2 = 120$.

Suppose we want to find extreme values of $f(x, y, z)$ and we have more than one constraint, say two constraints $g(x, y, z) = k$ and $h(x, y, z) = c$. So we are looking for the max and min values of f which lie on

the intersection of $g(x, y, z) = k$ and $h(x, y, z) = c$. In this case our ∇f is determined by both ∇g and ∇h . So we now have Lagrange multipliers for each constraint function. So in this example, we have

$$\nabla f = \lambda \nabla g + \mu \nabla h$$

■ Question 125.

Set up, but do not solve, the equations you would solve to find the extreme values of $f(x, y, z) = z$ subject to $x^2 + y^2 = z^2$ and $x + y + z - 24 = 0$

§18.5 Interpretation of λ as a rate of change

The value of λ can be interpreted as the rate of change of the optimum value of f as c increases (where $g(x, y) = c$). Suppose the optimum value of f is obtained at the point (a, b) , where both a and b depend on the choice of c , and suppose the optimum value is given by


$$f_{\min}(c) = f(a(c), b(c))$$

as a function of c . Then

$$\lambda = \frac{d}{dc} f_{\min}(c)$$


■ Question 126.

The image below shows the contours of a function f . For the graphs (a) and (b), Is P a maximizer or minimizer of f subject to $g(x, y) = x + y = c$ with $x, y \geq 0$? What is the sign of λ ? Where could the other extreme be located?



■ Question 127.

The images below show the optimal point (marked with a dot) in three optimization problems with the same constraint. Arrange the corresponding values of λ in increasing order. (Assume λ is positive.)



Chapter 19 | Global Optimization



§19.1 Definitions

- f has a global maximum on a two-dimensional region R at the point P_0 if $f(P_0) \geq f(P)$ for all points $P \in R$.
- f has a global minimum on a two-dimensional region R at the point P_0 if $f(P_0) \leq f(P)$ for all points $P \in R$.

A **Warning:** If the region R is not stated explicitly, we take it to be the whole xy -plane unless the context of the problem suggests otherwise.


§19.2 How do we know whether a function has a global maximum or minimum?

- A **closed** region is one which contains its boundary.
- A **bounded** region is one which does not stretch to infinity in any direction.

Theorem 19.2.40: Extreme Value Theorem

If f is a continuous function on a closed and bounded region R , then f achieves its global maximum and minimum at some points in R .

The global extrema could be on the boundary or in the interior of R .



■ Question 128.

For the function $f(x, y) = (x - 2)^2 + y^2$, find a region R such that

- f attains a global maximum value of 4 and a global minimum value of 0 over R .
- f attains a global maximum value of 3 and a global minimum value of 1 over R .
- f attains a global maximum value of 9 but has no global minimum over R .

§19.3 Algorithm for Global Optimization

■ Question 129.

Consider the function

$$F(x, y) = 3x^2y + y^3 - 3x^2 - 3y^2 + 1$$

defined on the disk D of radius $\sqrt{2}$ centered at the origin i.e.

$$D = \{(x, y) \mid x^2 + y^2 \leq 2\}.$$

Follow the steps below to find the global maximum and minimum of f on D.

- (a) Find all the stationary critical points of F. [HINT: There are 4 such points.]
- (b) In the list of critical points from part (a), identify the ones lying **inside** D (excluding the boundary).
- (c) Evaluate F at the critical point(s) from part (b).
- (d) Use Lagrange multiplier to find the maximum and minimum of F(x, y) subject to the constraint $x^2 + y^2 = 2$. Note that this gives the extreme values of F on the boundary circle of D.
- (e) Evaluate F at the critical point(s) from part (d).
- (f) Compare the extreme values of F from part (c), and the extreme values of F from part (e), to find the global maximum and minimum of F(x, y) on D.

§19.4 Practice Problems

■ Question 130.

For which of the regions D described below is it true that every continuous function $f(x, y)$ must attain an global maximum value and global minimum value on D? (There may be more than one.)

(i) D is the set of points (x, y) such that $|x| \leq 4$ and $|y| < 2$

(ii) D is the set of points (x, y) such that $|x + y| \leq 1$

(iii) D is the set of points (x, y) such that $x^2 + 4y^2 \leq 1$

(iv) D is the set of points (x, y) such that $x^2 + 4y \leq 1$

(v) D is the set of points (x, y) such that $-x \leq y \leq x$

Then do the following:

- (a) For one of the regions D that you picked, find the global minimum and global maximum value of $f(x, y) = x^2 - 4x + y^2$ on the region.
- (b) For one region you didn't pick, find a function $f(x, y)$ which has either no maximum or no minimum on the region D.

■ Question 131.

Find all global extrema of the function

$$f(x, y) = 2x^3 + 2y^3 - 3x^2 - 3y^2 + 6$$

on the disc $D = \{(x, y) \mid x^2 + y^2 \leq 4\}$.

Part IV

Integration of Multivariable Functions

Chapter 20 | Definite Integration in two variables



§20.1 Introduction

In one-variable calculus, we learned that the definite integral $\int_a^b f(x) dx$ gave the area under the graph of $f(x)$ over the interval $[a, b]$ on the x -axis. For two-variable functions, the input space is the xy -plane, so we compute volume integrals over two-dimensional regions R of the plane:

$$\int_R f(x, y) dA$$

which measure the amount of volume under the graph of $f(x, y)$ and over the region R . Note that R is NOT necessarily a rectangle. The dA indicates that the integration is taken over a region R with (A)rea in the xy -plane.

§20.2 Construction

Recall that we approximate definite integrals in one-variable calculus by chopping the interval $[a, b]$ into subintervals and summing the areas of rectangles (usually) based on each subinterval.

For volume integrals, we can chop R into subregions and sum the volumes of rectangular (usually) prisms based on each subregion. As with one-variable integrals, using smaller and smaller subregions of R leads to better and better approximations.




Figure 20.1: Approximating volume under a graph with finer and finer Riemann sums


Exploration Activity

The Riemann Sum definition: If R is a rectangle, we can express this approximation procedure more formally using Riemann sums and limits as follows.

Suppose the function $f(x, y)$ is continuous on R , the rectangle $a \leq x \leq b, c \leq y \leq d$. Consider the partitions

$$P : \{a = x_0 < x_1 < \dots < x_n = b\}$$

and



For a choice of points (x_{ij}^*, y_{ij}^*) in the (i, j) th subrectangle $[x_{i-1}, x_i] \times [y_{j-1}, y_j]$, we define the Riemann Sum of f over R as

$$S_{(P,Q)}^* = \sum_{i=1}^n \sum_{j=1}^m f(x_{ij}^*, y_{ij}^*) \Delta x_i \Delta y_j = \sum_{i=1}^n \sum_{j=1}^m f(x_{ij}^*, y_{ij}^*) (x_i - x_{i-1})(y_j - y_{j-1})$$

Also let $\|(P, Q)\|$ denote the maximum area of a subrectangle i.e.

$$\|(P, Q)\| = \max_{i,j} (x_i - x_{i-1})(y_j - y_{j-1})$$

Then the definite integral of f over R is given by

$$\int_R f dA = \lim_{\|(P, Q)\| \rightarrow 0} S_{(P,Q)}^*$$

In the case the partitions are uniform, above limit can be rewritten as

$$\int_R f dA = \lim_{n,m \rightarrow \infty} S_{(P,Q)}^*$$

■ Question 132.

If R is any region in the plane, what does the double integral $\int_R 1 dA$ represent? Why?

■ Question 133.

Suppose the shape of a flat plate is described as a region R in the plane, and $f(x, y)$ gives the density (mass of unit area) of the plate at the point (x, y) in kilograms per square meter. What does the double integral $\int_R f(x, y) dA$ represent? Why?

■ Question 134.

$\iint_R f(x, y) dA$
 What does the double integral $\frac{\iint_R f(x, y) dA}{\iint_R 1 dA}$ represent? Why?

■ Question 135.



Let R be the rectangle $-1 \leq x \leq 1, -1 \leq y \leq 1$, and T is the top half $-1 \leq x \leq 1, 0 \leq y \leq 1$, and L is the left half $-1 \leq x \leq 0, -1 \leq y \leq 1$.

Without evaluating any of the integrals, decide which of them are positive, which are negative, and which are zero.

- | | | | |
|---------------------------|----------------------------|-----------------------------|------------------------------|
| (i) $\int_R x dA$ | (ii) $\int_R y dA$ | (iii) $\int_T y dA$ | (iv) $\int_R (x - x^2) dA$ |
| (v) $\int_T (y - y^2) dA$ | (vi) $\int_L (x^2 - x) dA$ | (vii) $\int_L (y + y^3) dA$ | (viii) $\int_R (2x + 3y) dA$ |

■ Question 136.



For the same set of integrals above, arrange (a), (c), (e) and (f) in increasing order.

■ Question 137.




Evaluate $\iint_R (f(x, y) dA)$ for the function below

$$f(x, y) = \begin{cases} 2, & 0 \leq x \leq 5, 0 \leq y \leq 1 \\ 3, & 0 \leq x \leq 5, 1 \leq y \leq 3 \end{cases}$$

Chapter 21 | Iterated Integrals



Consider the volume integral of a function f over a rectangular region R . Instead of dicing R into a grid to approximate the integral, we can cut it in “vertical strips” (in the y -direction) in the xy -plane. Then, we build solid slices whose heights are determined by the value of $f(x, y)$ along one edges of each strip.



Volume of such a solid slice is given by

$$\underbrace{\left(\int_c^d f(x_{i-1}, y) dy \right)}_{\text{Area of a vertical cross-section parallel to YZ-plane}} \times \underbrace{(x_i - x_{i-1})}_{\text{width in X-direction}}$$

Now if we sum up the volumes of these solids and take a limit, we should get back the volume integral

$$\int_R f dA = \lim_{\Delta x_i \rightarrow 0} \sum_{i=1}^n \left(\int_c^d f(x_{i-1}, y) dy \right) (x_i - x_{i-1}) = \int_a^b \int_c^d f(x, y) dy dx$$

This is called an **iterated integral**.

§21.1 Type I and Type II regions

A region R is called **Type I** if it can be written in the following way:

$$R = \{(x, y) \mid a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\}$$

We can then compute a double integral as

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$$

A region R is called **Type II** if it can be written in the following way:

$$R = \{(x, y) \mid c \leq y \leq d, h_1(y) \leq x \leq h_2(y)\}$$

We can then compute a double integral as

$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

Note: For a quick reference, the rules on the limits of an iterated integral are as follows:

- The limits on the outer integral must be constants.
- The limits on the inner integral can involve only the variable in the outer integral. For example, if the inner integral is with respect to x , its limits can be functions of y .

■ Question 138.

Match the integral with the appropriate region of integration.

(a) R_1 : The triangle with vertices $(0, 0), (2, 0), (0, 1)$ (c) R_3 : The triangle with vertices $(0, 0), (2, 0), (2, 1)$

(b) R_2 : The triangle with vertices $(0, 0), (0, 2), (1, 0)$ (d) R_4 : The triangle with vertices $(0, 0), (1, 0), (1, 2)$

$$(i) \int_0^1 \int_0^{2-2x} f(x, y) dy dx$$


$$(ii) \int_0^1 \int_0^{2-2y} f(x, y) dx dy$$

$$(iii) \int_0^1 \int_0^{2x} f(x, y) dy dx$$


$$(iv) \int_0^1 \int_{2y}^2 f(x, y) dx dy$$

■ Question 139.


Here are some type I regions. For each of them set up the given double integral as an iterated integral and evaluate it.



$$(i) \iint_{D_1} xy dA$$



$$Ans: 45/8 \quad (ii) \iint_{D_2} e^{x^2} dA$$



$$Ans: (e-1)/2$$

$$(iii) \iint_{D_3} (x-1)y^2 dA$$



$$Ans: 0 \quad (iv) \iint_{D_1} 1 dA$$

$$Ans: 4.5$$

■ **Question 140.**

□

Here are some type II regions. For each of them set up the given double integral as an iterated integral and evaluate it. The curves in D_4 are $x = 2y - y^2$ and $x = y^2 - 2y + 4$.



$$(i) \iint_{D_2} (1-y)^3 dA$$

$$Ans: 1/5 \quad (ii) \iint_{D_4} (y-1)x^2 dA$$

$$Ans: 0$$

$$(iii) \iint_{D_2} \cos(x^2) dA$$

$$Ans: \frac{\sin(1)}{2} \quad (iv) \iint_{D_5} (x-1) dA$$

$$Ans: -2/3$$

§21.2 Switching the Order of Integration

If a region is both type I and type II, you may find that one order of integration will be simpler to deal with than the other. Sometimes, when converting a double integral to an iterated integral, we decide the order of integration based on the integrand, rather than the shape of the region - some integrands are easy to integrate with respect to one variable and much harder (or even impossible) to integrate with respect to the other.

■ **Question 141.**

□

For each of the following integrals, draw the region in question, write down an integral with the reverse order of integration, then finally integrate.

$$(i) \int_0^2 \int_x^2 (x+y) dy dx$$

$$Ans: 4 \quad (ii) \int_0^6 \int_{x/3}^2 x \sqrt{y^3 + 1} dy dx$$

$$Ans: 26$$

$$(iii) \int_0^1 \int_y^1 e^{x^2} dx dy$$

$$Ans: \frac{e-1}{2} \quad (iv) \int_0^{\pi/2} \int_y^{\pi/2} \frac{\sin(x)}{x} dx dy$$

$$Ans: 1$$

$$(v) \int_0^1 \int_{-\sqrt{y}}^{\sqrt{y}} (3x - x^3)^{10} dx dy$$

$$Ans: \frac{2^{12}}{33} \quad (vi) \int_1^e \int_{\ln(y)}^3 (e^x - x)^5 dx dy$$

$$Ans: \frac{e^3 - 4}{6}$$

Chapter 22 | Double Integral in Polar Coordinates




§22.1 When to Use it

We will use Polar coordinates when the region of integration is easier to express in polar coordinates. This might happen mostly two cases:

- The region is interior of a polar curve of the form $r = f(\theta)$.
- The region is circularly symmetric and either the bounds or the integrand has terms involving $x^2 + y^2$.

§22.2 What is dA in Polar Coordinates?

Consider the way a Riemann Sum is constructed on a region with polar coordinates. Subdividing a region of the form $R = \{(r, \theta) \mid a \leq r \leq b, \alpha \leq \theta \leq \beta\}$ into smaller partitions gives us the following pictures:



Accordingly, ΔA is approximated as $r \Delta r \Delta \theta$ and after taking a limit of the Riemann sum we get the following identity

$$\iint_R f \, dA = \iint_R f(r \cos(\theta), r \sin(\theta)) r \, dr \, d\theta$$

Warning: Note that extra r in the second integral. This shows up due to the fact that the area infinitesimal dA does not look like a small square in polar coordinates. A more detailed explanation will be given after we learn about Jacobians and change of variables.

Exploration Activity

When doing integrals in polar coordinates, you often need to integrate trigonometric functions. The double-angle formulas are very useful for this.

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\cos(2\theta) = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta = \cos^2 \theta - \sin^2 \theta$$

These two identities make it easy to integrate $\sin^2 \theta$ and $\cos^2 \theta$.

■ Question 142.



For the following problems

- Sketch the region of integration.
- Try to describe the region in polar coordinates and decide whether you should use polar coordinates or cartesian coordinates.
- Evaluate the integral.

(a) $\iint_R \sqrt{x^2 + y^2} dA$, where R is the region $x^2 + y^2 \leq 1$. Ans: $2\pi/3$

(b) $\iint_R x dA$, where R is the region $x^2 + y^2 \leq 1, x \geq 0$. Ans: $2/3$

(c) $\iint_R (x+y)^2 dA$, where R is the region $1 \leq x^2 + y^2 \leq 9, x \geq 0, y \geq 0$. Ans: $10\pi + 20$

(d) $\iint_R \sqrt{x^2 + y^2} dA$, where R is the region $0 \leq x \leq 1, 0 \leq y \leq 1$ Ans: $2/3$

(e) $\int_{-2}^2 \int_0^{\sqrt{4-x^2}} e^{x^2+y^2} dy dx$ Ans: $\pi \frac{e^4 - 1}{2}$

(f) $\int_{-1}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \sqrt{1-x^2-y^2} dx dy$ Ans: $2\pi/3$

(g) $\int_{\pi/4}^{\pi/2} \int_{1/\sin \theta}^{4/\sin \theta} r dr d\theta$ Ans: 7.5

■ Question 143.



- (a) Find the area of the region R lying between the curves $r = 2 + \sin(3\theta)$ and $r = 4 - \cos(3\theta)$.
- (b) The region R lying inside the curve $r = 2 + \sin 3\theta$ and outside the curve $r = 3 - \sin(3\theta)$ consists of three pieces. Find the area of one of those pieces.
- (c) Find the area of the region which lies inside the circle $x^2 + (y-1)^2 = 1$ but outside the circle $x^2 + y^2 = 1$.

Chapter 23 | Change of Variables in Multiple Integrals



In one-dimensional calculus, we often need to use a change of variable to evaluate integrals. This is known as the technique of u -substitution. The formula goes as follows:

$$\int_a^b f(x) dx = \int_c^d f(g(u))g'(u) du$$

where $x = g(u)$, $a = g(c)$, and $b = g(d)$. Note that we have reversed the roles of x and u , so f is a function of x and x is a function of u . This chapter deals with the question of how to deal with change of variables in double or triple integrals.

Example 23.0.41

We have already seen an example of a change of variables for double integrals: conversion to polar coordinates. The new variables r and θ are related to the old variables x and y by the equations

$$x = x(r, \theta) = r \cos \theta, \quad y = y(r, \theta) = r \sin \theta$$

and the change of variables formula can be written as

$$\iint_R f(x, y) dx dy = \iint_S \underbrace{f(r \cos(\theta), r \sin(\theta))}_{\text{analogue of } f(g(u))} \underbrace{r dr d\theta}_{\substack{\text{analogue of } g'(u)}} dr d\theta$$

where S is the region in the $r\theta$ -plane that corresponds to the region R in the xy -plane.

§23.1 One-to-one Transformation

More generally, we are going to consider a change of variables that is given by a transformation T from the uv -plane to the xy -plane:

$$T(u, v) = (x, y)$$

where

$$x = x(u, v), \quad y = y(u, v)$$

■ Question 144.

□

If we write x and y as functions of new variables u and v so if $x = 2u + 3v$ and $y = u - v$, what are the image of the following points in the $u - v$ in the xy plane?

- | | |
|------------|------------|
| (a) (0, 0) | (c) (3, 1) |
| (b) (3, 0) | (d) (0, 1) |

We usually assume that T is a C^1 transformation, which means that x and y have continuous first-order partial derivatives with respect to u and v . Additionally, we will assume that the function T is a bijection, so that T^{-1} exists.

■ Question 145.

Find the transformation from the xy plane into the uv plane for $u = 2x - 3y$ and $v = y - x$. In other words, find equations for x and y in terms of u and v .

■ Question 146.

A transformation is defined by the equations

$$x = u^2 - v^2, \quad y = 2uv.$$

Find the image of the square $S = \{(u, v) \mid 0 \leq u \leq 1, 0 \leq v \leq 1\}$.

Hint: First find the images of the boundary curves.

§23.2 The Jacobian

Now let's see how a change of variables affects a double integral. The first observation is that image of a rectangle in the uv -plane is not necessarily a rectangle any more. So we can't simply replace $du dv$ with $dx dy$. But we can approximate the area as a cross product of the boundary curves, which can be in turn approximated using tangent lines.

To write down the correct formula, we first need to define a quantity called the Jacobian.

Definition 23.2.42: The Jacobian

The Jacobian of the transformation T given by $x = x(u, v)$ and $y = y(u, v)$ is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

■ Question 147.

Compute the Jacobians for the transformations in the previous section.

Theorem 23.2.43

Suppose that T is a one-to-one C^1 transformation whose Jacobian is nonzero on the interior of a region S in uv -plane and that T maps the region S in the uv -plane onto a region R in the xy -plane. If $f(x, y)$ is continuous on R , then

$$\iint_R f(x, y) dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

So to perform a change of variables in a double integral, we need to complete three steps:

- (a) Substitute for x and y in the integrand in terms of u and v .
- (b) Use the absolute value of the Jacobian to change the dA term.
- (c) Change the bounds of the integrals accordingly.

■ Question 148.

Check that the Jacobian for the change to polar coordinates is equal to r .

□

■ Question 149.

Find the absolute value of the Jacobian for the given change of coordinates.

- (a) $x = 5s + 2t, y = 3s + t$
- (b) $x = s^2 - t^2, y = 2st$
- (c) $x = e^s \cos t, y = e^s \sin t$
- (d) $x = s^3 - 3st^2, y = 3s^2t - t^3$

□

■ Question 150.

Find the region R in the xy -plane corresponding to the region $T = \{(s, t) \mid 0 \leq s \leq 3, 0 \leq t \leq 2\}$ under the change of coordinates $x = 2s - 3t, y = s - 2t$. Check that

$$\int_R dxdy = \int_T \left| \frac{\partial(x, y)}{\partial(s, t)} \right| dsdt.$$

□

■ Question 151.

Use the change of coordinates $s = x + y, t = y$ to find the area of the ellipse $x^2 + 2xy + 2y^2 \leq 1$.

□

23.2.I Harder Problems**■ Question 152.**

Consider $\iint_R y^2 dA$ where R is the region in the upper half-plane bounded by the parabolas $x = y^2, x = y^2 - 9, x = 4 - y^2$, and $x = 16 - y^2$. Use change of variables to convert R into a rectangular region and set up the new integral.

□

■ Question 153.

Use the change of variables $x = u^2 - v^2, y = 2uv$ to evaluate the integral $\iint_R y dA$, where R is the region bounded by the x -axis and the parabolas $y^2 = 4 - 4x$ and $y^2 = 4 + 4x, y \geq 0$.

□

■ Question 154.

Evaluate the integral $\iint_R e^{(x+y)/(x-y)} dA$, where R is the trapezoidal region with vertices $(1, 0), (2, 0), (0, -2)$, and $(0, -1)$.

□

Chapter 24 | Triple Integrals



Just as we defined single integrals for functions of one variable and double integrals for functions of two variables, so we can define triple integrals for functions of three variables.

§24.1 Volume and Mass as an Iterated Integral

Consider first a 3D solid W that looks like a box and is described by the bounds

$$a \leq x \leq b, \quad c \leq y \leq d, \quad p \leq z \leq q$$

Then the triple integral $\iiint_W f(x, y, z) dV$ can be written as an iterated integral

$$\iiint_W f(x, y, z) dV = \int_a^b \int_c^d \int_p^q f(x, y, z) dz dy dx$$


■ Question 155.

What does the integral $\int_0^1 \int_0^1 \int_0^1 2 dz dy dx$ represent?

- (a) Twice the volume of a cube of side 1.
- (b) The volume of a cube of side length 1.
- (c) Twice the volume of a sphere of radius 1.
- (d) The volume under the plane $z = 2$ and over a square of side length 1 in the xy -plane.

Note: In general, the volume of a three-dimensional solid W is given by $\iiint_W dV$. If the density at point (x, y, z) is given by $f(x, y, z)$, then its mass is given by $\iiint_W f(x, y, z) dV$.

Now consider a space region E bounded below by the surface $z = u_1(x, y)$ and above by the surface $z = u_2(x, y)$, and whose 'shadow' (i.e. projection) in the XY-plane is a region D .




Then similar to the double integrals, the mass of the region E can be evaluated as

$$\iiint_E f(x, y, z) dV = \iint_D \left[\int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \right] dA$$

■ Question 156.

□

Now suppose further that the region D looks like either type I or type II region as defined earlier.



In each case, rewrite the triple integral as an iterated integral involving dz , dx , and dy in some order.

Note: Other orders of integration are possible and sometimes necessary to make the integration **feasible**. In general, the rules on the limits on a triple integral are as follows:

- The limits for the outer integral are constants.
- The limits for the middle integral can involve only one variable (that in the outer integral).
- The limits for the inner integral can involve two variables (those on the two outer integrals).

■ Question 157.

□

For each integral (i) - (iv) that makes sense, match it with its region of integration, I or II.

$$(i) \int_1^3 \int_{y-1}^2 \int_0^y f(x, y, z) dz dx dy$$

$$(ii) \int_1^3 \int_0^y \int_2^{y-1} f(x, y, z) dx dy dz$$

$$(iii) \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} f(x, y, z) dz dy dx$$

$$(iv) \int_{-1}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \int_0^{\sqrt{1-x^2-y^2}} f(x, y, z) dz dx dy$$

- I. The region below the plane $z = y$ and above the triangle with vertices $(0, 1), (2, 1), (2, 3)$ in the xy -plane.
II. The region between the upper hemisphere of $x^2 + y^2 + z^2 = 1$ and the xy -plane.

§24.2 Changing the Order of Integration - distinct viewpoints

Example 24.2.44

Consider the triple integral $\iiint_E f(x, y, z) dV$ where E is the solid in the first octant bounded by the surface $z = 2xy$ and the planes $y = x$ and $x = 1$.

- Use CalcPlot3D to add a “region” to your picture.
- Change the 2D Top to the function $y = x$ and the Bottom to $y = 0$ (because we are in the first octant)
- Change the bound for x to $0 \leq x \leq 1$
- Change the 3D Top to $f(x, y) = 2xy$.
- Increase the number of gridline to make your picture smoother.
- Click on the “Format Axes” panel and change it to “Quick View” to *First Octant*.
- Change the coordinate cutoffs until you get a nice picture.

We will start with a z -viewpoint. Rotate the solid until the z -axis is coming out of the screen directly towards you. In this viewpoint:

The ceiling has the equation $z = u_2(x, y) =$ _____

The floor has the equation $z = u_1(x, y) =$ _____

Check that the shadows (projection) of the ceiling and the floor onto the xy -plane match up perfectly. In fact, the projection of the vertical walls down to the xy -plane can be described as a triangle which is in fact both a type I and a type II region.

Written as a type I region, the bounds on the projection are: $\underline{\quad} \leq y \leq \underline{\quad}$, $\underline{\quad} \leq x \leq \underline{\quad}$.


Written as a type II region, the bounds on the projection are: $\underline{\quad} \leq x \leq \underline{\quad}$, $\underline{\quad} \leq y \leq \underline{\quad}$.

So the triple integral can be written as an iterated integral of the form

$f(x, y, z) dz dy dx$, and _____

$f(x, y, z) dz dx dy$, _____

Go back to CalcPlot3D and this time rotate the solid until the y -axis is coming out of the screen directly towards you.



In this viewpoint:

The ceiling has the equation $y = u_2(x, z) =$ _____

The floor has the equation $y = u_1(x, z) =$ _____

Once again, check that “shadows” (i.e. projection) onto the xz -plane of the ceiling and the floor in this viewpoint match each other perfectly. However, the projection of the walls onto the xz -plane does not give a triangle this time.

Clearly two of the sides of the projection are straight lines. The third side is the intersection of the surfaces $x = y$ and $z = 2xy$. Then the equation of the projection of this curve onto the xz -plane is

[Hint: the equation only involves x and z .]

Do the same steps as above to rewrite the triple integral as an iterated integral of the form

$f(x, y, z) dy dx dz$, and _____

$f(x, y, z) dy dz dx$.

Finally, we claim that the x -viewpoint is not feasible and there is no way to write the triple integral as a single iterated integral that starts with dx . Can you see why?

■ Question 158.

□

Express the iterated integral

$$\int_0^1 \int_0^{x^2} \int_0^y f(x, y, z) dz dy dx$$

as a triple integral and then rewrite it as an iterated integral in the following orders.


- (a) Integrate first with respect to x , then z , and then y .
- (b) Integrate first with respect to y , then x , and then z .

■ Question 159.

□

Consider the region T in the first octant bounded by the plane $x + y = 2$ and the parabolic cylinder $z = 4 - x^2$. Write the volume of this solid as an iterated integral involving

- (i) $dz dy dx$
- (ii) $dy dz dx$
- (iii) $dy dx dz$



■ Question 160.


□

Consider the region T in the first octant bounded by the elliptical cylinder $4x^2 + y^2 = 4$, the plane $2x + z = 2$ and the plane $y = 2$. For the given order of integration, write an iterated integral equivalent to the triple integral

$$\iiint_T 1 dV.$$

- (i) $dx dy dz$
- (ii) $dy dx dz$
- (iii) $dz dx dy$

Here are two pictures of the region T from two different viewpoints.



■ **Question 161.**

Consider the solid B bounded by $y = x^2 - 2, x + y = 0, z = -2, x + 2z = 2$. Which of the viewpoints are feasible?

□

■ **Question 162.**

Consider the solid B bounded by the parabolic cylinder $z = 6y^2$ and the elliptical cylinder $x^2 + 3y^2 = 12$ in the first octant. Draw the region using CalcPlot3D. Which of the viewpoints are feasible? For each of the following, write the volume of the solid B as an iterated integral if the viewpoint is feasible:

(i) $\mathrm{d}z \, \mathrm{d}x \, \mathrm{d}y$ (ii) $\mathrm{d}x \, \mathrm{d}z \, \mathrm{d}y$ (iii) $\mathrm{d}y \, \mathrm{d}z \, \mathrm{d}x$

□

■ **Question 163.**

Find a solid W such that the integral

$$I = \iiint_W (x^2 - x) \, dV$$

(i) is positive.

(ii) is negative

(iii) is zero.

□


§24.3 Practice Problems with Polar Coordinates

The following exercises require you to set up the base of integration in polar coordinates. The coordinate system (r, θ, z) is called a cylindrical coordinate system. Don't forget the extra r when setting up polar integrals.

■ **Question 164.**

A spherical ball of radius 5 cm is sliced off by a plane that leaves the following solid. Write an iterated triple integral in $\mathrm{d}z \, \mathrm{d}r \, \mathrm{d}\theta$ which represents the volume of this region.


□

**■ Question 165.**

□

A bead is made by drilling a cylindrical hole of radius 1 mm through a sphere of radius 5 mm.

- Set up a triple integral in $dzdrd\theta$ that represents the volume of the bead.
- Evaluate the integral.



Part V

Vector Calculus

Chapter 25 | Vector Fields and Flow Lines



§25.1 Motivation

Given a function $f(x, y)$ of two variables, consider what the gradient function $\vec{\nabla}f(x, y)$ represents. To each point in the plane it associates a vector that represents the magnitude and direction greatest increase of $f(x, y)$ at the point. We have learned earlier that vector $\vec{\nabla}f(x, y)$ is always perpendicular to the contour plot of f through the point (x, y) . Using this knowledge we can draw all these vectors in XY-plane as in the following picture:




Figure 25.1: Contour Plot and Gradients of $f(x, y) = x^2 - y^2$

Thus $\vec{\nabla}f$ is a function that takes vector inputs (position vector of a point) and gives a vector output. In general, functions of this kind are given a special name: they are called **Vector Fields**.

Definition 25.1.45

A vector field in 2-space is a function

$$\vec{F}(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j}$$

whose value at a point (x, y) is a 2-dimensional vector. Similarly, a vector field in 3-space is a function $\vec{F}(x, y, z)$ whose values are 3-dimensional vectors.

There are many other examples of vector fields that we come across in real life frequently e.g. Ocean currents, wind flow in your weather app, electromagnetic force field, gravitational force etc. It is not necessary that a vector field \vec{F} comes from gradient of a function.

§25.2 Examples of Vector Fields


Example 25.2.46

Consider the vector field $\vec{F}(x, y) = x\hat{i} + y\hat{j}$. We are interested in what it looks like.

- Along the x -axis, we have $y = 0$, so the arrows are purely horizontal.

- Along the y -axis, we have $x = 0$, so the arrows are purely vertical.
- What about the arrow at other points? According to the formula, the vector at the point (x, y) is equal to its position vector. So the arrow points radially outward and has length equal to the distance of the point from the origin.

With these observations, our vector field looks like:



■ Question 166.

Plot the following vector fields:

- $\vec{F}(x, y) = y\hat{i}$
- $\vec{F}(x, y) = (x + y)\hat{i} + (x - y)\hat{j}$

■ Question 167.

Give examples of vector fields where:

- all output vectors are parallel to the x -axis, and have the same magnitude if they are located on the same vertical line.
- all output vectors are unit length, and perpendicular to the position vectors $x\hat{i} + y\hat{j}$.

■ Question 168.

Match the following vector fields with their drawings in fig. 25.2.

- | | | | |
|-------------------------------|-------------------------------|-------------------------------|---------------------------------|
| (a) $\langle y, 1 \rangle$ | (d) $\langle -2y, 3x \rangle$ | (g) $\langle x^2y, 0 \rangle$ | (j) $\langle -y - x, x \rangle$ |
| (b) $\langle 0, 2y \rangle$ | (e) $\langle 0, x^2y \rangle$ | (h) $\langle -x, 0 \rangle$ | (k) $\langle -y, x \rangle$ |
| (c) $\langle -x, -2y \rangle$ | (f) $\langle -2y, -x \rangle$ | (i) $\langle -2y, 1 \rangle$ | (l) $\langle x^2, y^2 \rangle$ |




Figure 25.2

Chapter 26 | Differential Operators



In this section we define two operations that can be performed on vector fields and that play a basic role in the applications of vector calculus to fluid flow and electricity and magnetism. Each operation resembles differentiation, but one produces a vector field whereas the other produces a scalar field.

Consider the mathematical abstract ‘construct’

$$\vec{\nabla} = \frac{\partial}{\partial x} \hat{i} + \frac{\partial}{\partial y} \hat{j} + \frac{\partial}{\partial z} \hat{k}.$$

It looks like a vector, but clearly the components are neither functions, nor scalars. This object is called a “differential operator”. It’s an operation you can apply to a function or a Field. We will use it to define two operations on a vector field: curl and \div .

§26.1 Curl

Definition 26.1.47

Given a vector field $\vec{F} = P \hat{i} + Q \hat{j} + R \hat{k}$, we define the **curl** of \vec{F} to be the **vector field**

$$\text{curl } \vec{F} = \vec{\nabla} \times \vec{F}.$$

You should check that in the case of 2-dimensional vector fields, above definition gives the correct familiar formula for $\nabla \vec{F}$ and $\text{curl } \vec{F}$.

Warning: Since we have defined the differential operator $\vec{\nabla}$ which looks like a vector, we might naturally ask what is the dot product of the operator and the vector field. Note that the dot product $\vec{\nabla} \cdot \vec{F}$ is **not** the gradient because gradient of a vector field doesn’t make sense. The gradient of a function $g(x, y, z)$ can be written as $\vec{\nabla}g$ without any ‘dot’ (since it’s not a dot product). We will see how to interpret the dot product in the next section.

■ Question 169.

Write out the formula for $\text{curl } \vec{F}$ for a 3D vector field using $\hat{i}, \hat{j}, \hat{k}$ notation.

■ Question 170.

Now suppose \vec{F} is a gradient vector field. I.e.,

$$\vec{F} = P \hat{i} + Q \hat{j} + R \hat{k} = g_x \hat{i} + g_y \hat{j} + g_z \hat{k}$$

for some function $g(x, y, z)$, and suppose g has continuous second-order partial derivatives. Use Clairaut’s theorem to prove that $\text{curl } \vec{F} = 0$.

Note: This statement is sometimes written as $\vec{\nabla} \times (\vec{\nabla}g) = 0$.

Definition 26.1.48

A vector field is called **irrotational** if its curl is zero at every point where curl is defined.

■ Question 171.

Show that

$$\vec{F}(x, y, z) = \langle y^2 z^3, 2xyz^3, 3xy^2 z^2 \rangle$$

is an irrotational vector field.

26.I.I Understanding Curl in 2D

To understand the physical interpretation of curl, let's take a look at the two-dimensional version first. Given a planar vector field $\vec{F} = P(x, y)\hat{i} + Q(x, y)\hat{j}$, the curl of the vector field should be


$$\text{curl}(\vec{F}) = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \hat{k}$$

Of course, this is only defined at points where P and Q are differentiable with respect to y and x respectively. But what does the quantity inside the parentheses mean geometrically?

Curl and Rotation

The nomenclature suggests that $\text{curl}(\vec{F})$ has something to do with rotation in a vector field. There is a way to visually interpret the algebraic quantity that shows how curl corresponds to rotation.


Imagine an infinitesimally small paddle-wheel lying in a vector field $\vec{F}(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j}$. We will show that as long as \vec{F} is smooth, the curl measures how much (and which way) it turns at any point.




A positive curl means the paddle-wheel will rotate counterclockwise, and a negative curl means the paddle-wheel rotates clockwise. An irrotational vector field has no spinning paddle-wheels anywhere.

In general, the first term $\frac{\partial Q}{\partial x}$ in the curl function measures the effect of the vector field \vec{F} on the horizontal part of the paddle wheel, and a positive value corresponds to a counter-clockwise contribution to the rotation of the paddle-wheel. This follows because Q is the j -component of the vector field, and its partial derivative with respect to x measures its change when moving in horizontally, keeping y fixed.


The picture below shows a situation where $\frac{\partial Q}{\partial x}$ is **positive** since the (red) vectors representing the j -component $Q(x, y)$ of the vector field increases as we move horizontally. In this case, the paddle-wheel rotates counterclockwise.



Likewise, the second term $\frac{\partial P}{\partial y}$ in the curl function measures the effect of the vector field on the vertical part of the paddle wheel, but now a negative value corresponds to a counterclockwise contribution to the rotation of the paddle-wheel (that's why this term appears in the curl with a negative sign).



Note: In general, consider a three dimensional velocity vector field \vec{F} for fluid flow. Then $\text{curl } \vec{F}(x, y, z)$ is a vector at (x, y, z) that describes the rotation of particles near (x, y, z) in the following sense: if we picture a small paddlewheel in the fluid at (x, y, z) , the wheel rotates fastest when its axis is parallel to $\text{curl } \vec{F}(x, y, z)$. If $\text{curl } \vec{F}(x, y, z) = \vec{0}$, the fluid is free from rotations at the point and \vec{F} is irrotational at the point. In this case, a tiny paddle wheel moves with the fluid but doesn't rotate about its axis. If $\text{curl}(\vec{F}) \neq \vec{0}$, the paddle wheel rotates about its axis.



§26.2 Divergence

If the curl can be interpreted of as a **vector derivative** of the vector field, we define the **scalar derivative** of the vector field as follows.

Definition 26.2.49

If $\frac{\partial P}{\partial x}$, $\frac{\partial Q}{\partial y}$, and $\frac{\partial R}{\partial z}$ exist, then the **divergence** of \vec{F} is defined to be the **scalar** quantity

$$\operatorname{div} \vec{F} = \vec{\nabla} \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

■ Question 172.

Show that If $\vec{F} = P\hat{i} + Q\hat{j} + R\hat{k}$ is a vector field defined on \mathbb{R}^3 and P, Q, and R have continuous second-order partial derivatives, then

$$\operatorname{div} \operatorname{curl} \vec{F} = 0.$$

This is also sometimes written as

$$\vec{\nabla} \cdot (\vec{\nabla} \times \vec{F}) = 0.$$

■ Question 173.

Show that the vector field

$$\vec{F}(x, y, z) = xz\hat{i} + xyz\hat{j} - y^2\hat{k}$$

cannot be written as the curl of another vector field, that is, $\vec{F} \neq \operatorname{curl} \vec{G}$.

26.2.1 Understanding Divergence

The reason for the name divergence can be understood in the context of fluid flow. If $\vec{F} = P\hat{i} + Q\hat{j} + R\hat{k}$ the velocity of a fluid (or gas), then $\operatorname{div} \vec{F}$ represents the net rate of change (with respect to time) of the mass of fluid (or gas) flowing from the point (x, y, z) per unit volume. In other words, $\operatorname{div} \vec{F}$ measures the tendency of the fluid to diverge from the point (x, y, z) . if $\operatorname{div} \vec{F} = 0$ then the vector field \vec{F} is said to be **incompressible**.

Exploration Activity

Laplace Operator:

For the sake of completion we also mention another differential operator that occurs when we compute the divergence of a gradient vector field.

$$\operatorname{div}(\vec{\nabla} f) = \vec{\nabla} \cdot (\vec{\nabla} f)$$

is abbreviated as $\nabla^2 f$, and the operator ∇^2 is called the **Laplace operator**. $\nabla^2 f$ is also denoted as Δf .

You can easily check that

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.$$

Chapter 27 | Line Integrals on Parameterized Curves




§27.1 Motivation


Now that we know about parameterized curves and vector fields, we are ready to introduce a new kind of integral which uses both of these concepts. The line integral measures the extent to which a curve in a vector field is, overall, going with the vector field or against it. For instance, a flow line goes completely with the flow of its vector field, so it should have a comparatively high line integral value.

§27.2 Definition

Suppose we are given a vector field $\vec{F}(x, y)$ and a parametrized curve $C : \vec{r}(t), a \leq t \leq b$. First we will need to fix an **orientation** of C so that we know the direction of travel along the curve.



We will assume that the orientation is as in the first picture. Next, recall that we can use dot product to measure the extent to which two vectors point in the same or opposing directions. So we will break our parameterized curve C from $\vec{r}(a) = P$ to $\vec{r}(b) = Q$ into a linked trail of “secant” vectors $\Delta\vec{r}_i$, connecting points along the curve, and then compare each of these vectors to the corresponding vector from the vector field vector at the same points.



We then add the dot products and create a Riemann sum whose limit gives us the Line Integral.

$$\int_C \vec{F} \cdot d\vec{r} = \lim_{\|\Delta\vec{r}_i\| \rightarrow 0} \sum_{i=0}^{n-1} \vec{F}(\vec{r}_i) \cdot \Delta\vec{r}_i$$


Note that the value of the line integral also defines the work done by the force \vec{F} as displacement happens along the curve C , which is an important quantity in physics.

■ Question 174.

For **each curve C_i in the picture and for each of the following vector fields, determine if possible whether $\int_C \vec{F} \cdot d\vec{r}$ would be positive, negative, or zero.**

$$(i) \quad x\hat{i}$$

$$(ii) \quad y\hat{i} - x\hat{j}$$



27.2.1 Circulation

Definition 27.2.50


A curve C is called **closed** (or a **closed loop**) if it starts and ends at the same point.

Line integral on a closed loop is sometimes explicitly denoted by the symbol \oint_C , although it is not necessary. Recall that $d\vec{r} = \vec{T} ds$, and $\vec{F} \cdot \vec{T}$ is the component of \vec{F} in the direction of the unit tangent vector \vec{T} . This means that the closer the direction of \vec{F} is to the direction of \vec{T} , the larger the value of $\vec{F} \cdot \vec{T}$. Thus $\oint_C \vec{F} \cdot d\vec{r}$ is a measure of the tendency of the vector field to move around C in the same direction as the orientation of C , and is called the **circulation** of \vec{F} around C .

§27.3 Evaluating a line integral

The key to evaluating line integration is to relate the secant vectors $\Delta \vec{r}_i$ to the velocity vectors $\vec{r}'(t_i)$ as

$$\Delta \vec{r}_i \approx \vec{r}'(t_i) \Delta t$$



As $\Delta t \rightarrow 0$, we then find the limit of the Riemann sum as

$$\int_C \vec{F} \cdot d\vec{r} = \lim_{\Delta t \rightarrow 0} \sum_{a \leq t \leq b} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) \Delta t = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$$

Observe that this is essentially a u -substitution, where we are changing the variable of integration to t .

Example 27.3.51

Let's compute the line integral

$$\int_C \langle x - y, x \rangle \cdot d\vec{r}$$

along the upper-half C of the unit circle traversed counterclockwise. First we need to parameterize C and find $\vec{r}'(t)$.

$$\vec{r}(t) = \cos t \hat{i} + \sin t \hat{j}, \quad 0 \leq t \leq \pi$$

So $\vec{r}'(t) = \langle -\sin t, \cos t \rangle$. Then using the formula for line integrals,

$$\begin{aligned} \int_C \langle x - y, x \rangle \cdot d\vec{r} &= \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt \\ &= \int_0^\pi \underbrace{\langle \cos(t) - \sin(t), \cos(t) \rangle}_{\vec{F}(\vec{r}(t))} \cdot \underbrace{\langle -\sin(t), \cos(t) \rangle}_{\vec{r}'(t)} dt \\ &= \int_0^\pi (-\cos(t)\sin(t) + \sin^2(t) + \cos^2(t)) dt \\ &= \int_0^\pi -\cos(t)\sin(t) dt + \int_0^\pi 1 dt \\ &= \left(\frac{\cos^2(t)}{2} + t \right) \Big|_0^\pi = \pi \end{aligned}$$

■ Question 175.

Compute the line integral $\int_C \vec{F} \cdot d\vec{r}$ for the following pairs of vector fields \vec{F} and curves C . □

- (a) $\vec{F} = \langle y, x \rangle$ and C is the quarter-circle centered at the origin starting at $(2, 0)$ and proceeding counterclockwise to $(0, 2)$.
- (b) $\vec{F} = \langle y, x \rangle$ and C is the line segment starting at $(2, 0)$ and proceeding counterclockwise to $(0, 2)$
- (c) $\vec{F} = \langle x, y, -2z \rangle$ and C is the “twisted cubic” $\vec{r}(t) = \langle t, t^2, t^3 \rangle$ from $t = 0$ to $t = 1$
- (d) $\vec{F} = \langle x, y, -2z \rangle$ and C is the line segment from $(0, 0, 0)$ to $(1, 1, 1)$

■ Question 176.

True or False? The line integral of a vector field along a path depends on the parameterization of the curve. □

Chapter 28 | The Fundamental theorem of Line Integrals



Recall the Fundamental Theorem of (single-variable) Calculus:

$$\int_a^b f'(x)dx = f(b) - f(a)$$

This essentially says that integration is the opposite operation of differentiation (that's the reason we call the indefinite integral the anti-derivative). In this chapter we'll see a multivariable version of the Fundamental Theorem of Calculus, which uses the line integral of a (continuous) gradient $\vec{F} = \nabla f$. Among other things, this result will allow us to compute some line integrals quickly and without resorting the approach from last time involving parameterization.

§28.1 Gradient Vector Field

A vector field $\vec{F}(x, y)$ is called a gradient vector field if it can be written as the gradient of some underlying function $f(x, y)$ i.e.

$$\vec{F} = \vec{\nabla}f = \langle f_x, f_y \rangle$$

In that case, the function f is called the potential function for the field \vec{F} (the terminology comes from Physics and has to do with potential energy of objects).

■ Question 177.


Match the following functions with their gradient vector fields.

(a) $x^2 + y^2$

(c) $(x + y)^2$

(b) $x(x + y)$

(d) $\sin \sqrt{x^2 + y^2}$



Gradient vector fields have some very nice properties (which we'll explore soon), so it will be important for us to identify when a given vector field is a gradient vector field. Recall that Clairaut's theorem says $f_{xy} = f_{yx}$ for a smooth function. That means for a gradient vector field $\vec{F} = \nabla f$, we must have $P_y = Q_x$. As a contrapositive, we have the following test.

Theorem 28.1.52

If $\frac{\partial P}{\partial y} \neq \frac{\partial Q}{\partial x}$, then \vec{F} is not a gradient vector field.

A **Warning:** The natural follow-up question to ask is whether the equality of the two partials then guarantees a gradient vector field. Unfortunately, we cannot yet answer this question with the material covered so far. We will show that the answer is affirmative in a couple of chapters.

28.1.1 Finding the potential function

Assume that you are told that a given vector field $\vec{F}(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j}$ is in fact a gradient vector field. How do we then find the exact formula of a function $f(x, y)$ such that $\vec{F} = \nabla f$?

Here is an algorithm. Note that we want to find f such that

$$f_x = P \text{ and } f_y = Q$$

Integrating the first equation with respect to x gives us

$$f(x, y) = \int P(x, y)dx + g(y)$$

The extra $g(y)$ serves as a replacement for the arbitrary constant term ' $+c$ ' because $\frac{\partial g(y)}{\partial x} = 0$. With this presumptive model for $f(x, y)$ we must have

$$\frac{\partial}{\partial y} \left(\int P(x, y) dx + g(y) \right) = Q$$

From here you can solve for g . We can also start from Q if that seems easier.

It's best to learn the algorithm by doing examples:

Example 28.1.53

Suppose $\vec{F}(x, y) = (x+y)\hat{i} + (x-y)\hat{j} = \nabla f(x, y)$. Then $f_x = x+y$ and $f_y = x-y$. The first equation tells us

$$f(x, y) = \frac{x^2}{2} + xy + g(y)$$

Then

$$x-y = \frac{\partial}{\partial y} \left(\frac{x^2}{2} + xy + g(y) \right) = 0 + x + g'(y) \implies g'(y) = -y \implies g(y) = -\frac{y^2}{2} + c$$

So overall, $f(x, y) = \frac{x^2}{2} + xy - \frac{y^2}{2} + c$.

■ Question 178.

For each of the following vector fields, find a potential function or explain why one doesn't exist.

$$(a) xi$$

$$(xye^{xyz} + 2xz \cos(xz^2))\hat{k}$$

$$(b) yi$$

$$(g) \frac{-y\hat{i} + x\hat{j}}{x^2 + y^2}$$

$$(c) (x^2 - y^2)\hat{i} - 2xy\hat{j}$$

$$(h) \frac{\vec{r}}{\|\vec{r}\|^3} = \frac{x}{(x^2 + y^2 + z^2)^{3/2}}\hat{i} + \frac{y}{(x^2 + y^2 + z^2)^{3/2}}\hat{j} + \frac{z}{(x^2 + y^2 + z^2)^{3/2}}\hat{k}$$

$$(d) (2xy + 5)\hat{i} + (x^2 + 8y^3)\hat{j}$$

$$(e) 2xe^{x^2} \sin y\hat{i} + e^{x^2} \cos y\hat{j}$$

$$(f) (yze^{xyz} + z^2 \cos(xz^2))\hat{i} + xze^{xyz}\hat{j} +$$

§28.2 The Fundamental theorem of Line Integrals

28.2.1 Formulation

The single-variable FTC says that, to evaluate the integral of a function f' over an interval $[a, b]$, we find its antiderivative f , plug in the endpoints and subtract. By analogy, think of the potential function $f(x, y)$ as being an “antiderivative” for the gradient field $\vec{F} = \nabla f$. And let C be a piece-wise smooth oriented path with starting point A and endpoint B. Then we have the following theorem

Theorem 28.2.54: The Fundamental Theorem for Line Integral

$$\int_C \vec{\nabla} f \cdot d\vec{r} = f(B) - f(A)$$

Before we prove this to be true, note how it simplifies the computation of line integrals for gradient fields! Instead of parametrizing C , we just have to identify its endpoints, find the potential function, and plug in the endpoints and subtract.

Proof of theorem 54.


Suppose C is parametrized by $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j}$, $a \leq t \leq b$. Recall that from last chapter, we have

$$\begin{aligned}\int_C \vec{\nabla}f \cdot d\vec{r} &= \int_a^b \vec{\nabla}f \cdot \vec{r}'(t) dt = \int_a^b \langle f_x, f_y \rangle \cdot \langle x'(t), y'(t) \rangle dt \\ &= \int_a^b \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} \right) dt \\ &= \int_a^b \frac{df}{dt} dt \\ &= \int_A^B df = f(B) - f(A)\end{aligned}$$

■

■ Question 179.

Consider the line integrals, $\int_{C_i} \vec{F} \cdot d\vec{r}$ for $i = 1, 2, 3, 4$, where $\vec{F}(x, y) = \vec{\nabla}f$ and C_i is the path from P_i to Q_i shown below. Some level curves of f are also shown in the figure.




- (a) Which of the line integral(s) is (are) zero?
- (b) Arrange the four line integrals in ascending order (from least to greatest).
- (c) Two of the nonzero line integrals have equal and opposite values. Which are they? Which is negative and which is positive?

§28.3 Path Independence

Definition 28.3.55

A vector field is called **path-independent** or **conservative** if the value of its line integral is the same along any paths having the same endpoints.



In other words, the line integrals depend only on the starting and ending points and NOT on the particular path between those two points. Here ‘path’ means a piecewise smooth curves C that avoid any locations where \vec{F} is undefined.

How can we tell whether a given vector field is path-independent or path-dependent, without computing a bunch of line integrals to check? Note that with this new nomenclature, the FTI can be stated as

Gradient Vector Field \implies Path independent


In fact, the converse is true as well. We are going to include the proof below as a digression, but it is not within the scope of the current discussion.

Exploration Activity →


Suppose $\vec{F} = P\hat{i} + Q\hat{j}$ is a path-independent vector field. Consider the function f defined as follows:

$$f(x, y) = \int_C \vec{F} \cdot d\vec{r}$$

where C is a path from a fixed starting point O to a variable point $X = (x, y)$. This integral has the same value for any path from O to X because \vec{F} is path-independent. The two figures below then explain why $\nabla f = \vec{F}$.



The path $C' + L_y + L_x$ is used to show $f_x = P$



The path $C' + K_x + K_y$ is used to show $f_y = Q$

So we can conclude that

Gradient Vector Field \iff Path independent

So it is a natural question to ask if a given vector field is a gradient vector field or not since that would make our life considerably easier! And if it is a gradient vector field, how can find f so that $\vec{F} = \vec{\nabla}f$? Fortunately, we have already answered these questions in a previous chapter.




Warning: Note that we still haven't proved that a vector field is gradient v.f. if $Q_x = P_y$, although it is true in most cases.

■ Question 180.

Evaluate the following line integrals. Not all of them can be done using FTLL. Identify the ones that are gradient vector fields.

□

- (a) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y) = xy \vec{i} + x^2 \vec{j}$ and C is given by $\vec{r}(t) = \sin t \vec{i} + (1+t) \vec{j}$, $0 \leq t \leq \pi$
- (b) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y) = (4x^3y^2 - 2xy^3) \vec{i} + (2x^4y - 3x^2y^2 + 4y^3) \vec{j}$ and C is given by $\vec{r}(t) = (t + \sin \pi t) \vec{i} + (2t + \cos \pi t) \vec{j}$, $0 \leq t \leq 1$
- (c) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y,z) = \sin y \vec{i} + x \cos y \vec{j} - \sin z \vec{k}$, and C is the helix $x = 3 \cos t$, $y = t$, $z = 3 \sin t$ from $(3,0,0)$ to $(0,\pi/2,3)$
- (d) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y,z) = (3x^2yz - 3y) \vec{i} + (x^3z - 3x) \vec{j} + (x^3y + 2z) \vec{k}$ and C is the curve shown below.




- (e) $\int \vec{F} \cdot d\vec{r}$, where
 $\vec{F}(x,y,z) = \langle 4xe^{2x^2+3y^2+4z^2}, 6ye^{2x^2+3y^2+4z^2}, 8ze^{2x^2+3y^2+4z^2} \rangle$

and C is

$$\vec{r}(t) = \langle (2 + \cos(7t)) \cos(t), (2 + \cos(7t)) \sin(t), \sin(7t) \rangle$$

parameterized by $0 \leq t \leq \pi$ starting at $t = 0$ and ending at $t = \pi$.

- (f) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y) = xy^2 \vec{i} + x^2y \vec{j}$, and C is $\vec{r}(t) = \cos t \vec{i} + 2 \sin t \vec{j}$, $0 \leq t \leq \pi/2$
- (g) $\int \vec{F} \cdot d\vec{r}$, where $\vec{F}(x,y) = (\sin(\frac{x}{2}) \sin(\frac{y}{2})) \hat{i} - (\cos(\frac{x}{2}) \cos(\frac{y}{2})) \hat{j}$, and C is as in the picture below:



§28.4 Circulation Freedom

Definition 28.4.56

A vector field \vec{F} is called **circulation-free** if $\oint_C \vec{F} \cdot d\vec{r} = 0$ for **every** closed path C.

Again, here ‘path’ means a piecewise smooth curve C that avoids any locations where \vec{F} is undefined.

One of the immediate observations we could make from the definition is that for a vector field

Path independent \implies Circulation-free

The converse is non-trivial but still easy to prove.

■ Question 181.

Explain why circulation-free vector fields are path-independent.

§28.5 Conclusion

Overall, the results of this chapter show the following

Gradient v.f. \iff Path independent \iff Circulation-free

Chapter 29 | Parametric Surface


Chapter 30 | Irrotational Vector Fields



§30.1 Green's Idea: Volume Integrate the Curl

The fact that gradient vector fields are both circulation free and irrotational lead a famous mathematical physicist named George Green to guess that there might be a relationship between circulation integrals to volume integrals of the curl.

To approximate the volume integral of the curl function over a closed bounded region R , we can first break R into smaller rectangular sub-rectangles and then add up the volumes of the rectangular solids with heights equal to the curl at one of the corners of the sub-rectangles.



The key thing to notice here is that the vector field spins “adjacent” paddle-wheels in opposite directions: The above red vector gives counterclockwise spin to the left paddle-wheel and the same amount of clockwise spin to the right paddle-wheel. This means that the curls associated with interior points of the region R will essentially be cancelled out by each other. So, when we take the limit of the volume approximation, all that is left is the spin due to the (counterclockwise) push of the vector field along the boundary of the region. This is measured by the line integral of the vector field around the (counterclockwise oriented) boundary.

This is of course, very much of a heuristic reasoning. An exact algebraic proof can be found in the book if you are interested.

30.1.1 Careful Formulation of Green's Theorem

Green's Theorem is a statement about only the 2D plane, not 3D space. Here's the setup: Let C be a piecewise smooth simple closed curve that is the boundary of a simply-connected region R in the plane. Let $\vec{F} = P \vec{i} + Q \vec{j}$ be a smooth vector field defined on all of R and C .

Definition 30.1.57

To say a curve is **simple** means that it doesn't intersect itself, and to say a curve is **closed** means that it is a closed loop, that it starts and ends at the same point.

Definition 30.1.58

A region is called **simply-connected** if it is just one piece (connected) and doesn't have any holes in it.

Definition 30.1.59

A vector field is said to be **smooth** if its first partials are continuous functions.

Since C is the boundary of R , we will denote it as $C = \partial R$ and stop referring to C explicitly. When we consider the boundary curve of a simply-connected region, we always orient the curve so that the region is on the left as we follow the curve.

Theorem 30.1.60: Green's Theorem

Green's theorem relates the line integral of \vec{F} over ∂R to the volume integral of $\text{curl}(F)$ over R .

$$\oint_{\partial R} \vec{F} \cdot d\vec{r} = \iint_D \text{curl}(F) dA = \iint_R (Q_x - P_y) dA.$$

You may also see this written in the Leibniz notation for partial derivatives.

$$\oint_{\partial R} P dx + Q dy = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

■ Question 182.

□

Use Green's theorem to explain why smooth irrotational vector fields are guaranteed to be circulation free, and hence gradient vector fields.

§30.2 The Venn Diagram

Consider the following particularly illuminating example.


Example 30.2.61

We start with the vector field

$$\vec{F}(x, y) = \frac{y}{x^2 + y^2} \hat{i} - \frac{x}{x^2 + y^2} \hat{j}$$

- (a) Check that the vector field is irrotational.
- (b) Use a parameterization to compute $\oint_C \vec{F} \cdot d\vec{r}$ where C is the unit circle traversed once in a counterclockwise direction.
- (c) Explain how you can conclude that \vec{F} is not a gradient vector field.
- (d) Let $f(x, y) = \arctan \frac{y}{x}$. Calculate $\vec{\nabla} f$.
- (e) How do you explain the apparent contradiction?

This example highlights the requirement of smoothness in Green's theorem. In particular it shows that not all irrotational vector fields are gradient vector fields. Overall, the venn diagram of vector fields looks as follows:



Chapter 31 | Green's Theorem - Applications and Generalizations



§31.1 Applications of Green's Theorem

31.1.1 Evaluating Difficult Line Integrals

Example 31.1.62

If you're asked to find the line integral of an ugly vector field over a closed curve, you should look to see if Q_x and P_y are drastically more simple. If so, use Green's Theorem!

■ Question 183.

Evaluate

$$\oint_C (2y + \sqrt{9+x^3}) dx + (5x + e^{\tan^{-1} y}) dy,$$

where C is the circle $x^2 + y^2 = 4$ in the plane oriented counterclockwise.

Example 31.1.63

If you're asked to find the line integral over a closed curve that is clearly the boundary of a nice region, it's often a good idea to use Green's Theorem to switch to the double integral over the interior.

■ Question 184.

Integrate $\vec{F} = xy \vec{i} + e^x \vec{j}$ over the boundary of the rectangle determined by $0 \leq x \leq 2$, $0 \leq y \leq 3$, oriented clockwise around the boundary.

■ Question 185.

Find the line integral of $\vec{F} = 3xy \vec{i} + 2x^2 \vec{j}$ over the curve C defined as follows: follow the curve $y = x^2 - 2x$ from $(0,0)$ to $(3,3)$, then follow the line $y = x$ from $(3,3)$ back to $(0,0)$.

■ Question 186.

Evaluate

$$\oint_C (3y - e^{\sin x}) dx + (7x + \sqrt{y^4 + 1}) dy$$

where C is the circle $x^2 + y^2 = 9$, oriented clockwise.

Example 31.1.64

Consider the following problem.

Find the line integral of

$$\vec{G}(x,y) = (x+y) \vec{i} + \left(2x+y \ln(\csc \sqrt{1-y^5})\right) \vec{j}$$

over C_1 , the upper half of the unit circle from $(1, 0)$ to $(-1, 0)$.

The problem is that the vector field is ugly, so parametrizing C_1 is just going to lead to an impossible integral. So we would like to use Green's Theorem, but this isn't a closed curve!

Here's how to fix that issue. Let C_2 be the straight line segment from $(-1, 0)$ to $(1, 0)$. Now $C_1 + C_2$ is a closed loop.

(a) Let R be the region enclosed by $C_1 + C_2$. Use Green's Theorem to compute $\oint_{C_1+C_2} \vec{G} \cdot d\vec{r}$.

(b) Parametrize C_2 and directly calculate $\int_{C_2} \vec{G} \cdot d\vec{r}$. (Note that $y = 0$ everywhere on C_2 , which is helpful.)

(c) Write $\oint_{C_1+C_2} \vec{G} \cdot d\vec{r} = \int_{C_1} \vec{G} \cdot d\vec{r} + \int_{C_2} \vec{G} \cdot d\vec{r}$ and use your answers to part (a) and (b) to finish off the problem and find the line integral of \vec{G} along C_1 .

■ Question 187.

Evaluate the integral.

□

■ Question 188.

□

Compute the line integral of the vector field

$$\vec{F}(x, y, z) = \langle \cos(x), 2 + \cos(y), e^z + x(y^2 + z^2) \rangle$$

along the curve

$$\vec{r}(t) = \langle t, \cos(t), \sin(t) \rangle \text{ with } 0 \leq t \leq 3\pi.$$

[HINT: Write \vec{F} as $\vec{G} + \vec{H}$ where \vec{G} is a gradient vector field. Then do the two integrals separately.]

3I.I.2 Calculating Area

Consider the following vector fields:

$$\vec{F}_1 = x \vec{j}, \quad \vec{F}_2 = -y \vec{i}, \quad \vec{F}_3 = -\frac{1}{2}y \vec{i} + \frac{1}{2}x \vec{j}.$$

What is $Q_x - P_y$ for each of these fields? Applying Green's Theorem to a region D , we get that

$$\oint_{\partial D} \vec{F}_1 \cdot d\vec{r} = \oint_{\partial D} \vec{F}_2 \cdot d\vec{r} = \oint_{\partial D} \vec{F}_3 \cdot d\vec{r} = \iint_D 1 \, dA = \text{Area of } D. (!)$$

■ Question 189.


□

An ellipse with semi-major axis a and semi-minor axis b is parametrized by $x = a \cos t$, $y = b \sin t$ for $0 \leq t \leq 2\pi$. Use \vec{F}_3 to find the area inside this ellipse.

■ **Question 190.**

□

Let C be the curve parametrized by $\vec{r}(t) = (t^2 - 3)\vec{i} + (t^3 - 4t + 1)\vec{j}$, $-2 \leq t \leq 2$. This is a closed loop. Use Green's Theorem and \vec{F}_1 to find the area inside this loop.



§31.2 Extended Versions of Green's Theorem

31.2.1 Finite union of simple regions

Although we have proved Green's Theorem only for the case where D is simple, we can now extend it to the case where D is a **finite union of simple regions**. For example, if D is the region shown in Figure 31.1, then we can write $D = D_1 \cup D_2$, where D_1 and D_2 are both simple. The boundary of D_1 is $C_1 + C_3$ and the boundary of D_2 is $C_2 + (-C_3)$. so, applying Green's Theorem to D_1 and D_2 separately, we get




Figure 31.1

$$\oint_{C_1 + C_3} \vec{F} \cdot d\vec{r} = \iint_{D_1} \text{curl}(F) dA$$

$$\oint_{C_2 + (-C_3)} \vec{F} \cdot d\vec{r} = \iint_{D_2} \text{curl}(F) dA$$

Adding the two integrals above, we get,

$$\oint_{C_1+C_2} \vec{F} \cdot d\vec{r} = \iint_D \operatorname{curl}(F) dA$$


■ **Question 191.**

□

Let C be the boundary curve of the white Yang part of the Ying-Yang symbol in the disc of radius 6. You can see in the picture that the curve C has three parts, and that the orientation of each part is given. Find the line integral of the vector field

$$\vec{F}(x, y) = \langle -y + \sin(e^x), x \rangle$$

around C . Notice that the Ying and the Yang have the same area.



■ **Question 192.**


□

Look at the shaded region G bounded by a circle of radius 2 and an inner **figure eight lemniscate** with parametric equation

$$\vec{r}(t) = \langle \sin(t), \sin(t)\cos(t) \rangle$$

with $0 \leq t \leq 2\pi$. The picture shows the curve and the arrows indicate some of the velocity vectors of the curve. Find the area of this region G .


[HINT: Use Green's theorem and the vector field $x\hat{j}$.]



■ **Question 193.**

□

Compute the line integral of $\vec{F}(x, y) = \langle 5y + 3y^2, 6xy + y^5 \rangle$ along the boundary of the green **Cyclops region** given in the figure below. There are four boundary curves, **oriented as shown in the picture**: a large ellipse of area 16, two circles of area 2 (the eyeball) and 1 (the iris) as well as a small ellipse (the mouth) of area 3.



31.2.2 Finite intersection of simple regions

We can similarly extend the theorem to regions that are finite **intersections** of simple regions.

■ **Question 194.**

Evaluate

$$\oint_C y^2 dx + 3xy dy$$

where C is the boundary of the semiannular region D in the upper half plane between $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$.




Figure 31.2

31.2.3 Not simply-connected regions

Green's Theorem can be also extended to apply to regions with holes, that is, regions that are not simply-connected. Observe that the boundary C of the region D in Figure 31.2 consists of two simple closed curves C_1 and C_2 . We assume that these boundary curves are oriented so that the region D is always on the left as the curve C is traversed. **Thus the positive direction is counterclockwise for the outer curve**

C₁ but clockwise for the inner curve C₂. If we divide D into two regions D' and D'' by means of the lines shown in Figure 31.2 and then apply Green's Theorem to each of D' and D'', we get

$$\begin{aligned}\iint_D \operatorname{curl}(F) dA &= \iint_{D'} \operatorname{curl}(F) dA + \iint_{D''} \operatorname{curl}(F) dA \\ &= \oint_{\partial D'} \vec{F} \cdot d\vec{r} + \oint_{\partial D''} \vec{F} \cdot d\vec{r}\end{aligned}$$

Since the line integrals along the common boundary lines are in opposite directions, they cancel each other out and we get

$$\iint_D \operatorname{curl}(F) dA = \oint_{C_1} \vec{F} \cdot d\vec{r} + \oint_{C_2} \vec{F} \cdot d\vec{r} = \oint_{C_1+C_2} \vec{F} \cdot d\vec{r}$$

■ Question 195.

If $\vec{F}(x, y) = \frac{-y\vec{i} + x\vec{j}}{x^2 + y^2}$, show that $\oint_C \vec{F} \cdot d\vec{r} = 2\pi$ for every positively oriented closed path C that encloses the origin.

■ Question 196.

Evaluate $\oint_C \vec{F} \cdot d\vec{r}$ where

$$\vec{F}(x, y) = \frac{2xy\vec{i} + (y^2 - x^2)\vec{j}}{(x^2 + y^2)^2}$$


and C is a positively oriented closed path that encloses the origin.

§31.3 More Practice Problems

■ Question 197.

Evaluate the following line integrals.

- (a) $\oint_C ydx + (x + y^2)dy$ where C is the ellipse $4x^2 + 9y^2 = 36$ with counterclockwise orientation.
- (b) $\oint_C \sqrt{1+x^3}dx + 2xydy$ where C is the triangle with vertices (0, 0), (1, 0), and (1, 3)
- (c) $\oint_C \vec{F} \cdot d\vec{r}$, where $\vec{F}(x, y) = \frac{(2x^3+2xy^2-2y)\vec{i} + (2y^3+2x^2y+2x)\vec{j}}{x^2+y^2}$ and C is the curve shown below.




- (d) $\oint_C \vec{F} \cdot d\vec{r}$, where $\vec{F}(x, y, z) = \langle y \cos x - xy \sin x, xy + x \cos x \rangle$, and C is the triangle from $(0, 0)$ to $(0, 4)$ to $(2, 0)$ to $(0, 0)$.

■ Question 198.

□

Suppose \vec{F} is an irrotational vector field in the plane (that is, its curl is everywhere zero) that is not defined at the origin $O = (0, 0)$. Suppose the line integral of \vec{F} along the path p from A to B is 5 and the line integral of \vec{F} along the path q from A to B is -4 . Find the line integral of F along the paths a, b and c .



Projects

§A Distances

In this project, we will use Dot product and Cross product of vectors to derive formula for calculating distances between points, lines and planes. We will use the notation $d(\cdot, \cdot)$ to denote distance.

A.1 Distance between two points

To begin with, the distance between two points P and Q with position vectors \vec{P} and \vec{Q} is simply given by

$$d(P, Q) = \|\vec{Q} - \vec{P}\| = \|\vec{PQ}\|$$

where $\|\cdot\|$ denotes the magnitude of a vector.

■ Question 1001.

Find the distance between $(-5, 2, 4)$ and $(-2, 2, 0)$.

A.2 Distance from a point to a plane

The distance of a point P from a plane Σ is defined as the length of the perpendicular from P to Σ . Suppose the plane Σ passes through a point Q and has normal vector \vec{n} .

■ Question 1002.

Explain using a picture why the distance from P to Σ is the length of the projection of \vec{PQ} onto \vec{n} . Then

$$d(P, \Sigma) = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|}$$

■ Question 1003.

Find the distance of the point $(7, 1, 4)$ from the plane $2x + 4y + 5z = 9$.

■ Question 1004.

Without the absolute sign in the numerator of the distance formula, your answer in question (2) would have been negative. What does the negative sign signify here?

A.3 Distance from a point to a line

The distance of a point P from a line \mathcal{L} is defined as the length of the perpendicular from P to \mathcal{L} . Suppose the line \mathcal{L} passes through a point Q and is parallel to a vector \vec{u} (i.e. its parametric equation looks like $\vec{r}(t) = \vec{Q} + t\vec{u}$).

■ Question 1005.

Use the definition of cross product to derive the following formula:

$$d(P, \mathcal{L}) = \frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|}$$

■ Question 1006.

Find the distance of the point $(2, 3, 1)$ from the straight line $\vec{r}(t) = (1, 1, 2) + t\langle 5, 0, 1 \rangle$.

■ **Question 1007.**

What is the equation of the plane which contains the point P and the line \mathcal{L} ?

□

A.4 Distance between two straight lines

Suppose the two straight lines \mathcal{L}_1 and \mathcal{L}_2 are given by

$$\vec{r}_1(t) = \vec{P} + t\vec{u} \quad \text{and} \quad \vec{r}_2(t) = \vec{Q} + t\vec{v}$$

i.e. the straight lines pass through P (and Q respectively) and is parallel to \vec{u} (and \vec{v} respectively).

□

■ **Question 1008.**

Draw a picture and explain using geometry why the distance between the two straight lines is given by

$$d(\mathcal{L}_1, \mathcal{L}_2) = \frac{|\vec{PQ} \cdot (\vec{u} \times \vec{v})|}{\|\vec{u} \times \vec{v}\|}$$

□

■ **Question 1009.**

Find the distance between the lines $\vec{r}_1(t) = (2, 1, 4) + t\langle -1, 1, 0 \rangle$ and $\vec{r}_2(t) = (-1, 0, 2) + t\langle 5, 1, 2 \rangle$.

□

A.5 Distance between two planes

Before deriving the formula, observe that the distance between two planes is non-zero iff the two planes are parallel to each other, in which case they have the same normal vector $\vec{n} = \langle a, b, c \rangle$. Suppose the two planes Σ_1 and Σ_2 are given by

$$ax + by + cz = d \quad \text{and} \quad ax + by + cz = e$$

■ **Question 1010.**

□

Show that the distance formula is given by

$$d(\Sigma_1, \Sigma_2) = \frac{|d - e|}{\|\vec{n}\|}$$

■ **Question 1011.**

□

Find the distance between the planes $5x + 4y + 3z = 8$ and $5x + 4y + 3z = 1$.

■ **Question 1012.**

□

Find the distance between the planes $x + 3y - 2z = 2$ and $5x + 15y - 10z = 30$.

§B Conic Sections and Quadric Surfaces

B.I Conic Sections

A **conic section** (or simply conic) is a curve obtained as the intersection of the surface of a cone with a plane. The three types of conic sections are the **hyperbola**, the **parabola**, and the **ellipse**. The circle is type of ellipse, and is sometimes considered to be a fourth type of conic section.

A cone has two identically shaped parts called **nappes**. One nappe is what most people mean by “cone”, and has the shape of a dunce hat. It can be thought of as the surface of revolution of a straight line around an axis.

- If the intersecting plane is parallel to the axis of revolution of the cone, then the conic section is a hyperbola.
- If the plane is parallel to the generating line, the conic section is a parabola.
- If the plane is perpendicular to the axis of revolution, the conic section is a circle.
- If the plane intersects one nappe at an angle to the axis (other than 90°), then the conic section is an ellipse.




Figure 3: Conic Sections

Observe that when the intersecting plane is parallel to the axis of revolution and passes through the vertex of the cone, the conic section becomes a pair of straight lines (also known as a degenerate hyperbola).

Standard Equations in Cartesian Coordinates

The **Major Axis** is the chord between the two vertices: the longest chord of an ellipse, the shortest chord between the branches of a hyperbola. The **Minor Axis** is the shortest chord of an ellipse.

- In each of the above cases the center of the conic is at the origin. If the curve is translated h units horizontally and k units vertically, its new equation is obtained by replacing x with $(x - h)$ and y with $(y - k)$.

Conic Type	Standard Equation	Major Axis	Minor Axis
Circle	$x^2 + y^2 = r^2, \quad r \geq 0$	$2r$	$2r$
Ellipse	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \quad a, b > 0$	$2a$	$2b$
Parabola	$y^2 = 4ax$ $x^2 = 4ay$	N/A	N/A
Hyperbola	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \quad a, b > 0$ $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1, \quad a, b > 0$	$2a$	N/A
Pair of Straight Lines	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 0, \quad a, b > 0$ $\iff y = \pm \frac{b}{a}x$	N/A	N/A

■ Question 1013.

□

Use your precalculus memory or your favorite computer graphics software (e.g. DESMOS) to draw a picture of each of the above conic sections.

■ Question 1014.

□

For each of the following curves, find out what kind of conic section it is. See if you can answer without using a computer first.

- (a) $(x - 3)^2 + (y - 4)^2 = y^2$
- (b) $(x - 3)^2 + (y - 4)^2 = 2y^2$
- (c) $(x - 3)^2 + (y - 4)^2 = \frac{y^2}{2}$
- (d) $(x - 3)^2 + (y - 4)^2 = 1$

■ Question 1015.

□

Find the range of value(s) of p for which the curve $\frac{x^2}{9-p} + \frac{y^2}{p-3} = 1$ looks like a

- (a) Circle.
- (b) Ellipse.
- (c) Hyperbola.

B.2 Quadric Surface

Equations of surfaces in three dimension are of the form $f(x, y, z) = c$. One special case of interest is when $f(x, y, z)$ is a polynomial which is not linear and quadratic at most, containing terms involving x, y, z, x^2, y^2 , and z^2 only. This kind of surface is called a **Quadric Surface**. Quadric surfaces are often used as example surfaces since they are relatively simple. There are nine different basic quadric surfaces listed below. A catalog of the equations and pictures of the quadric surfaces is available on section 2.6 of your openstax textbook.

- **Cylinders:** A cylinder basically has no control over one of the variables. Take some sort of a curve in the plane, and draw a family of parallel lines so that each of the lines intersects the curve in a point.

For example, a cylinder over the line $(0, t, t)$ would be all points of the form (s, t, t) for any values of s and t . This is a plane (a plane is a cylinder over a line!), and has the equation $y = z$.

The most common picture you think of when you hear the term cylinder, is that of a cylinder over a circle. Some basic variations are

- **elliptical cylinder:** A cylinder over an ellipse
- **parabolic cylinder:** A cylinder over a parabola
- **hyperbolic cylinder:** A cylinder over a hyperbola
- **Ellipsoid**, the three-dimensional analogue of the ellipse. A sphere is an uniform ellipsoid.
- **Elliptic paraboloid**, a sort of cup or a bowl
- **Hyperbolic paraboloid**, looks like a horse-saddle or a pringle
- **Cone**, take a straight line intersecting the z -axis and consider its surface of revolution around the z -axis
- **Hyperboloids:** In three dimensions there are two different analogs of hyperbolas. The word "sheet" is used in an antique, specialized sense with surfaces: it means one connected "piece" of a surface. So a hyperboloid with one sheet is a surface with one (connected) piece, and a hyperboloid with two sheets is a surface with two (connected) pieces.
 - **hyperboloid of one sheet**, obtained by revolving a hyperbola around its minor axis. The surface is connected but there is a hole in it.
 - **hyperboloid of two sheet**, obtained by revolving a hyperbola around its major axis. This surface has two pieces.

Warning: In each case, note the direction of the axes relative to the surfaces and how the corresponding variables show up in the equation. For example, $z = 2y^2 - x^2$ is a hyperbolic paraboloid that goes downward in the x -axis direction and upwards in y -axis direction. The equation $y = 2x^2 - z^2$ is also a hyperbolic paraboloid that goes downward in the z -axis direction and upwards in x -axis direction. Similarly, $y = x^2 + z^2$ is an elliptical paraboloid that opens in the y -axis direction.

■ Question 1016.



Use CalcPlot3D to plot each of the following quadric surfaces, then use the catalog to decide which type of quadric surface it resembles.

- (a) $\frac{x^2}{9} - \frac{y^2}{16} = z$.
- (b) $\frac{x^2}{4} + \frac{y^2}{25} + \frac{z^2}{9} = 1$.
- (c) $\frac{x^2}{4} + \frac{y^2}{9} = \frac{z}{2}$.
- (d) $\frac{z^2}{4} - x^2 - \frac{y^2}{4} = 1$.
- (e) $x^2 + \frac{y^2}{9} = \frac{z^2}{16}$.

■ Question 1017.



For each of the above surfaces, describe what kind of conic sections are obtained by taking its cross-sections parallel to the YZ -plane, XZ -plane and XY -plane. These are called **traces**. Create a table to collect all your responses.

- Intersection with $x = k$:
- Intersection with $y = k$:
- Intersection with $z = k$:

■ Question 1018.

Now try identifying the following surfaces without the help of the catalog to see if you remember the names. Try to do this **without** graphing them first.

- (a) $9y^2 + 4z^2 = 36$
(b) $y^2 + 2y + z^2 = x^2$
(c) $4x^2 - y^2 + z^2 + 9 = 0$

■ Question 1019.

A non-basic Quadric Surface Use CalcPlot3D to plot the surface

$$x^2 - 17z^2 - 2y^2 - 2xz - 12yz - 1 = 0$$

Which of the nine basic quadric surface does this resemble most closely? You may need to rotate and translate the surface to make it look like one of the basic ones.

§C Epicycloids

Consider a (black) circle of radius R with its center at the origin O . A (blue) circle of radius r rolls around the **outside** of the circle of radius R . See figure 4 for diagrams of different values of r . A (red) point P is located on the circumference of the rolling circle. The path traced out by P is called an **epicycloid**.




Figure 4: Epicycloids for $R/r = 1, 2, 3, 4$

Assume that initially at time $t = 0$, the rolling circle sits to the right of the fixed circle and the point P is located at $(R, 0)$. After the rolling circle has moved a bit, draw a line from the center O of the large circle to the point of contact with the rolling circle and let t be the angle the line makes with the positive X-axis. If the location of P at this moment is given by $(x(t), y(t))$ (see figure 5), then the parametric equation of the **epicycloid** is given by

$$x(t) = (R + r) \cos t - r \cos \frac{(R + r)t}{r}$$

$$y(t) = (R + r) \sin t - r \sin \frac{(R + r)t}{r}$$

Warning: If the initial location of P and the rolling circle is chosen differently, you will get the same shape with a different orientation and the form of the parametric equations will change slightly. For example, if the \sin and \cos functions are interchanged ($\sin \leftrightarrow \cos$), we get a vertically oriented epicycloid. If we replace t with $t + \varphi$ we get an epicycloid that has been rotated by angle φ .

■ Question 1020.

Use the diagram above to analyze the geometry of the epicycloid. Write the detailed steps for the derivation of the parametric form of the epicycloid.

[**Hints:** The length of the blue arc and the green arc are equal, why? Use this to derive the angle $\angle PCA$ in terms of t . The x -coordinate of P is given by BD which is equal to $BC - CD$. The y -coordinate is equal to $OB - PD$.]




Figure 5

■ Question 1021.

Suppose $\frac{R}{r} = m$, a positive integer. Show that the number of times the point P touches the fixed circle is exactly m .

[**Hints:** When the point P touches the big fixed circle, $x(t) = R \cos t$ and $y(t) = R \sin t$ (why?).]

■ Question 1022.

Now we will use Mathematica to draw a couple of different epicycloids when $\frac{R}{r}$ is a rational number $\frac{p}{q}$ (in its lowest terms) for different values of p and q .

- First set $r = 2$, and let $R = 3$. What do you observe? Does the epicycloid complete a full revolution (i.e. does the point P completes a full cycle) for $t \in [0, 2\pi]$?
- Next set $r = 2$, and let $R = 4, 5, 6$. What do you observe? What are the values of p and q in each case? What does the quantity p correspond to in the pictures?
- Next set $R = 5$, and let $r = 2, 3, 4, 5$. Note that R/r has to be at least than 1 for an epicycloid to form. What do you observe? What does the quantity q correspond to in the pictures?

■ Question 1023.

Try plotting the curve with $R = e, r = 1$. How long does it take to complete a full revolution? Make a conjecture about what happens in general if $\frac{R}{r}$ is an irrational number.

■ Question 1024.

Find the length of the arc of the epicycloid between two points where it touches the fixed circle for a general value of $m = \frac{R}{r}$. For example, when $R/r = 3$, there are three such equal length arcs.