

MATH 1800 PROJECT 1: DISTANCES

Subhadip Chowdhury

In this project, we will use Dot product and Cross product of vectors to derive formula for calculating distances between points, lines and planes. We will use the notation $d(\cdot, \cdot)$ to denote distance.

DISTANCE BETWEEN TWO POINTS

To begin with, the distance between two points P and Q with position vectors \vec{P} and \vec{Q} is simply given by

$$d(P, Q) = \|\vec{Q} - \vec{P}\| = \|\vec{PQ}\|$$

where $\|\cdot\|$ denotes the magnitude of a vector.

Question 0

Find the distance between $(-5, 2, 4)$ and $(-2, 2, 0)$.

DISTANCE FROM A POINT TO A PLANE

The distance of a point P from a plane Σ is defined as the length of the perpendicular from P to Σ . Suppose the plane Σ passes through a point Q and has normal vector \vec{n} .

Question 1

Explain using a picture why the distance from P to Σ is the length of the projection of \vec{PQ} onto \vec{n} . Then

$$d(P, \Sigma) = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|}$$

Question 2

Find the distance of the point $(7, 1, 4)$ from the plane $2x + 4y + 5z = 9$.

Question 3

Without the absolute sign in the numerator of the distance formula, your answer in question (2) would have been negative. What does the negative sign signify here?

DISTANCE FROM A POINT TO A LINE

The distance of a point \mathbf{P} from a line \mathcal{L} is defined as the length of the perpendicular from \mathbf{P} to \mathcal{L} . Suppose the line \mathcal{L} passes through a point \mathbf{Q} and is parallel to a vector \vec{u} (i.e. its parametric equation looks like $\vec{r}(t) = \vec{Q} + t\vec{u}$).

Question 4

Use the definition of cross product to derive the following formula:

$$d(\mathbf{P}, \mathcal{L}) = \frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|}$$

Question 5

Find the distance of the point $(2, 3, 1)$ from the straight line $\vec{r}(t) = (1, 1, 2) + t(5, 0, 1)$.

Question 6

What is the equation of the plane which contains the point \mathbf{P} and the line \mathcal{L} ?

DISTANCE BETWEEN TWO STRAIGHT LINES

Suppose the two straight lines \mathcal{L}_1 and \mathcal{L}_2 are given by

$$\vec{r}_1(t) = \vec{P} + t\vec{u} \quad \text{and} \quad \vec{r}_2(t) = \vec{Q} + t\vec{v}$$

i.e. the straight lines pass through \mathbf{P} (and \mathbf{Q} respectively) and is parallel to \vec{u} (and \vec{v} respectively).

Question 7

Draw a picture and explain using geometry why the distance between the two straight lines is given by

$$d(\mathcal{L}_1, \mathcal{L}_2) = \frac{|\vec{PQ} \cdot (\vec{u} \times \vec{v})|}{\|\vec{u} \times \vec{v}\|}$$

Question 8

Find the distance between the lines $\vec{r}_1(t) = (2, 1, 4) + t(-1, 1, 0)$ and $\vec{r}_2(t) = (-1, 0, 2) + t(5, 1, 2)$.

DISTANCE BETWEEN TWO PLANES

Before deriving the formula, observe that the distance between two planes is non-zero iff the two planes are parallel to each other, in which case they have the same normal vector $\vec{n} = \langle a, b, c \rangle$. Suppose the two planes Σ_1 and Σ_2 are given by

$$ax + by + cz = d \quad \text{and} \quad ax + by + cz = e$$

Question 9

Show that the distance formula is given by

$$d(\Sigma_1, \Sigma_2) = \frac{|d - e|}{\|\vec{n}\|}$$

Question 10

Find the distance between the planes $5x + 4y + 3z = 8$ and $5x + 4y + 3z = 1$.

Question 11

Find the distance between the planes $x + 3y - 2z = 2$ and $5x + 15y - 10z = 30$.