

Lab 1: 3D Graphing with Mathematica

Subhadip Chowdhury

- Sign on to an iMac with your username and password.
- When you open *Mathematica*, click **New Document** and a blank screen will appear. This is known as a notebook. If you need to open a new notebook, go to **File**, choose **New** and choose **Notebook** or simply hit **Command + N** (or **Ctrl + N** for **Windows**).
- Make the Untitled window bigger if necessary by dragging the lower right corner. Choose at least 125% from the lower left of the Untitled window for a comfortable viewing size.
- Do not forget to **Save** the notebook periodically. Save new notebooks on the **Desktop**. The **Save** command is under the **File** menu. Give your file a name in the following format:

Lab0_Name1_Name2.nb

- Follow the instructions in the paper copy of the handout.
- At the end of Lab session, save and quit Mathematica. **Do not delete your file or any calculation you did.** Then log off or restart the computer. Do NOT click **Shut Down**.

We'll be learning to use two main commands in this lab. The first one (**Plot3D**) produces 3D surfaces, and the second (**ContourPlot**) produces contour plots for functions of two variables.

Exercise 1: Defining custom functions

1. To define a function $f(x, y)$ we write

$f[x_, y_] := \text{definition}$

Note the 'underscore' after x and y in the left hand side. Type

$f[x_, y_] := x^2 + y^2$

to define $f(x, y) = x^2 + y^2$.

2. Type $f[8, 6]$ to check that it gives the correct output of 100. Find square root of $f(12, 5)$ using Mathematica. Remember to use brackets and not parentheses.
3. Use Mathematica to calculate $\sin^3(f(2.31, 4.53))$ up to the 4th decimal point.

Exercise 2: 3D Graphing

1. Type

$\text{Plot3D}[f[x, y], \{x, -1, 1\}, \{y, -1, 1\}]$

to plot the graph of $f(x, y)$ over the domain $-1 \leq x \leq 1, -1 \leq y \leq 1$.

2. Type

$\text{ContourPlot}[f[x, y], \{x, -1, 1\}, \{y, -1, 1\}]$

to produce the contour plot of the same function.

We are going to explore the graph of the function

$$M(x, y) = 3xy^2 - x^3$$

The graph of this function is called the "monkey saddle".

3. Define the function M in Mathematica and plot the graph of M over the domain $\{(x, y) \mid -10 \leq x, y \leq 10\}$. Can you explain why it's called the "monkey saddle"?
4. Draw the **ContourPlot** of the monkey saddle function.
5. Explain what the cross sections parallel to x , y , and z axis look like.

Exercise 3: More on Contour Plots

Use the following format to get a better picture for the contour diagram. Explore other customization options as well.

```
ContourPlot[f[x,y], {x, -10, 10}, {y, -10, 10},  
ContourLabels -> True, ContourStyle -> Thickness[.005]]
```

Draw the graph and the contour plots of the following functions.

6. $2x + y + 4$

7. $\frac{x}{2} + \frac{y}{4} + 1$

8. Why do you get similar contour plots both times?

There are rare occasions when you may wish to use a set of non-evenly-spaced z values to compute contour lines. For the last example, we are going to start with evenly-spaced z values 1, 2, ..., 9, 10, but then switch to 10, 20, ..., 50, 60. The reason is when z gets large, the corresponding contour lines become too close together, hence it makes sense to "change the scale" by upping the Δz value from 1 to 10.

9. First plot the graph and the contour plot of $x^2 + 2y^2$ as before. Observe why we need to display contours selectively.

10. Next, add the option

```
Contours -> {1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60},
```

to selectively display contours in the last picture.

Exercise 4: Level Surface

The `ContourPlot3D` command is designed to plot the surfaces consisting of all the input triples (x, y, z) generating the same output from a function of three variables $g(x, y, z)$ (these are called "level surfaces").

11. Type the following and compare your output to problem 1.

```
ContourPlot3D[x^2 + y^2 - z == 0, {x, -1, 1}, {y, -1, 1}, {z, 0, 1}]
```

12. Recall that the graph of f is defined by the assignment $z = f(x, y)$. Use a simple algebra manipulation to show that, for the three-variable function defined by $g(x, y, z) = f(x, y) - z$, the set of triples (x, y, z) satisfying $g(x, y, z) = 0$ is the same as the set of points on the graph of f . Thus the 'level surface' of g at level 0 becomes the graph of f .

Exercise 5: Exploratory Plotting

Use the graphing commands to plot some surfaces. For instance, try plotting

13. a plane which has its x , y , and z intercepts all positive

14. a hemispherical dome

15. a cone