

# STAT 320: Principles of Probability

## Unit 1: A few Definitions and Notations from Set Theory

United Arab Emirates University

Department of Statistics

# Outline


- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product


## Definition of a Set


# Definition of a Set


## Definition ( Set)


*A Set is a well defined collection of **objects**.*

 **Elements:** The objects inside a set are called the **elements** of the Set


 Usually we denote sets with upper-case letters, elements with lower-case letters.

 The following notation is used to show a set membership.

  $x \in A$  :means that  $x$  is a member of the set  $A$ .

  $x \notin A$  :means that  $x$  is NOT a member of the set  $A$ .

# Venn Diagrams

 **Venn Diagrams** are graphical representation of the sets that are often used to depict the relation between various sets

For the rest of the slides, we shall include ‘Venn Diagrams’ for visualization of different set-theoretic concepts.

# Example

# Example: Different Ways of Describing a Set

- 1 Listing all the elements:  $A = \{1, 2, 3, 4, 5, 6\}$
- 2 Give a verbal description: A is the set of all integers from 1 to 6, inclusive.
- 3 Give a mathematical inclusion rule:

$$A = \{x \text{ is an Integer} : 1 \leq x \leq 6\}$$

# Example: Different Ways of Describing a Set

- 1 Listing all the elements:  $A = \{1, 2, 3, 4, 5, 6\}$
- 2 Give a verbal description: A is the set of all integers from 1 to 6, inclusive.
- 3 Give a mathematical inclusion rule:

$$A = \{x \text{ is an Integer} : 1 \leq x \leq 6\}$$



# Example: Different Ways of Describing a Set

- 1 Listing all the elements:  $A = \{1, 2, 3, 4, 5, 6\}$
- 2 Give a verbal description: A is the set of all integers from 1 to 6, inclusive.
- 3 Give a mathematical inclusion rule:

$$A = \{x \text{ is an Integer} : 1 \leq x \leq 6\}$$

# The Null Sets and the Universal Set

## Definition (Null Set)

*The **Null Set** (or also called the **Empty Set**) is a set with no elements in it.*

It is often denoted by  $\emptyset$

## Definition (Universal Set)

*The **Universal Set** is the set of all elements currently under consideration.*

In the context of probability theory, it is often denoted by  $\mathcal{S}$  (or  $\Omega$ )

The universal set contains all of the elements relevant to a given discussion.

# The Null Sets and the Universal Set

## Definition (Null Set)

*The **Null Set** (or also called the **Empty Set**) is a set with no elements in it.*

It is often denoted by  $\emptyset$

## Definition (Universal Set)

*The **Universal Set** is the set of all elements currently under consideration.*

In the context of probability theory, it is often denoted by  $\mathcal{S}$  (or  $\Omega$ )

The universal set contains all of the elements relevant to a given discussion.

# Outline

- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product

## Relationship Between Sets

**Subset:** A set  $A$  is called a subset of a set  $B$  if all the elements of  $A$  are also the elements of  $B$ .

If  $x \in A \implies x \in B$  then  $A \subseteq B$ .

**Subsets:** If  $A$  is a subset of  $B$  then we write  $A \subseteq B$ .

Note: The notation for subset is very similar to the notation for "less than or equal to" and means, "included in or equal to" in terms of the sets.



We write  $A \not\subseteq B$  if  $A$  is not included in  $B$

**Subset:** A set  $A$  is called a subset of a set  $B$  if all the elements of  $A$  are also the elements of  $B$ .

If  $x \in A \implies x \in B$  then  $A \subseteq B$ .

**Subsets:** If  $A$  is a subset of  $B$  then we write  $A \subseteq B$ .

Note: The notation for subset is very similar to the notation for "less than or equal to" and means, "included in or equal to" in terms of the sets.



We write  $A \not\subseteq B$  if  $A$  is not included in  $B$



We say "**A is a proper subset of B**" if all the elements of A are also the elements of B, but in addition there exists at least one element  $c$  such that  $c \in B$  but  $c \notin A$ .



**Proper Subset:**  $A \subset B$ , then we say **A is a proper subset of B**.

**Comment:** The notation for subset is very similar to the notation for "less than", and means, in terms of the sets, "included in but not equal to".



# Subset: Example

# Outline

- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product


## Few Set Operations

# Set Operation: Union

## A Union B:

A Union B, denoted by  $A \cup B$  is the set of all elements that are either in  $A$ , or in  $B$ , or inside both the sets.

The logical operator is OR.








# Set Operation: Union

**A Union B:**

A Union B, denoted by  $A \cup B$  is the set of all elements that are either in  $A$ , or in  $B$ , or inside both the sets.

The logical operator is **OR**.






# Set Operation: Intersection

**A Intersection B:** A Intersection B, denoted by  $A \cap B$  is the set of all elements that are both A and B.


The logical operator is AND .







# Set Operation: Intersection

**A Intersection B:** A Intersection B, denoted by  $A \cap B$  is the set of all elements that are both A and B.

The logical operator is **AND**.








# Set Operation: Complement

**A Complement :** A Complement, denoted by  $\bar{A}$  is the set of all elements that are not in  $A$ . Sometimes it is also denoted by  $A^c$



The logical operator is NOT .



# Set Operation: Complement

**A Complement :** A Complement, denoted by  $\bar{A}$  is the set of all elements that are not in  $A$ . Sometimes it is also denoted by  $A^c$

The logical operator is **NOT**.





# Set Operation: Set Difference

## A Minus B:

A Minus B, denoted by  $A - B$  is the set of all elements that are only an element of A, but not an element of the set B.

$A - B$  is same as:  $A \cap \bar{B}$ .

Comment:  $A - B$  is not same as  $B - A$





# Set Operation: Set Difference

## A Minus B:

A Minus B, denoted by  $A - B$  is the set of all elements that are only an element of A, but not an element of the set B.

$A - B$  is same as:  $A \cap \bar{B}$ .

Comment:  $A - B$  is not same as  $B - A$



# A Few Rules for Various Set Operations

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$\overline{(A \cap B)} = (\overline{A} \cup \overline{B})$$

$$\overline{(A \cup B)} = (\overline{A} \cap \overline{B})$$


# Set Operation: Examples

# Outline


- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product



## A Few Examples Using Venn Diagrams

 **Venn Diagrams** are graphical representation of the sets that are typically used to depict the relation between various sets


# Venn Diagrams: Examples



# Venn Diagrams: Examples

Let  $S = \{1, 2, 3, 4, 5, 6, 7, 8\}$ ,

$A = \{1, 2, 6, 7\}$ ,  $B = \{2, 3, 4, 7\}$ , and  $C = \{4, 5, 6, 7\}$



# Outline

- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product

## Disjoint Sets & Partition

# Disjoint Sets

**Disjoint Sets:**

Two sets  $A$ , and  $B$  are said to be **Disjoint Sets** or **mutually exclusive sets** if  $A$  and  $B$  does not have any elements in common.

$A$  and  $B$  are Disjoint  $\Leftrightarrow A \cap B = \emptyset$ .


# Disjoint Sets

**Disjoint Sets:**

Two sets  $A$ , and  $B$  are said to be **Disjoint Sets** or **mutually exclusive sets** if  $A$  and  $B$  does not have any elements in common.

$A$  and  $B$  are Disjoint  $\Leftrightarrow A \cap B = \emptyset$ .






# Exhaustive

**Exhaustive:** Two sets A and B are exhaustive for the set C if  $A \cup B = C$ .

A and B are exhaustive for C  $\Leftrightarrow A \cup B = C$ .



# Partition


**Partition:** A group of sets  $\{A, B\}$  is called a **partition** for a set  $C$  if

1

$A \cap B = \emptyset$  ( i.e. ,  $A$ , and  $B$  are Disjoint) , and


2

$A \cup B = C$  (i.e. ,  $A$  and  $B$  is exhaustive for  $C$ ).



# Example

$$S = \{1, 2, 3, 4, 5, 6\}$$



$$A_1 = \{1, 2\}, A_2 = \{3\}, A_3 = \{4\}, A_4 = \{5, 6\}$$

Is  $A_1 = \{1, 2, 3\}, A_2 = \{3\}, A_3 = \{4\}, A_4 = \{4, 5, 6\}$  a partition of  $S$ ?

# Examples

# A Few Questions:

Let  $\Omega$  be the universal set and  $\emptyset$  denotes the empty set.

- $A \cup \emptyset =$
- $A \cup \overline{A} =$
- $A \cap \emptyset =$
- $A - \overline{A} =$
- $A \cap \overline{A} =$
- $A \cup \Omega =$
- $A \cap \Omega =$
- If  $A \subset B$  then  $A \cap B =$
- If  $A \subset B$  then  $A \cup B =$




# Outline


- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product

## Real Numbers ( $\mathbb{R}$ ) & Intervals

# Real Numbers, Intervals

For the definitions below, assume  $a \in \mathbb{R}$ ,  $b \in \mathbb{R}$ , and  $a < b$ .

  $[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$

  $(a, b] := \{x \in \mathbb{R} : a < x \leq b\}$

  $[a, b) := \{x \in \mathbb{R} : a \leq x < b\}$

  $(a, b) := \{x \in \mathbb{R} : a < x < b\}$

# Examples of Intervals

$[0, 1] := \{x \in \mathbb{R} : 0 \leq x \leq 1\}$  = Set of all real numbers from 0 to 1, including the numbers 0 and 1.

$(0, 1) := \{x \in \mathbb{R} : 0 < x < 1\}$  = Set of all real numbers between 0 to 1. It does not include the numbers 0, and 1.

$(0, \infty) := \{x \in \mathbb{R} : 0 < x\}$  = Set of all positive real numbers. The interval **does not include 0**

$[0, \infty) := \{x \in \mathbb{R} : 0 \leq x\}$  = Set of all non-negative real numbers. The interval **does include 0**

# Examples: Union & Intersection of Intervals

# Outline

- 1 Definition of Sets
- 2 Relationship Between Sets
- 3 A Few Set Operations
- 4 A Few Examples Using Venn Diagrams
- 5 Disjoint Sets & Partition
- 6 Real Numbers & Intervals
- 7 The Notion of Cartesian Product

## The Notion of Cartesian Product

# The Notion of 'Tuple'

**Tuple:** Let  $k$  be an integer. A  **$k$ -tuple** is the a the ordered sequence of values often written inside parenthesis while different elements are separated by comma.

Example:

- $(1.5, 2)$  is a **2-tuple**.
- $(1.5, 2, 6, 2)$  is a **4-tuple**.
- $(H, T, H, H, H)$  is a **5-tuple**



# Set Operation: Cartesian Product

## Cartesian Product of A and B:

Cartesian Product of A and B, denoted by  $A \times B$  is the set of all “two-tuple” objects where the first element in the tuple is from the set A and the second element is taken from the set B.

$$A \times B = \{(x, y) : x \in A, y \in B\}.$$

**Comment:** In general  $A \times B$  is not same as  $B \times A$ . However,  $A \times A$  is denoted by  $A^2$ . Note that A is a set and not a number.

## Cartesian Product of Multiple Sets

Let  $A_1, A_2, \dots, A_n$  are non empty sets. Then their Cartesian product is defined as following:

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for } i = 1, \dots, n\}.$$

# Set Operation: Cartesian Product

## Cartesian Product of $A$ and $B$ :

Cartesian Product of  $A$  and  $B$ , denoted by  $A \times B$  is the set of all “two-tuple” objects where the first element in the tuple is from the set  $A$  and the second element is taken from the set  $B$ .

$$A \times B = \{(x, y) : x \in A, y \in B\}.$$

**Comment:** In general  $A \times B$  is not same as  $B \times A$ . However,  $A \times A$  is denoted by  $A^2$ . Note that  $A$  is a set and not a number.

## Cartesian Product of Multiple Sets

Let  $A_1, A_2, \dots, A_n$  are non empty sets. Then their Cartesian product is defined as following:  
 $A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for } i = 1, \dots, n\}.$

# A Few Standard Notation

$\mathbb{Z}$  : Set of all Integers. i.e.  $\mathbb{Z} = \{0, -1, 1, -2, 2, -3, 3, \dots\}$ .

$\mathbb{Z}_+$  : Set of all non-negative Integers. i.e.  $\mathbb{Z}_+ = \{1, 2, 3, \dots\}$ .

$\mathbb{R}$  : Set of all real numbers.

$\mathbb{R}_+$  : Set of all positive real numbers.

- $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R} = \{(x_1, x_2) : x_1 \in \mathbb{R}, x_2 \in \mathbb{R}\}$

- $\mathbb{R}^3 = \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{(x_1, x_2, x_3) : x_1 \in \mathbb{R}, x_2 \in \mathbb{R}, x_3 \in \mathbb{R}\}$

- $\mathbb{Z}_+^2 = \mathbb{Z}_+ \times \mathbb{Z}_+ = \{(z_1, z_2) : z_1 \in \mathbb{Z}_+, z_2 \in \mathbb{Z}_+\}$

# A Few Standard Notation

$\mathbb{Z}$  : Set of all Integers. i.e.  $\mathbb{Z} = \{0, -1, 1, -2, 2, -3, 3, \dots\}$ .

$\mathbb{Z}_+$  : Set of all non-negative Integers. i.e.  $\mathbb{Z}_+ = \{1, 2, 3, \dots\}$ .

$\mathbb{R}$  : Set of all real numbers.

$\mathbb{R}_+$  : Set of all positive real numbers.

- $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R} = \{ (x_1, x_2) : x_1 \in \mathbb{R}, x_2 \in \mathbb{R} \}$

- $\mathbb{R}^3 = \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{ (x_1, x_2, x_3) : x_1 \in \mathbb{R}, x_2 \in \mathbb{R}, x_3 \in \mathbb{R} \}$

- $\mathbb{Z}_+^2 = \mathbb{Z}_+ \times \mathbb{Z}_+ = \{ (z_1, z_2) : z_1 \in \mathbb{Z}_+, z_2 \in \mathbb{Z}_+ \}$

## Discussion on Cardinality of Sets

# A Discussion on Cardinality of Various Sets

Questions?