

STAT 320: Principles of Probability

Unit 5(PART:B)

A Few Discrete Random Variables

United Arab Emirates University

Department of Statistics

Outline

- 1 Binomial Distribution
- 2 Poisson Distribution
- 3 Geometric Distribution
- 4 Negative Binomial Distribution
- 5 Miscellaneous Problems

Binomial Distribution



A quality controlling team in TV manufacturer evaluates a products before they are send to market. They are interested in idensifying the number of defective products and the resources that should be optimally allocated to mitigate the defectives items.



A broker in a stock exchange speculates how many shares out of 10 newly introduced shares will go up next day.






A airline company is interested in identifying the number of last minute cancellations that may take place.



A car insurance company have sold 2000 insurance policies on a specific new cars. They need to estimate the funts that should be made available/reserved to compensate the losses that may occur to the insured cars during the next one year.

A Bernoulli Trial/ Experiment

-  The random experiment has only two outcomes. Namely **SUCCESS**, and **FAILURE**
-  Events corresponding to the successive trials/experiments are statistically independent.
-  All such trials/experiments have the **same chance/probability of success**.

If a sequence of n independent Bernoulli trials is performed under the same condition, then the random variable that records the **total number of successes** is called the Binomial Random variable.

Binomial Distribution

Definition (Binomial Experiment)

An experiment is called a Binomial experiment if it satisfies the following 4 conditions:

- The experiment consists of n Bernoulli trials.
- Each trial results in a success (S) or a failure (F).
- The trials are independent.
- The probability of a success, π , is fixed throughout n trials.

Binomial Distribution $\text{Binomial}(n, \pi)$

- 1 Given a Binomial experiment consisting of n Bernoulli trials with success probability π , the Binomial random variable X associated with this experiment is defined as the number of successes among the n trials.
- 2 The random variable X has the Binomial Distribution with parameters n and π ; denoted by $X \sim \text{Binomial}(n, \pi)$.
- 3 The behavior of Binomial Distribution with different n and π .

Binomial Distribution $\text{Binomial}(n, \pi)$

Definition (Binomial Distribution)

Let $\pi \in (0, 1)$, then the probability mass function of $\text{Binomial}(n, \pi)$ is given by

$$p(x) := \binom{n}{x} \pi^x (1 - \pi)^{n-x}, \text{ for } x \in \mathbb{S}_x, \text{ where } \mathbb{S}_x = \{0, 1, \dots, n\}$$

Mean

$$E(X) = n\pi$$

Variance

$$\text{VAR}(X) = n\pi(1 - \pi)$$

Expected Value of Binomial Distribution

$$\begin{aligned} E(X) &:= \sum_{y \in \mathbb{S}_X} y p_X(y) \\ &= \sum_{y=0}^n y \binom{n}{y} \pi^y (1 - \pi)^{n-y} \\ &= (1 - \pi)^n \sum_{y=0}^n y \binom{n}{y} \left(\frac{\pi}{1 - \pi} \right)^y \\ &= (1 - \pi)^n \frac{np}{(1 - \pi)^n} \\ &= np \end{aligned} \tag{1}$$

Expected Value of Binomial Distribution

$$\begin{aligned} E(X^2) &:= \sum_{y \in \mathbb{S}_X} y^2 p_X(y) \\ &= \sum_{y=0}^n y^2 \binom{n}{y} \pi^y (1 - \pi)^{n-y} \\ &= (1 - \pi)^n \sum_{y=0}^n y^2 \binom{n}{y} \left(\frac{\pi}{1 - \pi} \right)^y \\ &= (1 - \pi)^n \frac{np + n(n-1)\pi^2}{(1 - \pi)^n} \\ &= np + n(n-1)\pi^2 \end{aligned} \tag{2}$$

$$\text{Var}(X) = E(X^2) - (E(X))^2 = np + n(n-1)\pi^2 - n^2\pi^2 = np - np^2 = np(1 - \pi).$$

Expected Value of Binomial Distribution

$$\begin{aligned}M_X(t) &:= \sum_{y \in \mathbb{S}_X} e^{ty} p_X(y) \\&= (1 - \pi)^n \sum_{y=0}^n e^{ty} \binom{n}{y} \left(\frac{\pi}{1 - \pi} \right)^y \\&= (1 - \pi)^n \sum_{y=0}^n \binom{n}{y} \left(\frac{pe^t}{1 - \pi} \right)^y \\&= (1 - \pi)^n \sum_{y=0}^n e^{ty} \binom{n}{y} \left(\frac{\pi}{1 - \pi} \right)^y \\&= (1 - \pi)^n \left(1 + \frac{pe^t}{1 - \pi} \right)^n = (1 - \pi + pe^t)^n\end{aligned}$$

(3)

Example

Example : Five fair coins are flipped. If the outcomes are assumed independent.

- 1 Find the probability mass function of the number of heads obtained.
- 2 Find the probability that at least 3 heads are obtained.
- 3 Find the probability that at most 2 heads are obtained.

Example

Example : Five fair coins are flipped. If the outcomes are assumed independent.

- 1 Find the probability mass function of the number of heads obtained.
- 2 Find the probability that at least 3 heads are obtained.
- 3 Find the probability that at most 2 heads are obtained.

Solution: Let X = The number of heads in 5 tossed coins. $X \sim \text{Binomial}(n = 5, \pi = 0.5)$.

- 1 $P(X = 0) = 0.5^5 = 0.0313$
- 2 $P(X = 1) = \binom{5}{1} 0.5^5 = 0.1563$
- 3 $P(X = 2) = \binom{5}{2} 0.5^5 = 0.3125$
- 4 $P(X = 3) = \binom{5}{3} 0.5^5 = 0.3125$
- 5 $P(X = 4) = \binom{5}{4} 0.5^5 = 0.1563$
- 6 $P(X = 5) = \binom{5}{5} 0.5^5 = 0.0313$

Example

Example :

It is known that screws produced by a certain company will be defective with probability .01, independently of each other. The company sells the screws in packages of 10 and offers a money-back guarantee that at most 1 of the 10 screws is defective. What proportion of packages sold must the company replace? Use the Binomial Calculator or Statistical Tables.

Example

Example :

The following gambling game, known as the wheel of fortune (or chuck-a-luck), is quite popular at many carnivals and gambling casinos: A player bets on one of the numbers 1 through 6. Three dice are then rolled, and if the number bet by the player appears i times, $i = 1; 2; 3$, then the player wins i units; if the number bet by the player does not appear on any of the dice, then the player loses 1 unit. Is this game fair to the player?

Outline

- 1 Binomial Distribution
- 2 Poisson Distribution
- 3 Geometric Distribution
- 4 Negative Binomial Distribution
- 5 Miscellaneous Problems

Poisson Distribution

Poisson Distribution

The Poisson distribution models the number of occurrences of an event when there is a known average rate per unit time or space λ .

Definition (Poisson Distribution)

The requirements for a Poisson distribution are that:

- 1 no two events can occur simultaneously,
- 2 events occur independently in different intervals, and
- 3 the expected number of events in each time interval remain constant.

Poisson Distribution: pmf, Expected Value

The Poisson distribution models the number of occurrences of an event when there is a known average rate per unit time or space λ .

Definition (Poisson Distribution: pmf, Expected Value)

The requirements for a Poisson distribution are that:

- 1 The probability mass function of $\text{Poisson}(\lambda)$ is given by

$$p(x) = \frac{e^{-\lambda} \lambda^x}{x!} \text{ for } x = 0, 1, 2, 3, \dots$$

- 2 If $X \sim \text{Poisson}(\lambda)$, then $E(X) = \lambda$, and $\text{Var}(X) = \lambda$.

Expected Value of Binomial Distribution

$$\begin{aligned}M_X(t) &:= \sum_{y \in \mathbb{S}_X} e^{ty} p_X(y) \\&= \sum_{y=0}^{\infty} e^{ty} \frac{e^{-\lambda} \lambda^y}{y!} \\&= e^{-\lambda} \sum_{y=0}^{\infty} \frac{(\lambda e^t)^y}{y!} \\&= e^{\lambda e^t - \lambda}\end{aligned}\tag{4}$$

Example : The number of customers arriving at a service counter within one-hour period.

Example : The number of typographical errors in a book counted per page.

Example : The number of email messages received at the technical support center daily.

Example : The number of traffic accidents that occur on a specific road during a month.

A Few Examples of Poisson Distribution

Example :

Messages arrive at an electronic message center at random times, with an average of 9 messages per hour.

- 1 What is the probability of receiving exactly five messages during the next hour?
- 2 What is the probability that more than 10 messages will be received within the next two hours?

- 1 The number of messages received in an hour, X is modeled by Poisson distribution with $\lambda = 9$, i.e. $X \sim \text{Poisson}(9)$.

$$P(X = 5) = \frac{9^5 \exp(-9)}{5!}$$

- 2 The number of messages received within a 2-hour period, Y is another Poisson distribution with $Y = (2)(9) = 18$, i.e. $Y \sim \text{Poisson}(18)$. $P(Y > 10) = 1 - P(Y \leq 10) = \dots = 0.9696$

Group Work

- 1 Develop a real life example in which you can easily apply:
 - 1 Group 1: Poisson distribution.
 - 2 Group 2: Binomial distribution.
 - 3 Group 3: Poisson distribution
- 2 In each case, propose two problems which can be solved using the Statistical Calculator.
- 3 Can you propose an idea in which you can mix both distributions? (extra)

Outline

- 1 Binomial Distribution
- 2 Poisson Distribution
- 3 Geometric Distribution
- 4 Negative Binomial Distribution
- 5 Miscellaneous Problems

Geometric Distribution


Geometric Distribution

- 1 Suppose that independent trials, each having a probability π , $0 < \pi < 1$, of being a success, are performed until a success occurs.
- 2 Example: The first head in tossing coin several times.
- 3 Then, Geometric distribution models the number of trials performed until a success occurs.

Definition (Geometric Distribution)

The probability mass function of $Geometric(\pi)$ is given by

$$p(x) = (1 - \pi)^{x-1} \pi \text{ for } x = 1, 2, 3, \dots,$$

 If $X \sim Geometric(\pi)$ then $E(X) = \frac{1}{\pi}$, and $Var(X) = \frac{1-\pi}{\pi^2}$

Expected Value of Binomial Distribution

$$\begin{aligned}M_X(t) &:= \sum_{y \in \mathbb{S}_X} e^{ty} p_X(y) \\&= \sum_{y=1}^{\infty} e^{ty} (1 - \pi)^{y-1} \pi \\&= \pi \sum_{z=0}^{\infty} e^{tz+t} (1 - \pi)^z \\&= pe^t \sum_{z=0}^{\infty} ((1 - \pi)e^t)^z \\&= \frac{pe^t}{1 - (1 - \pi)e^t} \tag{5}\end{aligned}$$

Geometric Distribution: Example

Example : Suppose that the probability of engine malfunction during any one-hour period is $\pi = 0.02$. Find the probability that a given engine will survive two hours.

Geometric Distribution: Example

Example :

Suppose that the probability of engine malfunction during any one-hour period is $\pi = 0.02$. Find the probability that a given engine will survive two hours.

Solution:

Letting Y denote the number of one-hour intervals until the first malfunction, we have

$$\begin{aligned} & P(\text{Survival for Next Two Hours}) \\ = & P(Y \geq 3) \\ = & 1 - P(Y \leq 2) \\ = & 1 - \sum_{y=1}^2 p(y) \\ = & 1 - \{p(1) + p(2)\} \\ = & 1 - 0.02 - 0.98 \times 0.02 \\ = & 0.9604 \end{aligned}$$

Exercise Find the mean and standard deviation of Y .

Outline

- 1 Binomial Distribution
- 2 Poisson Distribution
- 3 Geometric Distribution
- 4 Negative Binomial Distribution
- 5 Miscellaneous Problems


Negative Binomial Distribution

- 1 Suppose that independent trials, each having probability π , $0 < \pi < 1$, of being a success are performed until a total of r successes is accumulated.
- 2 Example: The third head in tossing coin several times.
- 3 Then, Negative Binomial distribution models the number of trials performed until a the r th success occurs.

Definition (Negative Binomial Distribution)

The probability mass function of Negative Binomial RV, denoted by Negative-Binomial(r, π) is given by

$$p(x) = \binom{x-1}{r-1} \pi^{r-1} (1-\pi)^{x-r} \text{ for } x = r+1, r+2, r+3, \dots,$$

 If $X \sim \text{Negative-Binomial}(r, \pi)$ then $E(X) = \frac{r}{\pi}$, and $\text{Var}(X) = \frac{r(1-\pi)}{\pi^2}$

Geometric Distribution: Example

Example :

A machine produces 1% defective parts. Using the statistical calculator, calculate the probability that

- 1 10 parts have to be selected until to get 2 defective parts.
- 2 Between 20 to 25 parts have to be selected to get 2 defective parts.

Geometric Distribution: Example

Example :

A machine produces 1% defective parts. Using the statistical calculator, calculate the probability that

- 1 10 parts have to be selected until to get 2 defective parts.
- 2 Between 20 to 25 parts have to be selected to get 2 defective parts.

Solution:

Letting Y denote the number of

AA

(6)

Exercise Find the mean and standard deviation of Y .

Outline

- 1 Binomial Distribution
- 2 Poisson Distribution
- 3 Geometric Distribution
- 4 Negative Binomial Distribution
- 5 Miscellaneous Problems

Miscellaneous Problems

Example : Approximately 10% of the glass bottles coming off a production line have serious flaws in the glass. If two bottles are randomly selected, find the mean and variance of the number of bottles that have serious flaws.

Example : Cars arrive at a toll both according to a Poisson process with mean 80 cars per hour. If the attendant makes a one-minute phone call, what is the probability that at least 1 car arrives during the call?

Example :

Suppose that a lot of electrical fuses contains 5% defectives. If a sample of 5 fuses is tested, find the probability of observing at least one defective.

Example :

An oil exploration firm is formed with enough capital to finance ten explorations. The probability of a particular exploration being successful is 0.10 . Assume the explorations are independent.

- 1 Find the mean and variance of the number of successful explorations.
- 2 Suppose the firm has a fixed cost of \$20,000 in preparing equipment prior to doing its first exploration. If each successful exploration costs \$30,000 and each unsuccessful exploration costs \$15,000, find the expected total cost to the firm for its ten explorations.

Example :

A food manufacturer uses an extruder (a machine that produces bite-size cookies and snack food) that yields revenue for the firm at a rate of \$200 per hour when in operation. However, the extruder breaks down an average of two times every day it operates. If Y denotes the number of breakdowns per day, the daily revenue generated by the machine is $R = 1600 - 50Y^2$. Find the expected daily revenue for the extruder.

Example :

A particular sale involves four items randomly selected from a large lot that is known to contain 10% defectives. Let Y denote the number of defectives among the 20 sold. The purchaser of the items will return the defectives for repair, and the repair cost is given by $\text{Cost} = 3Y^2 + Y + 2$. Find the expected repair cost.

Example :

In a certain population, it is known that 80% of the individuals have the Rhesus (Rh) factor present in their blood.

- 1 If 5 volunteers are randomly selected from the population, what is the probability that at least one does not have the Rh factor?
- 2 If five volunteers are randomly selected, what is the probability that at most four have the Rh factor?

Example :

Consider rolling a fair dice multiple times until the first 6 appears.

- 1 Find the expected number of throws required to get the first 6.
- 2 What is the probability that more than 8 throws are required to obtain the first 6?

Questions?