

6. (a) Show that the function $f(z) = \frac{1}{z}$ is not analytic at origin although the Cauchy-Riemann equations are satisfied at the origin. (2+2)

(b) Define harmonic conjugate of a function. Show that $u(x,y) = \log(x^2+y^2)$ is a harmonic function and find the corresponding analytic function $f(z)$ in terms of z . (1+2+3]

7. (a) Find the Laurent's expansion of

$$f(z) = \frac{7z-2}{z(z+1)(z-2)}$$

in the region $1 < |z+1| < 3$.

(b) Determine the poles of the function

$$f(z) = \frac{\sin mz + \cos T2}{(z-1)(z+z-6)}$$

and the residue at each pole. Hence evaluate $\oint_C f(z) dz$, where C is the circle $|z|=2.5$. [1+3+2]

$$\begin{aligned} & (\) - \mathbf{y} , v(y) \\ a(o,) = & \int_{in} u(u,o) \, i_{\frac{1}{2}}(a,) \\ y(0,0) & \int u_{o,k} - u_{(0,)} \end{aligned}$$

$$y(o,4) \quad \text{din} \quad v(o,w) \, i_{\frac{1}{2}} v(9,)$$

$$\& \quad \mathbf{JUy} = \neg_v$$

at
rugin
Cauchy-
Renan
ea"s
ane
cah
rd.

$$\begin{aligned} & \text{Though} \\ \text{fo} & \int_{Ln} tce) I \, o) \\ & -CO \\ & o + 1y \quad \text{take} \sim 0_{alan} \\ & \quad \text{Hue} \, ie_{y2mz} \end{aligned}$$

$$(1+in)$$

$$\begin{aligned} & 0+ \\ & \quad 2 \, NIm \\ & \quad 1+M \\ & - \, n1 \end{aligned}$$

are
satisfied.
. . . 'co)
doesnot
eajst,
but
cR
Rguatians

(b) D Han manie ayugate
 A tunction $ua)$ high possess cout pantal
 dariva tiveo $1+ond$ $2dmdo$ ond cohs hieo Hhe Lapa
 $2+U$ $\cdot=0$) is called as a hannonic hune
 $-Ih$ 2 honmaetunetn $u(as$ and $v(,)$ satsly
 Hhe cauchy Teiwan eaMs a V , Lyz-h thaw tay ane
 hasmaie tneHan

$$2a$$

$$-(2).$$

$$)-y()$$

+

+

$\cdot:Tt$ sah's heo Laplaues Equahou
 It ls hanmoic tneHaw.

a4y

Aucmdng to M:1ne Thoso's r lahio, . . 1

te) =

(An) .

(a) 4) . $\frac{z^{-2}}{z(z+)} A + B +$

$A(2+4)^{-2} + BZ(z^2) + <xZ(4)$

$-2 = A_X 1_X(-2)$

A1

a -

2

B - 3

: .

-2

$1 < |z+| < 3$

3

$z+|$

$$i\bullet c) \quad -^3_{-2}+2$$

$$\begin{aligned} & \quad \quad \quad (+)-3 \\ & 1 \quad \quad -3(\quad 4^2 B(-) \\ & \quad \quad \quad (E) t^3)^-(1) \\ & + \quad \quad \quad (-) \quad \quad \quad -(-) \\ & +11_{+1} \quad \quad \quad t.. \end{aligned}$$

+

[An]

$$\begin{aligned} t_z) = & \quad Si \quad Tz'+C \quad zv \quad \quad \quad zt-6 \\ & \quad \quad \quad -)(2-) \\ & \quad \quad \quad \sin z+\cos zY \\ & \quad \quad \quad -)(32)(-3) \\ & 11 \quad \text{Pole} \quad \quad \text{amd}n2 \\ & \quad \quad 2-\text{simple} \quad \text{Prh} \\ & --3-\text{simple pole} \end{aligned}$$

At 1

$$\text{Res}(\text{tee})) = \frac{1}{z-1} \int_{\gamma} f(z) dz$$

+ 2

$$\text{Res}_{z=2} (f(z)) = \lim_{z \rightarrow 2} (z-2) f(z)$$

$$= \lim_{z \rightarrow 2} \frac{\sin z \cos^2 z}{(z-2)^{1/2}}$$

$$= \lim_{z \rightarrow 2} \frac{\sin z \cos^2 z}{(z-2)^{1/2}}$$

af 3

$$\text{Res}_{z=3} (f(z)) = \lim_{z \rightarrow 3} (z-3) f(z)$$

1

$$(4) * (-5)$$