

# FUNDAMENTAL GROUP AND FUNDAMENTAL PROPERTIES

ADITYA KARNATAKI

## 1. INTRODUCTION

Recall that we defined the fundamental group  $\pi_1(X, x_0)$  of a space  $X$  based at  $x_0$  as-

**Definition 1.1.** The set of path-homotopy classes of loops based at  $x_0$ , with the concatenation operation  $*$  is called the *fundamental group* of  $X$  for the base point  $x_0$ , denoted  $\pi_1(X, x_0)$ .

Today we will explore its properties and related concepts.

## 2. FUNDAMENTAL PROPERTIES OF FUNDAMENTAL GROUPS

First we need to figure out how much the group  $\pi_1(X, x_0)$  depends on the basepoint  $x_0$ . To this end, let us observe that if  $x_0$  and  $x_1$  are in the same path component of  $X$ , by definition there exists a path  $\alpha$  from  $x_0$  to  $x_1$ . Then, given any loop  $f$  based at  $x_0$ , we can get a loop based at  $x_1$ , by taking  $\bar{\alpha} * f * \alpha$ , i.e. traversing from  $x_1$  to  $x_0$  first via the reverse path  $\bar{\alpha}$ , looping at  $x_0$  via  $f$ , and then returning back to  $x_1$  via  $\alpha$ . Thus, we get a map

$$\hat{\alpha} : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$$

given by

$$[f] \rightarrow [\bar{\alpha} * f * \alpha] = [\alpha]^{-1} * [f] * [\alpha].$$

(Recall that  $*$  is well-defined on homotopy classes.)

**Proposition 2.1.**  $\hat{\alpha} : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$  is an isomorphism of groups.

*Proof.* This actually follows formally- in the fundamental groupoid we defined earlier, the objects  $x_0$  and  $x_1$  are isomorphic, with the isomorphism given by  $[\alpha]$ . A homework problem last week then would show us (if we have indeed solved it!) that  $\text{Aut}(x_0) \cong \text{Aut}(x_1)$  as groups, and the statement follows. Let us work this out explicitly in this case.

First of all, we need to show that  $\hat{\alpha}$  is a group homomorphism. This is a usual trick. If we have  $a, b \in \pi_1(X, x_0)$ , we have

$$\hat{\alpha}(a * b) = [\alpha]^{-1} * (a * b) * [\alpha] = [\alpha]^{-1} * a * [\alpha] * [\alpha]^{-1} * b * [\alpha]$$

and thus  $\hat{\alpha}(a * b) = \hat{\alpha}(a) * \hat{\alpha}(b)$ .

Let us denote  $\beta = \bar{\alpha}$ . Then we have  $\hat{\beta} : \pi_1(X, x_1) \rightarrow \pi_1(X, x_0)$  which is also a group homomorphism. We claim that  $\hat{\alpha}$  and  $\hat{\beta}$  are inverses of each other. Indeed, for any  $a \in \pi_1(X, x_0)$ , we have

$$\hat{\beta}(\hat{\alpha}(a)) = \hat{\beta}([\alpha]^{-1} * a * [\alpha]) = [\beta]^{-1} * [\alpha]^{-1} * a * [\alpha] * \beta$$

and since  $[\beta] = [\alpha]^{-1}$  in the groupoid, we get that  $\hat{\beta}(\hat{\alpha}(a)) = a$ . Similarly, we have  $\hat{\alpha}(\hat{\beta}(b)) = b$  for any  $b \in \pi_1(X, x_1)$ , so we are done.

**Corollary 2.2.** If  $X$  is path connected, then  $\pi_1(X, x_0)$  is independent of the base point  $x_0$  up to an isomorphism. Sometimes in such a situation, we will denote this object simply as  $\pi_1(X)$ .

**Corollary 2.3.** A loop  $f$  at  $x_0$  induces an automorphism  $\hat{f}$  of  $\pi_1(X, x_0)$ . This is simply of the form  $a \rightarrow [f]^{-1} * a * [f]$ , called an ‘inner automorphism’ (‘conjugation by  $f$ ’).

□

This illustrates how category language can be useful in simplifying proofs and ideas in our minds.

### 3. $\pi_1$ AS A FUNCTOR

Consider the category of *pointed topological spaces*: this has objects  $\{(X, x_0)\}$  ordered pairs of topological spaces  $X$  and the choice of a base point  $x_0 \in X$ . The morphisms are given by continuous maps that take base points to base points-  $f : (X, x_0) \rightarrow (Y, y_0)$  means that  $f : X \rightarrow Y$  is continuous, and  $f(x_0) = y_0$ .

**Definition 3.1.** Any morphism of pointed topological spaces  $h : (X, x_0) \rightarrow (Y, y_0)$  induces a group homomorphism  $h_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$  defined by

$$h_*([f]) := [h \circ f].$$

Note that  $h_*$  is well-defined. If  $f \simeq_p f'$  with homotopy  $F(s, t)$ , then  $h \circ f \simeq_p h \circ f'$  with homotopy  $h \circ F(s, t)$ . It is also a group homomorphism because composition with  $h$  is compactible with concatenation of loops:  $h \circ (f * g) = (h \circ f) * (h \circ g)$ . Thus,  $h_*([f] * [g]) = h_*([f]) * h_*([g])$ .

**Lemma 3.2.**  $\pi_1$  is a functor from the category of pointed topological spaces to Groups.

*Proof.* On objects,  $\pi_1$  takes  $(X, x_0)$  to  $\pi_1(X, x_0)$ . On morphisms, we have  $\pi_1(h) = h_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ . We have to check the two required properties:

- (i) **Composition:** Let  $h : (X, x_0) \rightarrow (Y, y_0)$  and  $k : (Y, y_0) \rightarrow (Z, z_0)$  be two morphisms of pointed topological spaces. Then we have to show that  $\pi_1(k \circ h) = \pi_1(k) \circ \pi_1(h)$ , i.e. for  $(k \circ h)_* = k_* \circ h_*$ . For  $[f] \in \pi_1(X, x_0)$ , we have  $(k \circ h)_*([f]) = [(k \circ h) \circ f] = [k \circ (h \circ f)] = k_*([h \circ f]) = k_*(h_*([f]))$ .
- (ii) **Identity:** We have  $(id_{x_0})_*([f]) = [id_{x_0} \circ f] = [f]$  so that the identity element in the category of pointed topological spaces induces the identity on the group  $\pi_1(X, x_0)$ .

□

Since functors carry isomorphisms to isomorphisms (**exercise!**), we get the following corollary:

**Corollary 3.3.** If  $h : (X, x_0) \rightarrow (Y, y_0)$  is a homeomorphism, then  $h_*$  is an isomorphism of groups.

In fact there are much more general statements of this kind, once we get to homotopy equivalence for topological spaces.