Ans 1:

(i)	*	FIRST	1000	FOLLOW	Conje Buguns	1
	S	fa,63		283		
	A	fa, E3		{a,b} x = x		
	В	₹ 6,83			3111 - 1	1
				esteria a	· Mar e-T	7
	4	1	ei Lucia	8.1	2 - NK	
			, Q.,.		MARK - XI	
(11)		a		b	\$ C-17	
	S	S-A AaAb	Sa BbBa		N-F	
	A	AAA	A-98		Share Canerical	
		A-9 E			man A. A. Santa	
в		B-2 81-1.21	- B - B		of the	
		B-8 B-8		78	1 X . 62 SON	i i
7.	(88.84)	- \$ }	M	- Anna and a second	TON OF K	-
(ii)	No, it	8 not LL(1)			1 2 - 1 . DAL	
	Since t	here are mu	lliple	onthies in	n a single cell	
m.	25	N. S. X		1		
m/	23	A.C. X	n 1's	Am biguo us	TONO C X	1
in /	Also, s	since grammar	not 1	L(1)	40011	
	Also, s	since grammar	not 1	L(1)	70611	
	Also, s	since grammas	not 1	L(1)	40011	
(m/4 - 30)	Also, s	sina gramman	1 ton	L(1)	70611	
(iv) f	Also, s	since gramman it is	mot u	can never	40011	
(iv) f	Also, s	since gramman it is	not 1	L(1)	70611	
(iv) f	Also, s	t of a varia	not l	can never	contain an	
(iv) f	Follow seepsilon	since gramman it is	not l	can never	contain an	
(iv) s	Follow seepsilon	t of a variate because,	hot l	can never	contain an	
(iv) s	Follow seepsilon Since, and Error ca	t of a variate because,	hot l	can never	contain an	