


Image Enhancement:

Frequency representation

Dr. Tushar Sandhan


Introduction

- Signal decomposition




Introduction

- Signal decomposition



Introduction




- Signal decomposition



$$\sin(\omega t + \phi) = \sin(\omega t) \cos(\phi) + \cos(\omega t) \sin(\phi)$$

Introduction



- Signal decomposition





$$\sin(\omega t + \phi) = \sin(\omega t) \cos(\phi) + \cos(\omega t) \sin(\phi)$$

Introduction


- Signal decomposition



$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$





Frequency





$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$

Frequency




$$\sin(\omega t + \phi) = \sin(\omega t) \cos(\phi) + \cos(\omega t) \sin(\phi)$$




Frequency




$$\sin(\omega t + \phi) = \sin(\omega t) \cos(\phi) + \cos(\omega t) \sin(\phi)$$






Frequency




$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$






Frequency




$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$





Frequency



$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$




Frequency



basis: sin()

$$\sin(wt + \phi) = \sin(wt) \cos(\phi) + \cos(wt) \sin(\phi)$$



Frequency

- 2D harmonics
 - $v_o = 0$?

$$f(x, y) = A \cos(u_0 x + v_0 y + \varphi)$$

Frequency

- 2D harmonics

- $v_o = 0$?

- identical sinusoids with

- amplitude A
 - spatial period $P_x = \frac{2\pi}{u_o}$
 - vertical stripes in the image

$$f(x, y) = A \cos(u_o x + v_o y + \varphi)$$

Frequency


- 2D harmonics

- $v_o = 0$?

- identical sinusoids with

- amplitude A
 - spatial period $P_x = \frac{2\pi}{u_0}$
 - vertical stripes in the image

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$



Frequency



- 2D harmonics

- $v_o = 0$?

- identical sinusoids with

- amplitude A
 - spatial period $P_x = \frac{2\pi}{u_0}$
 - vertical stripes in the image

$$f(x, y) = A \cos(u_0 x + v_0 y + \varphi)$$



Frequency

- 2D harmonics

- $v_0 = 0$?



- identical sinusoids with

- amplitude A
 - spatial period $P_x = \frac{2\pi}{u_0}$
 - vertical stripes in the image

- space frequency

- in both x, y directions
 - x direction: $\frac{u_0}{2\pi} (m^{-1})$
 - number of stripes per meter in x direction

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$



Frequency

- 2D harmonics

- $v_o = 0$?


- identical sinusoids with

- amplitude A
 - spatial period $P_x = \frac{2\pi}{u_0}$
 - vertical stripes in the image

- space frequency


- in both x, y directions
 - x direction: $\frac{u_0}{2\pi} (m^{-1})$
 - number of stripes per meter in x direction

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$



- phase


- determines shift d_x of ridges from origin of coordinates
 - d_x is from origin of coordinates as a fraction of harmonic's period
 - $d_x = P_x \frac{\phi}{2\pi} = \frac{\phi}{u_0}$



Frequency

- 2D harmonics
 - $u_0 \neq 0$ & $v_0 \neq 0$?

$$f(x, y) = A \cos(u_0 x + v_0 y + \varphi)$$



Frequency


- 2D harmonics

- $u_0 \neq 0$ & $v_0 \neq 0$?

- stripes become oblique

- They make angle ϑ with x axis
 - $\vartheta = \arctan(\frac{v_0}{u_0})$
 - ratio of both frequencies determines orientation
 - ridges of stripes are characterized by
$$u_0x + v_0y + \varphi = \pm k2\pi$$

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$



Frequency

- 2D harmonics

- $u_0 \neq 0$ & $v_0 \neq 0$?

- stripes become oblique

- They make angle ϑ with x axis

- $\vartheta = \arctan(\frac{v_0}{u_0})$

- ratio of both frequencies determines orientation

- ridges of stripes are characterized by

- $$u_0x + v_0y + \varphi = \pm k2\pi$$

- family of linear equations

- oblique parallel lines


- distance between them = period P of harmonics

- $P = \frac{2\pi}{\sqrt{u_0^2 + v_0^2}}, w_0 = \sqrt{u_0^2 + v_0^2}$

- shifting of stripes by distance d wrt origin & \perp to ridges

- $d = \frac{P\varphi}{2\pi} = \frac{\varphi}{\omega}$

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$



Frequency



- 2D harmonics

- $u_0 \neq 0$ & $v_0 \neq 0$?

- stripes become oblique





- They make angle ϑ with x axis
 - $\vartheta = \arctan(\frac{v_0}{u_0})$
 - ratio of both frequencies determines orientation
 - ridges of stripes are characterized by
$$u_0x + v_0y + \varphi = \pm k2\pi$$
 - family of linear equations
 - oblique parallel lines
 - distance between them = period P of harmonics
 - $P = \frac{2\pi}{\sqrt{u_0^2 + v_0^2}}$, $w_0 = \sqrt{u_0^2 + v_0^2}$
 - shifting of stripes by distance d wrt origin & \perp to ridges
 - $d = \frac{P\varphi}{2\pi} = \frac{\varphi}{\omega}$

$$f(x, y) = A \cos(u_0x + v_0y + \varphi)$$




Frequency

Time Domain




Frequency Domain




Frequency




○ 1D



○ 2D



Time Domain




Frequency Domain




Frequency

○ 1D




○ 2D





Frequency

○ 1D




○ 2D





Frequency

○ 1D




○ 2D





Frequency

○ 1D




○ 2D





Frequency

○ 1D



○ 2D



Fourier

- Fourier series
 - any periodic function can be approximated with series of harmonics
 - sinusoids are considered as harmonics
 - $f(x)$ with period L

Fourier

- Fourier series
 - any periodic function can be approximated with series of harmonics
 - sinusoids are considered as harmonics
 - $f(x)$ with period L

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

Fourier

- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

Fourier

- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

Fourier

- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

$$\int \sin x \sin(2x) dx = 0$$


Fourier

- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

$$\int \sin x \sin(2x) dx = 0$$



Fourier


- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

$$\int \sin(nx) \cos(mx) dx = 0$$

$$\int \sin x \sin(2x) dx = 0$$



Fourier


- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

$$\int \sin(nx) \cos(mx) dx = 0$$

$$\int \sin x \sin(2x) dx = 0$$



$$\int \sin x \cos x dx = 0$$

Fourier


- Fourier series

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$


$$\int \sin(nx) \sin(mx) dx = 0 \quad (m \neq n)$$

$$\int \sin(nx) \cos(mx) dx = 0$$

$$\int \sin x \sin(2x) dx = 0$$



$$\int \sin x \cos x dx = 0$$



Fourier

- Fourier series
 - orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

Fourier

- Fourier series

- orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int_{-L/2}^{L/2} \cos \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx = \begin{cases} L/2 & \text{for } n = m \\ 0 & \text{for } n \neq m \end{cases}$$

$$\int_{-L/2}^{L/2} \sin \frac{2\pi nx}{L} \sin \frac{2\pi mx}{L} dx = \begin{cases} L/2 & \text{for } n = m \\ 0 & \text{for } n \neq m \end{cases}$$

$$\int_{-L/2}^{L/2} \sin \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx = 0$$

Fourier

- Fourier series
 - orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$\int_{-L/2}^{L/2} \cos \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx = \begin{cases} L/2 & \text{for } n = m \\ 0 & \text{for } n \neq m \end{cases}$$

$$\int_{-L/2}^{L/2} \sin \frac{2\pi nx}{L} dx = 0$$

$$\int_{-L/2}^{L/2} \sin \frac{2\pi nx}{L} \sin \frac{2\pi mx}{L} dx = \begin{cases} L/2 & \text{for } n = m \\ 0 & \text{for } n \neq m \end{cases}$$

$$\int_{-L/2}^{L/2} \cos \frac{2\pi nx}{L} dx = 0$$

$$\int_{-L/2}^{L/2} \sin \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx = 0$$

Fourier

- Fourier series

- equation

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$a_0 = \frac{1}{L} \int_{-L/2}^{L/2} f(x) dx$$

$$a_n = \frac{2}{L} \int_{-L/2}^{L/2} f(x) \cos \frac{2\pi nx}{L} dx$$

$$b_n = \frac{2}{L} \int_{-L/2}^{L/2} f(x) \sin \frac{2\pi nx}{L} dx$$

$$n = 1, 2, \dots$$

Fourier

- Fourier series

- equation

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$a_0 = \frac{1}{L} \int_{-L/2}^{L/2} f(x) dx$$

$$a_n = \frac{2}{L} \int_{-L/2}^{L/2} f(x) \cos \frac{2\pi nx}{L} dx$$

$$b_n = \frac{2}{L} \int_{-L/2}^{L/2} f(x) \sin \frac{2\pi nx}{L} dx$$

$$n = 1, 2, \dots$$

Fourier

- Fourier series

- Complex
- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

Fourier

- Fourier series

- Complex

- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$e^{ix} = \cos x + i \sin x$$

Fourier

- Fourier series

- Complex

- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$e^{ix} = \cos x + i \sin x$$

$$\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix})$$

Fourier

- Fourier series

- Complex

- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$e^{ix} = \cos x + i \sin x$$

$$\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix}) \qquad \sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix}) = -\frac{i}{2} (e^{ix} - e^{-ix})$$

Fourier

- Fourier series

- Complex

- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$e^{ix} = \cos x + i \sin x$$

$$\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix}) \qquad \sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix}) = -\frac{i}{2} (e^{ix} - e^{-ix})$$

$$\begin{aligned} f(x) &= a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right) \\ &= a_0 + \frac{1}{2} \sum_{n=1}^{\infty} a_n (e^{i\frac{2\pi nx}{L}} + e^{-i\frac{2\pi nx}{L}}) - \frac{i}{2} \sum_{n=1}^{\infty} b_n (e^{i\frac{2\pi nx}{L}} - e^{-i\frac{2\pi nx}{L}}) \end{aligned}$$

Fourier

- Fourier series

- Complex

- $e^{i\theta} = ?$

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$e^{ix} = \cos x + i \sin x$$

$$\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix}) \qquad \sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix}) = -\frac{i}{2} (e^{ix} - e^{-ix})$$

$$\begin{aligned} f(x) &= a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right) \\ &= a_0 + \frac{1}{2} \sum_{n=1}^{\infty} a_n (e^{i\frac{2\pi nx}{L}} + e^{-i\frac{2\pi nx}{L}}) - \frac{i}{2} \sum_{n=1}^{\infty} b_n (e^{i\frac{2\pi nx}{L}} - e^{-i\frac{2\pi nx}{L}}) \\ &= a_0 + \frac{1}{2} \sum_{n=1}^{\infty} (a_n - ib_n) e^{i\frac{2\pi nx}{L}} + \frac{1}{2} \sum_{n=1}^{\infty} (a_n + ib_n) e^{-i\frac{2\pi nx}{L}} \end{aligned}$$

Fourier

- Fourier series
 - complex
 - orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = a_0 + \frac{1}{2} \sum_{n=1}^{\infty} (a_n - ib_n) e^{i\frac{2\pi nx}{L}} + \frac{1}{2} \sum_{n=1}^{\infty} (a_n + ib_n) e^{-i\frac{2\pi nx}{L}}$$

Fourier

- Fourier series
 - complex
 - orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = a_0 + \frac{1}{2} \sum_{n=1}^{\infty} (a_n - ib_n) e^{i\frac{2\pi nx}{L}} + \frac{1}{2} \sum_{n=1}^{\infty} (a_n + ib_n) e^{-i\frac{2\pi nx}{L}}$$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

Fourier

- Fourier series
 - complex
 - orthogonality

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = a_0 + \frac{1}{2} \sum_{n=1}^{\infty} (a_n - ib_n) e^{i\frac{2\pi nx}{L}} + \frac{1}{2} \sum_{n=1}^{\infty} (a_n + ib_n) e^{-i\frac{2\pi nx}{L}}$$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$\int_{-\frac{L}{2}}^{\frac{L}{2}} e^{-\frac{i2\pi mx}{L}} e^{\frac{i2\pi nx}{L}} = \begin{cases} L & \text{for } m = n \\ 0 & \text{otherwise} \end{cases}$$

Fourier

- Fourier series

- complex

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

Fourier

- Fourier series

- complex

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

Fourier

- Fourier series

- complex

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right)$$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

Fourier

- Fourier Transform
 - any signal
 - finite discontinuities in finite interval
 - $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

$$L c_n = \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-j\frac{2\pi nx}{L}} dx$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

$$L c_n = \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$L \rightarrow \infty$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

$$L c_n = \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$L \rightarrow \infty$$

Frequency n/L becomes continuous

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

$$L c_n = \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$L \rightarrow \infty$$

Frequency n/L becomes continuous

$$n/L \rightarrow u$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$c_n = \frac{1}{L} \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-i\frac{2\pi nx}{L}} dx, n = 0, \pm 1, \pm 2, \dots$$

$$L c_n = \int_{-\frac{L}{2}}^{\frac{L}{2}} f(x) e^{-j\frac{2\pi nx}{L}} dx \quad \longrightarrow \quad F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx$$

$$L \rightarrow \infty$$

Frequency n/L becomes continuous

$$n/L \rightarrow u$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx = L c_n$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx = L c_n$$

$$n/L \rightarrow u$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx = L c_n$$

$$n/L \rightarrow u$$

$$1/L \rightarrow du$$

Fourier

- Fourier Transform

- any signal
- finite discontinuities in finite interval
- $\int_{-\infty}^{\infty} |f(t)| dt < \infty$

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi nx/L}$$

$$F(u) = \int_{-\infty}^{\infty} f(x) \exp(-j2\pi ux) dx = L c_n$$

$$n/L \rightarrow u$$

$$1/L \rightarrow du$$

$$f(x) = \int_{-\infty}^{\infty} F(u) \exp(j2\pi ux) du$$

Fourier

- Discrete Fourier Transform
 - any digital signal
 - N samples in $[0, L]$
 - Δx sample step in x direction
 - $L = ?$

$$c_n = \frac{1}{L} \int_{-L/2}^{L/2} f(x) e^{-j\frac{2\pi nx}{L}} dx$$
$$= \frac{1}{L} \int_0^L f(x) e^{-j\frac{2\pi nx}{L}} dx$$

Fourier

- Discrete Fourier Transform
 - any digital signal
 - N samples in $[0, L]$
 - Δx sample step in x direction
 - $L = ?$

$$c_n = \frac{1}{L} \int_{-L/2}^{L/2} f(x) e^{-j\frac{2\pi nx}{L}} dx$$
$$= \frac{1}{L} \int_0^L f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$f(0), f(\Delta x), f(2\Delta x), \dots, f((N-1)\Delta x)$$

Fourier

- Discrete Fourier Transform
 - any digital signal
 - N samples in $[0, L]$
 - Δx sample step in x direction
 - $L = ?$

$$c_n = \frac{1}{L} \int_{-L/2}^{L/2} f(x) e^{-j\frac{2\pi nx}{L}} dx$$
$$= \frac{1}{L} \int_0^L f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$f(0), f(\Delta x), f(2\Delta x), \dots, f((N-1)\Delta x)$$

$$f(k) = f(k\Delta x), k = 0, 1, 2, \dots, N-1$$

Fourier

- Discrete Fourier Transform
 - any digital signal
 - N samples in $[0, L]$
 - Δx sample step in x direction
 - $L = ?$

$$c_n = \frac{1}{L} \int_{-L/2}^{L/2} f(x) e^{-j\frac{2\pi nx}{L}} dx$$
$$= \frac{1}{L} \int_0^L f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$f(0), f(\Delta x), f(2\Delta x), \dots, f((N-1)\Delta x)$$

$$f(k) = f(k\Delta x), k = 0, 1, 2, \dots, N-1 \qquad f(x) = f(k)$$

Fourier

- Discrete Fourier Transform

- any digital signal
- N samples in $[0, L]$
- Δx sample step in x direction
- $L = ?$

$$c_n = \frac{1}{L} \int_{-L/2}^{L/2} f(x) e^{-j\frac{2\pi nx}{L}} dx$$
$$= \frac{1}{L} \int_0^L f(x) e^{-j\frac{2\pi nx}{L}} dx$$

$$f(0), f(\Delta x), f(2\Delta x), \dots, f((N-1)\Delta x)$$

$$f(k) = f(k\Delta x), k = 0, 1, 2, \dots, N-1 \quad f(x) = f(k)$$

$$c_n = \frac{\Delta x}{N\Delta x} \sum_{k=0}^{N-1} f(k) e^{-j\frac{2\pi nk\Delta x}{N\Delta x}}$$
$$= \frac{1}{N} \sum_{k=0}^{N-1} f(k) e^{-j\frac{2\pi nk}{N}} n = 0, 1, 2, \dots, N-1$$

Fourier

- Discrete Fourier Transform
 - DFT
 - N samples in $[0, L]$
 - $\frac{1}{N}$ outside scaling constant is interchangeably used either in IDFT or in DFT (like below)

Fourier

- Discrete Fourier Transform
 - DFT
 - N samples in $[0, L]$
 - $\frac{1}{N}$ outside scaling constant is interchangeably used either in IDFT or in DFT (like below)

$$F(u) = \frac{1}{N} \sum_{x=0}^{N-1} f(x) \exp(-j2\pi ux/N) \quad u = 0, 1, 2, \dots, N-1$$

Fourier

- Discrete Fourier Transform

- DFT
- N samples in $[0, L]$
- $\frac{1}{N}$ outside scaling constant is interchangeably used either in IDFT or in DFT (like below)

$$F(u) = \frac{1}{N} \sum_{x=0}^{N-1} f(x) \exp(-j2\pi ux/N) \quad u = 0, 1, 2, \dots, N-1$$

$$f(x) = \sum_{u=0}^{N-1} F(u) \exp(j2\pi ux/N) \quad x = 0, 1, 2, \dots, N-1$$

Fourier

- 2D DFT

- MxN samples

$$0 \leq x < M$$
$$0 \leq y < N$$

- image $f(x, y)$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

□ DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

□ DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$u = x = 0, 1, 2, \dots, M - 1$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

□ DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$u = x = 0, 1, 2, \dots, M-1$$

$$v = y = 0, 1, 2, \dots, N-1.$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

□ DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$f(x, y) = \frac{1}{MN} \sum_{u=0}^{M-1} \sum_{v=0}^{N-1} F(u, v) e^{j2\pi(ux/M + vy/N)}$$

$$u = x = 0, 1, 2, \dots, M-1$$

$$v = y = 0, 1, 2, \dots, N-1.$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

- DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

- IDFT:
$$f(x, y) = \frac{1}{MN} \sum_{u=0}^{M-1} \sum_{v=0}^{N-1} F(u, v) e^{j2\pi(ux/M + vy/N)}$$

$$u = x = 0, 1, 2, \dots, M-1$$

$$v = y = 0, 1, 2, \dots, N-1.$$

Fourier

- 2D DFT

- MxN samples

- image $f(x, y)$

- DFT:
$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

- IDFT:
$$f(x, y) = \frac{1}{MN} \sum_{u=0}^{M-1} \sum_{v=0}^{N-1} F(u, v) e^{j2\pi(ux/M + vy/N)}$$

$$u = x = 0, 1, 2, \dots, M-1$$

$$v = y = 0, 1, 2, \dots, N-1.$$

Fourier

- 2D DFT
 - separability

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

Fourier

- 2D DFT
 - separability

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$F(u, v) = \sum_{x=0}^{M-1} \left[\sum_{y=0}^{N-1} f(x, y) e^{-j2\pi vy/N} \right] e^{-j2\pi ux/M}$$

Fourier

- 2D DFT
 - separability

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$F(u, v) = \sum_{x=0}^{M-1} \left[\sum_{y=0}^{N-1} f(x, y) e^{-j2\pi vy/N} \right] e^{-j2\pi ux/M}$$

$$F(u, v) = \sum_{x=0}^{M-1} [F(x, v)] e^{-j2\pi ux/M}$$

Fourier

- 2D DFT

- separability

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$F(u, v) = \sum_{x=0}^{M-1} \left[\sum_{y=0}^{N-1} f(x, y) e^{-j2\pi vy/N} \right] e^{-j2\pi ux/M}$$

$$F(u, v) = \sum_{x=0}^{M-1} [F(x, v)] e^{-j2\pi ux/M}$$

$$F(u, v) = FT_x \{ FT_y [f(x, y)] \}$$

Fourier

- 2D DFT

- separability

$$F(u, v) = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} f(x, y) e^{-j2\pi(ux/M + vy/N)}$$

$$F(u, v) = \sum_{x=0}^{M-1} \left[\sum_{y=0}^{N-1} f(x, y) e^{-j2\pi vy/N} \right] e^{-j2\pi ux/M}$$

$$F(u, v) = \sum_{x=0}^{M-1} [F(x, v)] e^{-j2\pi ux/M}$$

$$F(u, v) = FT_x \{ FT_y [f(x, y)] \}$$

FT_x and FT_y are the 1D FTs on row and column, respectively.


Fourier

- 2D DFT
 - image
 - use grayscale image for DFT



Fourier


- 2D DFT
 - image
 - use grayscale image for DFT



Fourier

- 2D DFT
 - image
 - use grayscale image for DFT

$f(x, y)$



Fourier


- 2D DFT

- image
- use grayscale image for DFT

$f(x, y)$



Enhanced($|F(u, v)|$)



Fourier

- 2D DFT
 - image



Fourier

- 2D DFT

- image

$f(x, y)$




Fourier

- 2D DFT

- image

$f(x, y)$



Fourier


- 2D DFT

- image

$f(x, y)$



$|F(u, v)|$



Fourier



- 2D DFT

- image

$f(x, y)$



$|F(u, v)|$



Fourier


- 2D DFT

- image


$f(x, y)$



$|F(u, v)|$



$\angle F(u, v)$



Conclusion

- Frequency representation
- Fourier
 - Series
 - Transform
 - 2D DFT

Conclusion

- Frequency representation
- Fourier
 - Series
 - Transform
 - 2D DFT



Conclusion

- Frequency representation
- Fourier
 - Series
 - Transform
 - 2D DFT



"Four years at IIT
transform ($f_F^* T$)
life phenomenally"

-TS

Conclusion

- Frequency representation
- Fourier
 - Series
 - Transform
 - 2D DFT



"Four years at IIT
transform ($f_F^* T$)
life phenomenally"

-TS



Conclusion

- Frequency representation
- Fourier
 - Series
 - Transform
 - 2D DFT



"Four years at IIT
transform ($f_F^* T$)
life phenomenally"

-TS



Lagrange, Laplace, Monge