

# COMP 680

## Statistics for Computing and Data Science

### Week 13: Beyond Linear Models

Su Chen, Assistant Teaching Professor,  
Rice D2K Lab

# Outline

- 1 Beyond Linearity
- 2 Regularization
- 3 Splines and GAM
- 4 Nonparametric Models
- 5 Code Demo

# The Truth is probably NOT Linear

- But often the linearity assumption is good enough
  - “all models are wrong, some are useful.”

# The Truth is probably NOT Linear

- But often the linearity assumption is good enough
  - “all models are wrong, some are useful.”
- When linearity is clearly not enough:
  - polynomial regression
  - spline models
  - generalized additive models (GAM)
  - local regression
  - fully non-parametric models

# Outline

- ① Beyond Linearity
- ② Regularization
- ③ Splines and GAM
- ④ Nonparametric Models
- ⑤ Code Demo

# Motivation

- High dimensional data
  - but OLS solution is **not available** for  $p > n$

# Motivation

- High dimensional data
  - but OLS solution is **not available for  $p > n$**
- Prediction accuracy:
  - when  $p$  is large, need to control the variance
  - trade bias with variance to decrease MSE

# Motivation

- High dimensional data
  - but OLS solution is **not available for  $p > n$**
- Prediction accuracy:
  - when  $p$  is large, need to control the variance
  - trade bias with variance to decrease MSE
- Model Interpretability:
  - penalize large models and large slopes
  - automatically perform variable selection



# Shrinkage Estimate

- We fit a model involving all  $p$  covariates
- But the estimated slopes are shrunken towards 0 relative to OLS.
- This shrinkage is known as **regularization**
  - penalize “large”  $\beta$  by shrinking them - reduce variance
  - shrink some  $\beta$  to exactly 0 - variable selection

# Ridge Regression

- Recall that OLS estimates  $\hat{\beta}^{OLS}$  using the values that minimize

$$\text{RSS} = \sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 .$$

# Ridge Regression

- Recall that OLS estimates  $\hat{\beta}^{OLS}$  using the values that minimize

$$\text{RSS} = \sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2.$$

- In contrast, the ridge regression coefficient estimates  $\hat{\beta}^R$  are the values that minimize

$$\sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p \beta_j^2 = \text{RSS} + \lambda \|\beta\|_{L_2}^2,$$

where  $\lambda \geq 0$  is a **tuning parameter**, to be determined separately

# Ridge Regression

- As with least squares, ridge regression seeks coefficient estimates that fit the data well, by making the RSS small.

# Ridge Regression

- As with least squares, ridge regression seeks coefficient estimates that fit the data well, by making the RSS small.
- However, the second term,  $\lambda \|\beta\|_{L_2}^2$ , called a **shrinkage penalty**, is small when  $\beta_1, \dots, \beta_p$  are close to zero, and so it has the effect of shrinking the estimates of  $\beta_j$  towards zero.



# Ridge Regression

- As with least squares, ridge regression seeks coefficient estimates that fit the data well, by making the RSS small.
- However, the second term,  $\lambda \|\boldsymbol{\beta}\|_{L_2}^2$ , called a **shrinkage penalty**, is small when  $\beta_1, \dots, \beta_p$  are close to zero, and so it has the effect of shrinking the estimates of  $\beta_j$  towards zero.
- The tuning parameter  $\lambda$  serves to control the relative impact of these two terms on the regression coefficient estimates.

# Ridge Regression

- As with least squares, ridge regression seeks coefficient estimates that fit the data well, by making the RSS small.
- However, the second term,  $\lambda \|\boldsymbol{\beta}\|_{L_2}^2$ , called a **shrinkage penalty**, is small when  $\beta_1, \dots, \beta_p$  are close to zero, and so it has the effect of shrinking the estimates of  $\beta_j$  towards zero.
- The tuning parameter  $\lambda$  serves to control the relative impact of these two terms on the regression coefficient estimates.
- Selecting a good value for  $\lambda$  is critical!

# Ridge Solution Path



Example from ISLR



## Details of Previous Figure

- In the left-hand panel, each curve corresponds to the ridge regression coefficient estimate for one of the ten variables, plotted as a function of  $\lambda$ .

## Details of Previous Figure

- In the left-hand panel, each curve corresponds to the ridge regression coefficient estimate for one of the ten variables, plotted as a function of  $\lambda$ .
- The right-hand panel displays the same ridge coefficient estimates as the left-hand panel, but instead of displaying  $\lambda$  on the x-axis, we now display  $\|\hat{\beta}_\lambda^R\|_2 / \|\hat{\beta}\|_2$ , where  $\hat{\beta}$  denotes the vector of least squares coefficient estimates.

## Details of Previous Figure

- In the left-hand panel, each curve corresponds to the ridge regression coefficient estimate for one of the ten variables, plotted as a function of  $\lambda$ .
- The right-hand panel displays the same ridge coefficient estimates as the left-hand panel, but instead of displaying  $\lambda$  on the x-axis, we now display  $\|\hat{\beta}_\lambda^R\|_2 / \|\hat{\beta}\|_2$ , where  $\hat{\beta}$  denotes the vector of least squares coefficient estimates.
- The notation  $\|\beta\|_2$  denotes the  $L^2$  norm of a vector, and is defined as  $\|\beta\|_2 = \sqrt{\sum_{j=1}^p \beta_j^2}$ .

# Lasso Regression

- Disadvantage of Ridge regression
  - shrink slopes towards 0 but not exactly 0
  - final model still include all  $p$  covariates
  - does not perform variable selection

# Lasso Regression

- Disadvantage of Ridge regression
  - shrink slopes towards 0 but not exactly 0
  - final model still include all  $p$  covariates
  - does not perform variable selection
- The **Lasso** coefficients,  $\hat{\beta}_\lambda^L$ , minimize the quantity

$$\sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p |\beta_j| = \text{RSS} + \lambda \|\beta\|_{L1}.$$

# Lasso Regression

- Disadvantage of Ridge regression
  - shrink slopes towards 0 but not exactly 0
  - final model still include all  $p$  covariates
  - does not perform variable selection
- The **Lasso** coefficients,  $\hat{\beta}_\lambda^L$ , minimize the quantity

$$\sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p |\beta_j| = \text{RSS} + \lambda \|\beta\|_{L1}.$$

- **Lasso uses an L1 penalty while Ridge uses an L2 penalty.**

# Lasso Regression

- As with ridge regression, the Lasso also shrinks the coefficient estimates towards zero.

# Lasso Regression

- As with ridge regression, the Lasso also shrinks the coefficient estimates towards zero.
- However, the L1 penalty has the effect of forcing some of the coefficient estimates to be exactly equal to zero when the tuning parameter  $\lambda$  is sufficiently large.



# Lasso Regression

- As with ridge regression, the Lasso also shrinks the coefficient estimates towards zero.
- However, the L1 penalty has the effect of forcing some of the coefficient estimates to be exactly equal to zero when the tuning parameter  $\lambda$  is sufficiently large.
- Therefore, Lasso performs **variable selection** automatically when estimating the model



# Lasso Regression

- As with ridge regression, the Lasso also shrinks the coefficient estimates towards zero.
- However, the L1 penalty has the effect of forcing some of the coefficient estimates to be exactly equal to zero when the tuning parameter  $\lambda$  is sufficiently large.
- Therefore, Lasso performs **variable selection** automatically when estimating the model
  - yields **sparse** models — that is, models that involve only a subset of the variables

# Lasso Regression

- As with ridge regression, the Lasso also shrinks the coefficient estimates towards zero.
- However, the L1 penalty has the effect of forcing some of the coefficient estimates to be exactly equal to zero when the tuning parameter  $\lambda$  is sufficiently large.
- Therefore, Lasso performs **variable selection** automatically when estimating the model
  - yields **sparse** models — that is, models that involve only a subset of the variables
- Same as in ridge regression, selecting a good value of  $\lambda$  for the lasso is critical!

# Lasso Solution Path



Example from [ISLR](#)

# The Variable Selection Property of the Lasso

Why is it that the lasso, unlike ridge regression, results in coefficient estimates that are exactly equal to zero?

# The Variable Selection Property of the Lasso

Why is it that the lasso, unlike ridge regression, results in coefficient estimates that are exactly equal to zero?

$$\hat{\beta}^R = \arg \min \left[ \sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p \beta_j^2 \right]$$

this is **equivalent** to an optimization with constrain:

$$\text{minimize}_{\beta} \sum_{i=1}^n \left( y_i - \sum_{j=1}^p \beta_j x_{ij} \right)^2 \quad \text{subject to} \quad \sum_{j=1}^p \beta_j^2 \leq s$$

i.e. there is a **1-1 correspondence of  $\lambda$  and  $s$**  that produce the same solution!

# The Variable Selection Property of the Lasso



Use two covariates as example for visualization purpose:

$$\text{minimize}_{\beta} \sum_{i=1}^n (y_i - \beta_1 x_{i1} - \beta_2 x_{i2})^2$$

Ridge penalty:  $\lambda(\beta_1^2 + \beta_2^2) \implies \min \text{RSS subject to } \beta_1^2 + \beta_2^2 \leq s$

Lasso penalty:  $\lambda(|\beta_1| + |\beta_2|) \implies \min \text{RSS subject to } |\beta_1| + |\beta_2| \leq s$

# The Intuition





# What about Inference

- Notice the Ridge and Lasso solutions are no longer MLE
  - no more sampling distributions of  $\hat{\beta}^R$  or  $\hat{\beta}^L$
  - no more p-values and CI ???

# What about Inference

- Notice the Ridge and Lasso solutions are no longer MLE
  - no more sampling distributions of  $\hat{\beta}^R$  or  $\hat{\beta}^L$
  - no more p-values and CI ???
- Lasso selected variables have nothing to do with statistical significance!
  - in practice of course there is still some agreement
  - cares more about prediction than what is the “true” model

# What about Inference


- Notice the Ridge and Lasso solutions are no longer MLE
  - no more sampling distributions of  $\hat{\beta}^R$  or  $\hat{\beta}^L$
  - no more p-values and CI ???
- Lasso selected variables have nothing to do with statistical significance!
  - in practice of course there is still some agreement
  - cares more about prediction than what is the “true” model
- Post selection inference is an active area of research:
  - can we recover true signals if we have infinity amount of data
  - what if number of variables increases with sample size

# Connection to Bayesian Methods

Solutions are posterior mode with corresponding prior on  $\beta$ :


- Ridge prior: Gaussian

$$g(\beta_j) = \frac{1}{\sqrt{2\pi\tau}} \exp\left(-\frac{\beta_j^2}{2\tau^2}\right)$$



- Lasso prior: Laplace

$$g(\beta_j) = \frac{1}{2b} \exp\left(-\frac{|\beta_j|}{b}\right)$$



# In Practice

- Standardize covariates!
  - why?
  - most software does automatically

# In Practice

- Standardize covariates!
  - why?
  - most software does automatically
- How to choose  $\lambda$ ?
  - hyper-parameter tuning in ML
  - choose a grid of  $\lambda$  values, and compute validation/cross-validation error for each value of  $\lambda$ - choose the smallest validation error

# In Practice



- Standardize covariates!
  - why?
  - most software does automatically
- How to choose  $\lambda$ ?
  - hyper-parameter tuning in ML
  - choose a grid of  $\lambda$  values, and compute validation/cross-validation error for each value of  $\lambda$ - choose the smallest validation error
- Ridge or Lasso?
  - neither will universally dominate the other
  - Lasso wins when true regression function is sparse!

# In Practice

- Standardize covariates!
  - why?
  - most software does automatically
- How to choose  $\lambda$ ?
  - hyper-parameter tuning in ML
  - choose a grid of  $\lambda$  values, and compute validation/cross-validation error for each value of  $\lambda$ - choose the smallest validation error
- Ridge or Lasso?
  - neither will universally dominate the other
  - Lasso wins when true regression function is sparse!
- Can be applied to any model fitting using optimization:
  - GLM, spline models, GAM...
  - all parametric ML models...



# Lasso with CV



Example from [ISLR](#)

# Outline

- ① Beyond Linearity
- ② Regularization
- ③ Splines and GAM
- ④ Nonparametric Models
- ⑤ Code Demo

# Polynomial Regression

- Introduce nonlinearity by adding polynomial terms

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + \cdots + \epsilon_i$$

# Polynomial Regression

- Introduce nonlinearity by adding polynomial terms

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + \cdots + \epsilon_i$$

- Degree of the polynomial controls the flexibility of the model
  - degree of freedom  $\approx$  number of parameters

# Polynomial Regression

- Introduce nonlinearity by adding polynomial terms

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + \cdots + \epsilon_i$$

- Degree of the polynomial controls the flexibility of the model
  - degree of freedom  $\approx$  number of parameters
- How to select the degree of polynomial?
  - Stat approach - ANOVA test to compare nested models
  - ML approach - treat as a tuning parameter

# Piecewise Polynomials

- Instead of a single polynomial in  $X$  over its whole domain, we can rather use different polynomials in regions defined by **knots**

$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \epsilon_i, & \text{if } x_i < c \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \epsilon_i, & \text{if } x_i \geq c \end{cases}$$

# Piecewise Polynomials

- Instead of a single polynomial in  $X$  over its whole domain, we can rather use different polynomials in regions defined by **knots**

$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \epsilon_i, & \text{if } x_i < c \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \epsilon_i, & \text{if } x_i \geq c \end{cases}$$

- Better to add constraints to the polynomials, e.g. continuity.

# Piecewise Polynomials


- Instead of a single polynomial in  $X$  over its whole domain, we can rather use different polynomials in regions defined by **knots**

$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \epsilon_i, & \text{if } x_i < c \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \epsilon_i, & \text{if } x_i \geq c \end{cases}$$


- Better to add constraints to the polynomials, e.g. continuity.
- **Splines** have the “maximum” amount of continuity
  - piece together local polynomials smoothly




Piecewise Cubic




Continuous Piecewise Cubic



Cubic Spline



Linear Spline



# Cubic Splines

- A cubic spline with knots at  $\xi_k$ ,  $k = 1, 2, \dots, K$  is a piecewise cubic polynomial with continuous derivatives up to order 2 at each knot.

# Cubic Splines

- A cubic spline with knots at  $\xi_k$ ,  $k = 1, 2, \dots, K$  is a piecewise cubic polynomial with continuous derivatives up to order 2 at each knot.
- We can represent this model with truncated power basis functions:

$$y_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \dots + \beta_{K+3} b_{K+3}(x_i) + \epsilon_i$$

where the  $b_k$  are [basis functions](#)

# Cubic Spline Basis

- The  $b_k$  are **basis functions**

$$b_1(x_i) = x_i$$

$$b_2(x_i) = x_i^2$$

$$b_3(x_i) = x_i^3$$

$$b_{k+3}(x_i) = (x_i - \xi_k)_+^3, \quad k = 1, 2, \dots, K$$

# Cubic Spline Basis

- The  $b_k$  are **basis functions**

$$b_1(x_i) = x_i$$



$$b_2(x_i) = x_i^2$$

$$b_3(x_i) = x_i^3$$

$$b_{k+3}(x_i) = (x_i - \xi_k)_+^3, \quad k = 1, 2, \dots, K$$

- where

$$(x_i - \xi_k)_+^3 = \begin{cases} (x_i - \xi_k)^3 & \text{if } x_i > \xi_k \\ 0 & \text{otherwise} \end{cases}$$



# More Constrains

- Natural Cubic Splines
  - extrapolates linearly beyond the boundary knots
  - cubic spline with  $K$  knots =  $K + 4$  df
  - natural spline with  $K$  knots =  $K$  df

# More Constrains

- Natural Cubic Splines
  - extrapolates linearly beyond the boundary knots
  - cubic spline with  $K$  knots =  $K + 4$  df
  - natural spline with  $K$  knots =  $K$  df
- How many knots and where?
  - choose number of knots = choose flexibility
  - some ad-hoc: fixed intervals, percentiles of  $X$ , etc.



# More Constrains

- Natural Cubic Splines
  - extrapolates linearly beyond the boundary knots
  - cubic spline with  $K$  knots =  $K + 4$  df
  - natural spline with  $K$  knots =  $K$  df
- How many knots and where?
  - choose number of knots = choose flexibility
  - some ad-hoc: fixed intervals, percentiles of  $X$ , etc.
- Smoothing splines
  - choose knot at each data point  $x_i$
  - add smoothing penalty to control df

# Smoothing Splines

- Consider a regression model  $y_i = g(x_i) + \epsilon_i$  where we solve for:

$$\hat{g} = \arg \min_{g \in \mathbb{S}} \left( \sum_{i=1}^n (y_i - g(x_i))^2 + \lambda \int g''(t)^2 dt \right)$$

# Smoothing Splines

- Consider a regression model  $y_i = g(x_i) + \epsilon_i$  where we solve for:

$$\hat{g} = \arg \min_{g \in \mathbb{S}} \left( \sum_{i=1}^n (y_i - g(x_i))^2 + \lambda \int g''(t)^2 dt \right)$$

- Minimize RSS tries to make  $g(x)$  match the data at each  $x_i$ 
  - notice  $g$  is restricted to be smooth

# Smoothing Splines

- Consider a regression model  $y_i = g(x_i) + \epsilon_i$  where we solve for:

$$\hat{g} = \arg \min_{g \in \mathbb{S}} \left( \sum_{i=1}^n (y_i - g(x_i))^2 + \lambda \int g''(t)^2 dt \right)$$

- Minimize RSS tries to make  $g(x)$  match the data at each  $x_i$ 
  - notice  $g$  is restricted to be smooth
- Add a **roughness penalty** to control how wiggly  $g(x)$  is

# Smoothing Splines

- Consider a regression model  $y_i = g(x_i) + \epsilon_i$  where we solve for:

$$\hat{g} = \arg \min_{g \in \mathbb{S}} \left( \sum_{i=1}^n (y_i - g(x_i))^2 + \lambda \int g''(t)^2 dt \right)$$

- Minimize RSS tries to make  $g(x)$  match the data at each  $x_i$ 
  - notice  $g$  is restricted to be smooth
- Add a **roughness penalty** to control how wiggly  $g(x)$  is
  - $\lambda \rightarrow 0$ ?

# Smoothing Splines

- Consider a regression model  $y_i = g(x_i) + \epsilon_i$  where we solve for:

$$\hat{g} = \arg \min_{g \in \mathbb{S}} \left( \sum_{i=1}^n (y_i - g(x_i))^2 + \lambda \int g''(t)^2 dt \right)$$

- Minimize RSS tries to make  $g(x)$  match the data at each  $x_i$ 
  - notice  $g$  is restricted to be smooth
- Add a **roughness penalty** to control how wiggly  $g(x)$  is
  - $\lambda \rightarrow 0$ ?
  - $\lambda \rightarrow \infty$ ?

# Smoothing Splines

- The solution is a natural cubic spline, with a knot at every unique value of  $x_i$

# Smoothing Splines


- The solution is a natural cubic spline, with a knot at every unique value of  $x_i$
- The roughness penalty controls the df!
  - $df \ll$  number of knots
  - avoid the knot-selection issue
  - a single  $\lambda$  to be chosen



# Smoothing Splines

- The solution is a natural cubic spline, with a knot at every unique value of  $x_i$
- The roughness penalty controls the df!
  - $df \ll$  number of knots
  - avoid the knot-selection issue
  - a single  $\lambda$  to be chosen
- Most software can specify df rather than  $\lambda$ 
  - ML approach: treat  $\lambda$  as a tuning parameter
  - same regularization idea

## Smoothing Spline



# Generalized Additive Models (GAM)

- Allow nonlinearity in GLM but still **additive** in covariates:

$$g(\mathbb{E}[Y|X]) = \beta_0 + f_1(X_{i1}) + f_2(X_{i2}) + \cdots + f_p(X_{ip})$$

# Generalized Additive Models (GAM)

- Allow nonlinearity in GLM but still **additive** in covariates:

$$g(\mathbb{E}[Y|X]) = \beta_0 + f_1(X_{i1}) + f_2(X_{i2}) + \cdots + f_p(X_{ip})$$

- $f_j$  can be linear, polynomial, spline...
  - nonlinear terms only for numerical covariates
- Coefficients no longer interpretable
  - fitted function values are (partial plot)

# Generalized Additive Models (GAM)

- Allow nonlinearity in GLM but still **additive** in covariates:

$$g(\mathbb{E}[Y|X]) = \beta_0 + f_1(X_{i1}) + f_2(X_{i2}) + \cdots + f_p(X_{ip})$$

- $f_j$  can be linear, polynomial, spline...
  - nonlinear terms only for numerical covariates
- Coefficients no longer interpretable
  - fitted function values are (partial plot)
- Relax additive assumption?
  - bivariate smoothers
  - low-order interactions

# Outline

- ① Beyond Linearity
- ② Regularization
- ③ Splines and GAM
- ④ Nonparametric Models
- ⑤ Code Demo

# Kernel Density Estimate

- Nonparametric method to estimate a density function:

$$\hat{f}(x) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h} K\left(\frac{x - X_i}{h}\right)$$

- $K$  is the kernel function: uniform, triangular, Gaussian...
- $h$  is the bandwidth

# Kernel Density Estimate

- Nonparametric method to estimate a density function:

$$\hat{f}(x) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h} K\left(\frac{x - X_i}{h}\right)$$

- $K$  is the kernel function: uniform, triangular, Gaussian...
  - $h$  is the bandwidth
- “Smoothed out” histogram
  - converges faster



# Kernel Density Estimate

- Nonparametric method to estimate a density function:

$$\hat{f}(x) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h} K\left(\frac{x - X_i}{h}\right)$$

- $K$  is the kernel function: uniform, triangular, Gaussian...
  - $h$  is the bandwidth
- “Smoothed out” histogram
  - converges faster
- In practice, need to choose a kernel and a bandwidth
  - some asymptotic guideline
  - software default choice

# Kernel Regression

- Want to estimate the regression function as the conditional mean:

$$f(x) = \mathbb{E}[Y|X = x]$$

with observed data  $(x_i, y_i)$  for  $i = 1, 2, \dots, n$ .

# Kernel Regression

- Want to estimate the regression function as the conditional mean:

$$f(x) = \mathbb{E}[Y|X = x]$$

with observed data  $(x_i, y_i)$  for  $i = 1, 2, \dots, n$ .

- Kernel regression with kernel  $K$  and bandwidth  $h$ :

$$\hat{f}(x) = \sum_{i=1}^n \omega_i(x) y_i, \quad \text{where} \quad \omega_i(x) = \frac{K\left(\frac{x-x_i}{h}\right)}{\sum_{k=1}^n K\left(\frac{x-x_k}{h}\right)}$$

# Kernel Regression

- Want to estimate the regression function as the conditional mean:

$$f(x) = \mathbb{E}[Y|X = x]$$

with observed data  $(x_i, y_i)$  for  $i = 1, 2, \dots, n$ .

- Kernel regression with kernel  $K$  and bandwidth  $h$ :

$$\hat{f}(x) = \sum_{i=1}^n \omega_i(x) y_i, \quad \text{where} \quad \omega_i(x) = \frac{K\left(\frac{x-x_i}{h}\right)}{\sum_{k=1}^n K\left(\frac{x-x_k}{h}\right)}$$

- $K$  and  $h$  play the similar role in KDE


# Local Regression

- Locally Weighted Scatterplot Smoothing
  - fit linear regression locally by weighted least squares
  - (weighted) nearest neighbor regression as a special case

# Local Regression

- Locally Weighted Scatterplot Smoothing
  - fit linear regression locally by weighted least squares
  - (weighted) nearest neighbor regression as a special case

Local Regression




Example from [ISLR](#)

# Summary

- With GAM in your statistics toolbox, you are able to:
  - model any type of **response variable** in the GLM family:
    - continuous: Gaussian, Gamma, Beta
    - counts: Poisson, Negative Binomial
    - binary: Binomial
    - categorical: Multinomial
  - include both numerical and categorical **predictors**
  - include a **mix of linear and nonlinear** effects
    - how do you decide?
  - include interaction terms to relax additive assumption
    - interpretation is key
  - apply regularization with Ridge or Lasso penalty

# Which means...

You are now officially a supermodeler. Bravo!!!





# Outline

- ① Beyond Linearity
- ② Regularization
- ③ Splines and GAM
- ④ Nonparametric Models
- ⑤ Code Demo