

# COMP 680

## Statistics for Computing and Data Science

### Week 1: Probability and Random Variable I

Su Chen, Assistant Teaching Professor,  
Rice D2K Lab

# Outline

- ① Course Summary
- ② Basic Probability Theory
- ③ Random Variables
- ④ Bivariate Distributions
- ⑤ Code Demo

# Your Instructor

- Su Chen
- Email: [Su.Chen@rice.edu](mailto:Su.Chen@rice.edu)
- Office: Duncan 2051
- About me

# Administrative Stuff

You can find everything on the [Canvas website](#) for this course

- The syllabus
- Weekly slides, code demo and quiz
- Bi-weekly homework assignments
- Announcements and emails...

# Prerequisites

- College level calculus: single & multivariate
- Basics of linear algebra: vectors and matrices
- Comfortable with mathematical reasoning...
- Basic programming skills (or not afraid to learn :)

# Learning Objectives

- Identify and calculate appropriate summary and inferential statistics based on data.

# Learning Objectives

- Identify and calculate appropriate summary and inferential statistics based on data.
- Gain fluency in basic programming skills in Python with a focus on simulation-based inference and statistical modeling.

# Learning Objectives

- Identify and calculate appropriate summary and inferential statistics based on data.
- Gain fluency in basic programming skills in Python with a focus on simulation-based inference and statistical modeling.
- Use applied statistical knowledge to analyze real-world data, test hypotheses, build regression models, and make scientific inference.



# Learning Objectives

- Identify and calculate appropriate summary and inferential statistics based on data.
- Gain fluency in basic programming skills in Python with a focus on simulation-based inference and statistical modeling.
- Use applied statistical knowledge to analyze real-world data, test hypotheses, build regression models, and make scientific inference.
- Interpret inferences and modeling results in real-world contexts and communicate the findings effectively.

# Course Work

- Weekly classes schedule: MW 2-3:15pm
  - Attendance
  - Participation
  - Feedback

# Course Work

- Weekly classes schedule: MW 2-3:15pm
  - Attendance
  - Participation
  - Feedback
- Major assignments:
  - Bi-weekly homework assignments
  - Weekly quizzes
  - Midterm in-class exam
  - Final takehome exam

# Grades

- Assignment: 45%
  - homework: 30%
  - quizzes: 15%

# Grades

- Assignment: 45%
  - homework: 30%
  - quizzes: 15%
- Exams: 50%
  - midterm: 20%
  - final: 30%

# Grades

- Assignment: 45%
  - homework: 30%
  - quizzes: 15%
- Exams: 50%
  - midterm: 20%
  - final: 30%
- Other: 5%
  - class participation and attendance: 3%
  - course exit survey: 2%

# Etc.

- Disability accommodations.
- Tentative schedule.
- Syllabus is subject to change with reasonable advance notice by the instructor.

# What is Data Science?

- A definition: draw useful information and actionable knowledge from data using computation: so that we can better understand the world and solve problems!



# What is Data Science?

- A definition: draw useful information and actionable knowledge from data using computation: so that we can better understand the world and solve problems!
- math + stats + computing + **domain knowledge**

# What is Data Science?

- A definition: draw useful information and actionable knowledge from data using computation: so that we can better understand the world and solve problems!
- math + stats + computing + **domain knowledge**
- [A Brief History of Data Science](#)

# What is Data Science?

- A definition: draw useful information and actionable knowledge from data using computation: so that we can better understand the world and solve problems!
- math + stats + computing + **domain knowledge**
- [A Brief History of Data Science](#)
- Rooted in Statistics, powered by Computer Science, driven by application, interdisciplinary in nature
  - big data, cloud computing, data mining
  - machine learning/deep learning, artificial intelligence ...

# What is Statistics?



Statistics are like  
bikinis. What  
they reveal is  
suggestive but  
what they  
conceal is vital.

-Aaron Levenstein



# What is Statistics??

- Origin of the word “statistics”
  - information useful to the state
  - early human statistical and census methods

# What is Statistics??

- Origin of the word “statistics”
  - information useful to the state
  - early human statistical and census methods
- [Wiki: history of statistics](#)
  - [Seven Pillars of Statistical Wisdom](#)

# What is Statistics??

- Origin of the word “statistics”
  - information useful to the state
  - early human statistical and census methods
- [Wiki: history of statistics](#)
  - [Seven Pillars of Statistical Wisdom](#)
- Statistics as a discipline:
  - procedures and techniques used to collect, process, and analyze the numerical data to make inferences and to reach appropriate decisions in situations of uncertainty

# What is Statistics??

- Origin of the word “statistics”
  - information useful to the state
  - early human statistical and census methods
- [Wiki: history of statistics](#)
  - [Seven Pillars of Statistical Wisdom](#)
- Statistics as a discipline:
  - procedures and techniques used to collect, process, and analyze the numerical data to make inferences and to reach appropriate decisions in situations of uncertainty
- Statistics as problem solving: harnessing the power of uncertainty
  - descriptive / exploratory
  - predictive / decision-making
  - inferential / causal



# Some Misconceptions

- Stats require hardcore math

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data
  - bigger data  $\neq$  better data

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data
  - bigger data  $\neq$  better data
- Stats provide unambiguous results & unique solution

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data
  - bigger data  $\neq$  better data
- Stats provide unambiguous results & unique solution
  - “I report, you decide.”

# Some Misconceptions


- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data
  - bigger data  $\neq$  better data
- Stats provide unambiguous results & unique solution
  - “I report, you decide.”
- “Lies, damn lies, and statistics.”

# Some Misconceptions

- Stats require hardcore math
  - applied stats is about reason with data
- Stats need lot of data
  - bigger data  $\neq$  better data
- Stats provide unambiguous results & unique solution
  - “I report, you decide.”
- “Lies, damn lies, and statistics.”
  - statistics don't lie, people do.



# Statistics vs. Data Science



# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA

# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA
- Is data science just applied stats???

# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA
- Is data science just applied stats???
  - origin of the field

# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA
- Is data science just applied stats???
  - origin of the field
  - goals and the types of problems

# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA
- Is data science just applied stats???
  - origin of the field
  - goals and the types of problems
  - data and technology involved

# Statistics vs. Data Science

- “Statistics is the science of learning from data...” – ASA
- Is data science just applied stats???
  - origin of the field
  - goals and the types of problems
  - data and technology involved
- “fair to consider applied statistics as a subset of data science”:
  - harvesting, processing, storing and cleaning are more central to data science than hard core statistics


# Outline

- ① Course Summary
- ② Basic Probability Theory
- ③ Random Variables
- ④ Bivariate Distributions
- ⑤ Code Demo



# Probability and Statistics

"Probability is the vehicle of statistics."



Copyright (c) 2004 Brooks/Cole, a division of Thomson Learning, Inc.

# Probability Spaces

Definition: A probability space is a triple  $(\Omega, \mathcal{F}, \mathbb{P})$  consisting of:

- A sample space  $\Omega$ , which contains all possible outcomes.
- A set of events  $\mathcal{F}$ , which must be a  $\sigma$ -algebra defined on  $\Omega$ .  $\mathcal{F}$  is a collection of subsets of  $\Omega$  such that:

# Probability Spaces

Definition: A probability space is a triple  $(\Omega, \mathcal{F}, \mathbb{P})$  consisting of:

- A sample space  $\Omega$ , which contains all possible outcomes.
- A set of events  $\mathcal{F}$ , which must be a  $\sigma$ -algebra defined on  $\Omega$ .  $\mathcal{F}$  is a collection of subsets of  $\Omega$  such that:
  - $\Omega \in \mathcal{F}$
  - if  $A \in \mathcal{F}$ , then  $A^c \in \mathcal{F}$
  - if  $A_1, A_2, \dots, A_i, \dots \in \mathcal{F}$ , then  $\cup_i A_i \in \mathcal{F}$
- A probability measure  $\mathbb{P}$ , which can assign probabilities to the events in  $\mathcal{F}$ .

# Probability Spaces Example

- Flip a coin:
  - sample space  $\Omega$ :
  - $\sigma$ -algebra  $\mathcal{F}$ :
  - probability measure  $\mathbb{P}$ :

# Probability Spaces Example

- Flip a coin:
  - sample space  $\Omega$ :
  - $\sigma$ -algebra  $\mathcal{F}$ :
  - probability measure  $\mathbb{P}$ :
- Generalize to any finite sample space:

# Probability Measure

Definition: A probability measure  $\mathbb{P}$  assigns each even  $A \in \mathcal{F}$  to a real number, and it satisfies the following **three axioms**:

①  $0 \leq \mathbb{P}(A) \leq 1, \quad \forall A \in \mathcal{F}$

# Probability Measure

Definition: A probability measure  $\mathbb{P}$  assigns each even  $A \in \mathcal{F}$  to a real number, and it satisfies the following **three axioms**:

- ①  $0 \leq \mathbb{P}(A) \leq 1, \quad \forall A \in \mathcal{F}$
- ②  $\mathbb{P}(\emptyset) = 0$  and  $\mathbb{P}(\Omega) = 1$

# Probability Measure

Definition: A probability measure  $\mathbb{P}$  assigns each even  $A \in \mathcal{F}$  to a real number, and it satisfies the following **three axioms**:

- ①  $0 \leq \mathbb{P}(A) \leq 1, \quad \forall A \in \mathcal{F}$
- ②  $\mathbb{P}(\emptyset) = 0$  and  $\mathbb{P}(\Omega) = 1$
- ③ If  $A_1, A_2, \dots, A_n \dots$  are disjoint then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$$



# Probability Measure

Definition: A probability measure  $\mathbb{P}$  assigns each even  $A \in \mathcal{F}$  to a real number, and it satisfies the following **three axioms**:

- ①  $0 \leq \mathbb{P}(A) \leq 1, \quad \forall A \in \mathcal{F}$
- ②  $\mathbb{P}(\emptyset) = 0$  and  $\mathbb{P}(\Omega) = 1$
- ③ If  $A_1, A_2, \dots, A_n \dots$  are disjoint then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$$

Based on 1 - 3, can you show  $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$  ?

# Conditional Probability

- Intuition:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$

# Conditional Probability

- Intuition:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$

- **Bayes Theorem:**

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

# Conditional Probability

- Intuition:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$

- **Bayes Theorem:**

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

- **Law of total probability:**  $A_1, A_2, \dots$  be a partition of  $\Omega$

$$\mathbb{P}(S) = \sum_{i=1} \mathbb{P}(S \cap A_i) = \sum_{i=1} \mathbb{P}(S|A_i)\mathbb{P}(A_i)$$

# Bayes Theorem

- Apply Bayes Theorem and (simple case) of Law of total probability:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c)}$$

# Bayes Theorem

- Apply Bayes Theorem and (simple case) of Law of total probability:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c)}$$

- Example: Suppose 1% of population has certain cancer. A medical test can detect this cancer with 100% accuracy, but also has 1% false positive rate. If a person randomly selected tested positive, what is the probability this person actually has the cancer?

# Independence

Definition: Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space. Two events  $A, B \in \mathcal{F}$  are independent **if and only if**:

$$\mathbb{P}(A|B) = \mathbb{P}(A)$$

# Independence

Definition: Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space. Two events  $A, B \in \mathcal{F}$  are independent **if and only if**:

$$\mathbb{P}(A|B) = \mathbb{P}(A)$$

Equivalently, if  $A$  and  $B$  are independent then:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$$



# Conditional Independence

Definition: Two events  $A, B \in \mathcal{F}$  are independent conditioned on another event  $C$  **if and only if**:

$$\mathbb{P}(A|B, C) = \mathbb{P}(A|C)$$

- It is possible that  $A$  and  $B$  are not independent, but conditioned on  $C$  they are (and vice versa!)

# Conditional Independence

Definition: Two events  $A, B \in \mathcal{F}$  are independent conditioned on another event  $C$  **if and only if**:

$$\mathbb{P}(A|B, C) = \mathbb{P}(A|C)$$

- It is possible that  $A$  and  $B$  are not independent, but conditioned on  $C$  they are (and vice versa!)
- A box contains two coins: a regular coin and one fake two-headed coin. I choose a coin at random and toss it twice. Define the following events:
  - $A$  = 1st coin toss is a head.
  - $B$  = 2nd coin toss is a head.
  - $C$  = the regular coin has been selected.

# Monte Carlo Simulation

- How to calculate the probability without doing the math?
  - interpretation of probability: long-term (relative) frequency

# Monte Carlo Simulation

- How to calculate the probability without doing the math?
  - interpretation of probability: long-term (relative) frequency
- Run a computer program to repeatedly simulate a random process!
  - Monte Carlo simulation
  - random number generator in computers

# Monte Carlo Simulation

- How to calculate the probability without doing the math?
  - interpretation of probability: long-term (relative) frequency
- Run a computer program to repeatedly simulate a random process!
  - Monte Carlo simulation
  - random number generator in computers
- An approximation of the true probability
  - good enough approximation for “sufficient” repetition

# Outline

- ① Course Summary
- ② Basic Probability Theory
- ③ Random Variables
- ④ Bivariate Distributions
- ⑤ Code Demo

# Definition

What is a random variable?

- English: a variable whose value depends on some random event.

# Definition

What is a random variable?

- English: a variable whose value depends on some random event.
- Mathy: given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , a random variable  $X$  is a function that maps the random events in  $\mathcal{F}$  to some real numbers  $\mathbb{R}$ .



# Definition

What is a random variable?

- English: a variable whose value depends on some random event.
- Mathy: given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , a random variable  $X$  is a function that maps the random events in  $\mathcal{F}$  to some real numbers  $\mathbb{R}$ .
- Rigorous: in measure theory and advanced probability theory, for example [here](#).

# Definition

What is a random variable?

- English: a variable whose value depends on some random event.
- Mathy: given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , a random variable  $X$  is a function that maps the random events in  $\mathcal{F}$  to some real numbers  $\mathbb{R}$ .
- Rigorous: in measure theory and advanced probability theory, for example [here](#).

Notation:

- capital letter  $X$  for the random variable
- lower case  $x$  for values the random variable can take

# Discrete Random Variables

- A discrete random variable  $X$  can only take countable values in  $\mathbb{R}$ .

# Discrete Random Variables

- A discrete random variable  $X$  can only take countable values in  $\mathbb{R}$ .
- A probability mass function (pmf)  $P_X(x)$  is defined as:

$$P_X(x) = \mathbb{P}(X = x)$$

# Discrete Random Variables

- A discrete random variable  $X$  can only take countable values in  $\mathbb{R}$ .
- A probability mass function (pmf)  $P_X(x)$  is defined as:

$$P_X(x) = \mathbb{P}(X = x)$$

- Important discrete random variables:
  - Uniform
  - Bernoulli
  - Binomial
  - Poisson
  - See more [here](#)

# Continuous Random Variables

- A continuous random variable  $X$  can take values in interval(s) in  $\mathbb{R}$ .

# Continuous Random Variables

- A continuous random variable  $X$  can take values in interval(s) in  $\mathbb{R}$ .
- A cumulative distribution function (cdf)  $F_X(x)$  is defined as:

$$F_X(x) = \mathbb{P}(X \leq x)$$

# Continuous Random Variables

- A continuous random variable  $X$  can take values in interval(s) in  $\mathbb{R}$ .
- A cumulative distribution function (cdf)  $F_X(x)$  is defined as:

$$F_X(x) = \mathbb{P}(X \leq x)$$

- A probability density function (pdf)  $f_X(x)$  is defined as:

$$f_X(x) = F'_X(x)$$

for all points  $x$  at which  $F_X(x)$  is differentiable.



# Continuous Random Variables

Some properties of cdf  $F_X(x)$  and pdf  $f_X(x)$ :

# Continuous Random Variables

Some properties of cdf  $F_X(x)$  and pdf  $f_X(x)$ :

- $0 \leq F_X(x) \leq 1$ ,  $F_X(-\infty) = 0$ ,  $F_X(\infty) = 1$ ,

# Continuous Random Variables

Some properties of cdf  $F_X(x)$  and pdf  $f_X(x)$ :

- $0 \leq F_X(x) \leq 1$ ,  $F_X(-\infty) = 0$ ,  $F_X(\infty) = 1$ ,
- $F_X(x)$  is non-decreasing:  $F_X(x_1) \leq F_X(x_2)$  for all  $x_1 \leq x_2$

# Continuous Random Variables

Some properties of cdf  $F_X(x)$  and pdf  $f_X(x)$ :

- $0 \leq F_X(x) \leq 1$ ,  $F_X(-\infty) = 0$ ,  $F_X(\infty) = 1$ ,
- $F_X(x)$  is non-decreasing:  $F_X(x_1) \leq F_X(x_2)$  for all  $x_1 \leq x_2$
- $f_X(x)$  is non-negative:  $f_X(x) \geq 0$  for all  $x \in \mathbb{R}$

# Continuous Random Variables

Some properties of cdf  $F_X(x)$  and pdf  $f_X(x)$ :

- $0 \leq F_X(x) \leq 1$ ,  $F_X(-\infty) = 0$ ,  $F_X(\infty) = 1$ ,
- $F_X(x)$  is non-decreasing:  $F_X(x_1) \leq F_X(x_2)$  for all  $x_1 \leq x_2$
- $f_X(x)$  is non-negative:  $f_X(x) \geq 0$  for all  $x \in \mathbb{R}$
- $\int_{-\infty}^{\infty} f_X(x) = 1$

# Continuous Random Variables

- Important continuous random variables:
  - Uniform
  - Normal (Gaussian)
  - Student-t
  - Cauchy
  - Exponential
  - Gamma
  - Beta
  - $\chi^2$
- See more [here](#)

# Functions of Random Variables

- Let  $X$  be a random variable and  $Y = g(X)$  be another random variable defined by a function  $g : \mathbb{R} \rightarrow \mathbb{R}$ .

# Functions of Random Variables

- Let  $X$  be a random variable and  $Y = g(X)$  be another random variable defined by a function  $g : \mathbb{R} \rightarrow \mathbb{R}$ .
- The cdf of  $Y$  is given by:

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(g(X) \leq y) = \int_{\{x|g(x) \leq y\}} f_X(x) dx$$



# Functions of Random Variables

- Let  $X$  be a random variable and  $Y = g(X)$  be another random variable defined by a function  $g : \mathbb{R} \rightarrow \mathbb{R}$ .
- The cdf of  $Y$  is given by:

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(g(X) \leq y) = \int_{\{x | g(x) \leq y\}} f_X(x) dx$$

- If  $g$  is a strictly increasing function, then

$$F_Y(y) = \int_{-\infty}^{g^{-1}(y)} f_X(x) dx = F_X(g^{-1}(y))$$

$$f_Y(y) = f_X(g^{-1}(y)) \frac{d}{dy} g^{-1}(y)$$

# Generating Random Variables

- Let  $X$  be a random variable with a continuous CDF  $F_X(x)$  and define the random variable  $Y = F_X(x)$ . Can you show  $Y \sim U[0, 1]$ ?

# Generating Random Variables

- Let  $X$  be a random variable with a continuous CDF  $F_X(x)$  and define the random variable  $Y = F_X(x)$ . Can you show  $Y \sim U[0, 1]$ ?
- This is how most computer programs are generating random variables of common distributions:
  - sample from Uniform  $[0, 1]$   $u_1, u_2, \dots, u_n$
  - get  $x_i = F_X^{-1}(u_i)$  for a specific distribution with cdf  $F_X(\cdot)$

# Outline

- ① Course Summary
- ② Basic Probability Theory
- ③ Random Variables
- ④ Bivariate Distributions
- ⑤ Code Demo

# Joint Distributions

A joint probability mass function:

	Y=0	Y=1	
X=0	1/10	2/10	3/10
X=1	3/10	4/10	7/10
	4/10	6/10	1

- Marginal distributions of  $X$  and  $Y$ ?

# Joint Distributions

A joint probability mass function:

	Y=0	Y=1	
X=0	1/10	2/10	3/10
X=1	3/10	4/10	7/10
	4/10	6/10	1

- Marginal distributions of  $X$  and  $Y$ ?
- Conditional distribution of  $X|Y = 0$  and  $Y|X = 1$ ?

# Joint Distributions

A joint probability density function:

$$f_{X,Y}(x,y) = \begin{cases} x+y & \text{if } 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Marginal distributions of  $X$  and  $Y$ ?

# Joint Distributions

A joint probability density function:

$$f_{X,Y}(x,y) = \begin{cases} x+y & \text{if } 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Marginal distributions of  $X$  and  $Y$ ?
- Conditional distribution of  $X|Y = 1$ ?



# Independence and IID Samples

- Two random variables  $X$  and  $Y$  are independent **if and only if** the joint pdf  $f_{X,Y}(x,y) = f_X(x) \cdot f_Y(y)$  for all values  $x$  and  $y$ .

# Independence and IID Samples

- Two random variables  $X$  and  $Y$  are independent **if and only if** the joint pdf  $f_{X,Y}(x,y) = f_X(x) \cdot f_Y(y)$  for all values  $x$  and  $y$ .
- If  $X_1, X_2, \dots, X_n$  are independent and each has the same marginal distribution of pdf  $f_X(\cdot)$ , we say that  $X_1, X_2, \dots, X_n$  are IID (independent and identically distributed) and has joint pdf:

$$f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = \prod_{i=1}^n f_X(x_i)$$

# Outline

- ① Course Summary
- ② Basic Probability Theory
- ③ Random Variables
- ④ Bivariate Distributions
- ⑤ Code Demo

# Python Jupyter Notebook

- Birthday problem.
- Monty Hall problem.
- Su's favorite probability brain teaser.

# Python Jupyter Notebook

- Birthday problem.
- Monty Hall problem.
- Su's favorite probability brain teaser.
- Assignment this week:
  - Install Python 3 and Jupyter Notebook through [Anaconda](#)
    - install tutorial for [mac](#) and [window](#)
  - Run the demo notebook and make sure nothing breaks.
  - Homework 1 posted, due in 2 weeks, please start early!