

Point Separation and Obstacle Removal by Finding and Hitting Odd Cycles

Jie Xue

New York University Shanghai

joint work with


Neeraj Kumar (University of California, Santa Barbara)

Daniel Lokshtanov (University of California, Santa Barbara)


Saket Saurabh (Institute of Mathematical Sciences, Chennai)

Subhash Suri (University of California, Santa Barbara)

- **Obstacles:** connected geometric objects in the plane




- **Obstacles:** connected geometric objects in the plane




The obstacles separate s and t

- **Obstacles:** connected geometric objects in the plane



The obstacles do not separate s and t

- **Obstacles:** connected geometric objects in the plane




The obstacles do not separate s and t

- **Point separation**

Input: $s, t \in \mathbb{R}^2$ and a set \mathcal{O} of obstacles in \mathbb{R}^2


Output: a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ that separates s and t



- **Point separation**


Input: $s, t \in \mathbb{R}^2$ and a set \mathcal{O} of obstacles in \mathbb{R}^2

Output: a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ that separates s and t




- **A generalization of point separation**

Given p point pairs $(s_1, t_1), \dots, (s_p, t_p)$ in \mathbb{R}^2 , want a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ that separates every pair (s_i, t_i) .



- **A generalization of point separation**


Given p point pairs $(s_1, t_1), \dots, (s_p, t_p)$ in \mathbb{R}^2 , want a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ that separates every pair (s_i, t_i) .



- **Obstacle removal**

Input: $s, t \in \mathbb{R}^2$ and a set \mathcal{O} of obstacles in \mathbb{R}^2


Output: a minimum $\mathcal{O}' \subseteq \mathcal{O}$ so that $\mathcal{O} \setminus \mathcal{O}'$ does not separate s and t



- **Obstacle removal**

Input: $s, t \in \mathbb{R}^2$ and a set \mathcal{O} of obstacles in \mathbb{R}^2

Output: a minimum $\mathcal{O}' \subseteq \mathcal{O}$ so that $\mathcal{O} \setminus \mathcal{O}'$ does not separate s and t



- **Cabello and Giannopoulos [SoCG'13]**

Point separation for one pair is polynomial-time solvable.

- **Cabello and Giannopoulos [SoCG'13]**

Point separation for one pair is polynomial-time solvable.

- **Eiben and Lokshtanov [SoCG'20]**

Obstacle removal is fixed-parameter tractable (FPT) parameterized by the solution size k : it can be solved in $k^{O(k^3)} \cdot n^{O(1)}$ time.

- **Cabello and Giannopoulos [SoCG'13]**

Point separation for one pair is **polynomial-time solvable**.

- **Eiben and Lokshtanov [SoCG'20]**

Obstacle removal is **fixed-parameter tractable (FPT)** parameterized by the solution size k : it can be solved in $k^{O(k^3)} \cdot n^{O(1)}$ time.

- **Kumar et al. [SODA'21]**

Obstacle removal admits a (complicated) **constant approximation**.

Main results of our paper

- **Point separation (with multiple pairs)**
 - Can be solved in $n^{O(p)}$ time for p pairs of points
 \implies Polynomial-time solvable for any fixed p
 - Impossible to achieve $n^{\Omega(p/\log p)}$ time assuming ETH

Main results of our paper

- **Point separation (with multiple pairs)**

- Can be solved in $n^{O(p)}$ time for p pairs of points
 \implies **Polynomial-time solvable** for any fixed p
- Impossible to achieve $n^{\Omega(p/\log p)}$ time assuming ETH
- For **pseudo-disk** obstacles:
 $n^{O(\sqrt{p})}$ time upper bound and $n^{\Omega(\sqrt{p}/\log p)}$ time lower bound

Main results of our paper

- **Point separation (with multiple pairs)**

- Can be solved in $n^{O(p)}$ time for p pairs of points
 \implies **Polynomial-time solvable** for any fixed p
- Impossible to achieve $n^{\Omega(p/\log p)}$ time assuming ETH
- For **pseudo-disk** obstacles:
 $n^{O(\sqrt{p})}$ time upper bound and $n^{\Omega(\sqrt{p}/\log p)}$ time lower bound

- **Obstacle removal**

- Improved FPT algorithm: $2.3146^k \cdot n^{O(1)}$ time
- **Simpler** constant-approximation algorithm

A structural characterization

- All our algorithmic results are based on a **structural characterization** for determining whether a set of obstacles separate a pair of points (and its generalization to **multiple point pairs**).

A structural characterization

- All our algorithmic results are based on a **structural characterization** for determining whether a set of obstacles separate a pair of points (and its generalization to **multiple point pairs**).

Theorem (one-pair version)

*Given two points $s, t \in \mathbb{R}^2$ and a set \mathcal{O} of obstacles in \mathbb{R}^2 , there exists a (multi-)graph G with vertex set \mathcal{O} and edges labeled from $\{0, 1\}$ such that a subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains **a cycle whose sum of labels is odd**.*

The graph G can be built in $n^{O(1)}$ time given s, t and \mathcal{O} where $n = |\mathcal{O}|$.

A structural characterization

- More pairs to be separated?

A structural characterization

- More pairs to be separated?

Theorem (multi-pair version)

Given p point pairs $(s_1, t_1), \dots, (s_p, t_p)$ in \mathbb{R}^2 and a set \mathcal{O} of obstacles in \mathbb{R}^2 , there exists a multi-graph G with vertex set \mathcal{O} and edges labeled from $\{0, 1\}^p$ such that a subset $\mathcal{O}' \subseteq \mathcal{O}$ separates all p point pairs iff for every $i \in \{1, \dots, p\}$ the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels has an odd i -th entry.

The graph G can be built in $2^{O(p)} \cdot n^{O(1)}$ time given the pairs and \mathcal{O} .

- **Obstacle removal**
- Using our characterization theorem, we construct the multi-graph G .
- We want a minimum subset $\mathcal{X} \subseteq \mathcal{O}$ such that $\mathcal{O}' := \mathcal{O} \setminus \mathcal{X}$ does not separate s and t , i.e., $G[\mathcal{O}']$ does not contain an odd labeled cycle.

- **Obstacle removal**

- Using our characterization theorem, we construct the multi-graph G .
- We want a minimum subset $\mathcal{X} \subseteq \mathcal{O}$ such that $\mathcal{O}' := \mathcal{O} \setminus \mathcal{X}$ does not separate s and t , i.e., $G[\mathcal{O}']$ does not contain an odd labeled cycle.
- (MINIMUM) ODD CYCLE TRANSVERSAL
Given a graph G , find a minimum subset X of vertices of G that hit all odd cycles in G (or equivalently, $G - X$ is bipartite).

- **Obstacle removal**

- Using our characterization theorem, we construct the multi-graph G .
- We want a minimum subset $\mathcal{X} \subseteq \mathcal{O}$ such that $\mathcal{O}' := \mathcal{O} \setminus \mathcal{X}$ does not separate s and t , i.e., $G[\mathcal{O}']$ does not contain an odd labeled cycle.

- (MINIMUM) ODD CYCLE TRANSVERSAL

Given a graph G , find a minimum subset X of vertices of G that hit all odd cycles in G (or equivalently, $G - X$ is bipartite).

- (PARAMETERIZED) ODD CYCLE TRANSVERSAL

Given a graph G and a parameter $k \geq 0$, find a subset X of at most k vertices of G that hit all odd cycles in G .


- Relate obstacle removal to ODD CYCLE TRANSVERSAL?

Applications

- Relate obstacle removal to ODD CYCLE TRANSVERSAL?
- Subdivide edges of label 0 (odd labeled cycles \rightarrow odd cycles)


Applications

- Relate obstacle removal to ODD CYCLE TRANSVERSAL?
- Subdivide edges of label 0 (odd labeled cycles \rightarrow odd cycles)



Applications

- Relate obstacle removal to ODD CYCLE TRANSVERSAL?
- Subdivide edges of label 0 (odd labeled cycles \rightarrow odd cycles)



- **FPT algorithm:** Directly apply the $2.3146^k \cdot n^{O(1)}$ -time algorithm for (PARAMETERIZED) ODD CYCLE TRANSVERSAL.
- **Constant approximation:** LP rounding using the technique of Lee (similar to the basic idea of Kumar et al. [SODA'21]).

- **Point separation**
- Using our characterization theorem, we construct the multi-graph G .
- We want a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ such that for any $i \in \{1, \dots, p\}$, $G[\mathcal{O}']$ contains a cycle with odd i -th entry.

- **Point separation**

- Using our characterization theorem, we construct the multi-graph G .
- We want a minimum subset $\mathcal{O}' \subseteq \mathcal{O}$ such that for any $i \in \{1, \dots, p\}$, $G[\mathcal{O}']$ contains a cycle with odd i -th entry.

- **Key observation:**

There is a subgraph H of $G[\mathcal{O}']$ that consists of $O(p)$ special vertices and $O(p)$ paths connecting these special vertices such that for every $i \in \{1, \dots, p\}$, H contains a cycle with odd i -th entry.

- Now our task becomes computing a subgraph H of G with fewest vertices satisfying the condition in the key observation.

Applications

- Now our task becomes computing a subgraph H of G with fewest vertices satisfying the condition in the key observation.
- Solve this problem in $n^{O(p)}$ time?

- Now our task becomes computing a subgraph H of G with fewest vertices satisfying the condition in the key observation.
- Solve this problem in $n^{O(p)}$ time?
 - 1 Guess the structure of H and the label parities of the paths in H .
 - 2 Guess the special vertices of H .
 - 3 For each path in H connecting X and Y with label parity $\ell \in \{0, 1\}^p$, find the **shortest path** in G between X and Y with label parity ℓ .


- Now our task becomes computing a subgraph H of G with fewest vertices satisfying the condition in the key observation.
- Solve this problem in $n^{O(p)}$ time?
 - 1 Guess the structure of H and the label parities of the paths in H .
 - 2 Guess the special vertices of H .
 - 3 For each path in H connecting X and Y with label parity $\ell \in \{0, 1\}^p$, find the **shortest path** in G between X and Y with label parity ℓ .
- In this way, we can compute H and solve the problem in $n^{O(p)}$ time.

Proof sketch of the characterization theorem

- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
 - (*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.

Proof sketch of the characterization theorem


- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
 - (*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.



Fix a curve π connecting s and t in the plane.

Proof sketch of the characterization theorem


- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
 - (*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.



For each obstacle X , fix a reference point inside X .

Proof sketch of the characterization theorem


- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
(*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.



If there exists a curve connecting the ref points of X and Y in $X \cup Y$ that crosses π an **even** number of times, add an edge (X, Y) to G with label 0.

Proof sketch of the characterization theorem


- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
 - (*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.



If there exists a curve connecting the ref points of X and Y in $X \cup Y$ that crosses π an **odd** number of times, add an edge (X, Y) to G with label 1.

Proof sketch of the characterization theorem

- Recall that we need a multi-graph G whose vertex set is \mathcal{O} and edges are labeled from $\{0, 1\}$ such that
 - (*) A subset $\mathcal{O}' \subseteq \mathcal{O}$ of obstacles separates s and t iff the induced subgraph $G[\mathcal{O}']$ contains a cycle whose sum of labels is odd.



An odd labeled cycle in the graph corresponds to a closed curve in the union of the obstacles that crosses π an odd number of times.

Open questions

- More restricted obstacles?

Open questions

- More restricted obstacles?
- Approximation for point separation?

Open questions

- More restricted obstacles?
- Approximation for point separation?
- Higher dimensions?

Thank you!
Q & A