

SOLUTION FOR HOMEWORK 5, STAT 6331

Welcome to your fifth homework. It is the beginning of our point-estimation part.

Reminder: if you find a mistake/misprint, do not e-mail or call me. Write it down on the first page of your solutions and you may give yourself a partial credit — but keep in mind that the total for your homeworks cannot exceed 20 points.

Now let us look at your problems.

1. Exerc.7.1. Note that $\hat{\theta}_{MLE} = \psi(X)$ meaning that it depends on a single outcome. Then for each value of X we choose the value of parameter that yields the largest likelihood (here $f(X|\hat{\theta}_{MLE}) := \max_{\theta \in \{1,2,3\}} f(X|\theta)$).

Answer: $\hat{\theta}_{MLE}$ is 1,1,(2 or 3),3,3 for X being 0,1,2,3,4, respectively.

2. Exerc. 7.3. In short, this is because likelihood is nonnegative and $\log(x)$ is a monotonically increasing function.

Nonetheless, let us present a rigorous solution. We want to show that if

$$\hat{\theta} := \operatorname{argmax}_{\theta \in \Omega} L(\theta|\mathbf{X}), \quad \hat{\mu} := \operatorname{argmax}_{\theta \in \Omega} \log(L(\theta|\mathbf{X})),$$

then $\hat{\theta} = \hat{\mu}$ a.s. Assume that the latter is wrong. Then

$$L(\hat{\theta}|\mathbf{X}) > L(\hat{\mu}|\mu), \quad \log(L(\hat{\theta}|\mathbf{X})) < \log(L(\hat{\mu}|\mu)).$$

But $\log(z)$ is the monotonically increasing function for $z > 0$ meaning that $z_1 < z_2$ iff $\log(z_1) < \log(z_2)$. The contradictory proves that $\hat{\theta} = \hat{\mu}$, and we can search after the MLE using either likelihood or loglikelihood.

3. Exers. 7.6. Given: X is distributed according to the pdf

$$f(x|\theta) = \theta x^{-2} I(0 < \theta \leq x < \infty).$$

Note how I wrote the pdf using the indicator function! This is a very important step!

Solution: (a) Write for a sample of size n ,

$$f(\mathbf{x}|\theta) = \theta^n \prod_{l=1}^n x_l^{-2} I(0 < \theta \leq x_{(1)}).$$

By the Factorization Theorem $X_{(1)}$ is a SS.

(b) To find MLE we need to use graphic (this is always the case when parameter defines the support). From the graphic it is clear that $\hat{\theta}_{MLE} = X_{(1)}$. Note that it is our sufficient statistic!

(c) Let us calculate the expectation of X :

$$\mu = E(X) = \theta \int_{\theta}^{\infty} xx^{-2}dx = \infty.$$

As a result, there is no standard/classical method of moments estimator for the problem.

Remark: If you desire to get a generalized method of moments estimator, you can choose, for instance $E(X^{1/2})$ or something like this.

4. Exerc. 7.7 Let us look at where $1^n \geq \prod_{l=1}^n 1/[2x_l^{1/2}]$. This is equivalent to $\prod_{l=1}^n x_l \geq 2^{-2n}$. As a result, the MLE is 0 if $\prod_{l=1}^n X_l \geq 2^{-2n}$ and the MLE is 1 otherwise.

5. Exerc. 7.8. Given: $X \sim N(0, \sigma^2)$. The sample size $n = 1$.

(a). Clearly $\hat{\theta} = X^2$ is unbiased estimator of σ^2 because

$$E(\hat{\theta}) = E(X^2) = \text{Var}X + E^2(X) = \sigma^2 + 0 = \sigma^2.$$

(b) Here

$$L(\sigma|X) = (2\pi\sigma^2)^{-1/2}e^{-X^2/(2\sigma^2)}.$$

It is more convenient to work with loglikelihood,

$$l(\sigma|X) = -(1/2)\log(2\pi) - (1/2)\log(\sigma^2) - X^2/(2\sigma^2) = -(1/2)\log(2\pi) - \log(\sigma) - X^2/(2\sigma^2).$$

Before taking the derivative, let us look at a graphic.

Now take derivative with respect to σ and get

$$dl(\sigma|X)/d\sigma = -\frac{1}{\sigma} + \frac{X^2}{\sigma^3}.$$

Equate to zero, remember that $\sigma > 0$, and the solution is $\sigma_{MLE} = |X|$. Then check, via graphic or second derivative, that this solution is indeed the maximum of the likelihood (do not forget about this step on your exams — you may not get a credit for solution if you do not check that the extreme point is indeed the maximum!).

(c) Let us begin with the first moment. We have $E_{\sigma}(X) = 0$, and because the population mean does not depend on σ , we cannot use the equality of the population and sample mean for estimation of σ . For the second moment the situation is better, and we have

$$E(X^2) = \sigma^2$$

which should be equated to X^2 . Again, remember that $\sigma > 0$ so solution of

$$\sigma^2 = X^2$$

yields $\hat{\sigma}_{MME} = |X|$.

6. Exerc. 7.9. Here we have a sample of size n from a distribution with the pdf $f(x|\theta) = \theta^{-1}I(0 \leq x \leq \theta)$ and $\theta \in \Omega := (0, \infty)$.

(a) Method of Moments. Write

$$\mu_1 = E(X) = \theta^{-1} \int_0^\theta x dx = \theta/2.$$

As a result, the MME is defined as the solution of

$$\hat{\theta}_{MME}/2 = \bar{X}.$$

We get $\hat{\theta}_{MME} = 2\bar{X}$.

(b) Let us calculate the MLE. I may use $\underline{X} := \mathbf{X} := X^n$ to denote a sample. These are more popular notations used in the literature. Write

$$L(\theta|X^n) = \theta^{-n}I(0 \leq X_{(1)})I(X_{(n)} \leq \theta).$$

Look at the corresponding graphic.

This yields $\hat{\theta}_{MLE} = X_{(n)}$.

(c) Now let us compare statistical properties of these two estimators. For the method of moments we have

$$E(\hat{\theta}_{MME}) = \theta,$$

(note that this we have by definition of the used MME). For iid observations,

$$\begin{aligned} \text{Var}(\hat{\theta}_{MME}) &= \text{Var}(2\bar{X}) = 4n^{-2}\text{Var}\left(\sum_{l=1}^n X_l\right) \\ &= (4/n)\text{Var}(X) = (4/n)[E(X^2) - (E(X))^2] = (4/n)\left[\theta^{-1} \int_0^\theta x^2 dx - (\theta/2)^2\right] \\ &= (4/n)\theta^2[(1/3) - (1/4)] = n^{-1}\theta^2/3. \end{aligned}$$

We conclude that

$$E(\hat{\theta}_{MME}) = \theta; \quad \text{Var}(\hat{\theta}_{MME}) = n^{-1}[\theta^2/3]. \quad (1)$$

For the MLE we must begin with the distribution of $X_{(n)}$. Its cdf is, for $0 < x < \theta$,

$$\begin{aligned} F(x|\theta) &= P(X_{(n)} \leq x) = P(X_1 \leq x, \dots, X_n \leq x) = \prod_{l=1}^n P(X_l \leq x < \theta) \\ &= [\theta^{-1} \int_0^x du]^n I(x < \theta) = \theta^{-n} x^n I(0 < x < \theta). \end{aligned}$$

This yields the pdf

$$f_{X_{(n)}}(x) = dF(u|\theta)/du \Big|_{u=x} = n\theta^{-n} x^{n-1} I(0 < x < \theta)$$

Now we are in a position to make all calculations for the MLE. Write,

$$\begin{aligned} E(X_{(n)}) &= \int_0^\theta n\theta^{-n} x^{n-1} x dx = n\theta^{-n} \int_0^\theta x^n dx \\ &= n\theta^{-n} (n+1)^{-1} \theta^{n+1} = [n/(n+1)]\theta. \end{aligned}$$

Further,

$$E(X_{(n)}^2) = \int_0^\theta n\theta^{-n} x^{n-1} x^2 dx = n\theta^{-n} \int_0^\theta x^{n+1} dx = [n/(n+2)]\theta^2.$$

We conclude that

$$\text{Var}(\hat{\theta}_{MLE}) = E(\hat{\theta}_{MLE}^2) - [E(\hat{\theta}_{MLE})]^2 = [n/(n+2)]\theta^2 - [n/(n+1)]^2\theta^2 = \frac{n}{(n+2)(n+1)^2}\theta^2.$$

Let us combine for the MLE the results

$$E(\hat{\theta}_{MLE}) = \frac{n}{n+1}\theta, \quad \text{Var}(\hat{\theta}_{MLE}) = \frac{n}{(n+2)(n+1)^2}\theta^2. \quad (2)$$

Now we are comparing (1) and (2). The MME is unbiased while the MLE is only asymptotically unbiased. On the other hand, the MLE has smaller variance. As a result, the choice is based on practical interests. To avoid such a dilemma, typically another risk function, the mean squared error, $E(\hat{\theta} - \theta)^2$, is used to choose the better estimator.

7. Exerc. 7.10(a,b). Note that the cdf is given here, which can be written as

$$F(x|\alpha, \beta) = \min(1, (x/\beta)^\alpha) I(x > 0), \quad (\alpha, \beta) \in (0, \infty)^2.$$

Remark: can α be negative?

This is a continuous RV with the pdf (take accurately derivative of the cdf)

$$f(x|\alpha, \beta) = \beta^{-\alpha} \alpha x^{\alpha-1} I(0 \leq x \leq \beta).$$

Now we can solve the problems.

(a) To find a SS, via using the Factorization Theorem, we write

$$f(\mathbf{x}|\alpha, \beta) = \beta^{-n\alpha} \alpha^n e^{(\alpha-1)\sum_{l=1}^n \ln(x_l)} I(x_{(1)} \geq 0) I(x_{(n)} \leq \beta).$$

By the FT the statistic $(T_1, T_2) := (\sum_{l=1}^n \ln(X_l), X_{(n)})$ is a two-dimensional SS.

(b) To find the MLE we write using the above-written pdf,

$$L(\alpha, \beta | \mathbf{X}) = \beta^{-n\alpha} \alpha^n e^{(\alpha-1) \sum_{l=1}^n \ln(X_l)} I(X_{(1)} \geq 0) I(X_{(n)} \leq \beta).$$

For a given $\alpha = \alpha_0$, we can graph $L(\alpha_0, \beta | \mathbf{X})$ as a function in β . Note that $\beta \geq 0$ by the assumption.

It follows from the graphic that, regardless of α_0 , the MLE of β is $\hat{\beta}_{MLE} = X_{(n)}$. Now let $\beta = \beta_0$. The log-likelihood is

$$\begin{aligned} l(\alpha, \beta_0 | \mathbf{X}) &:= \ln(L(\alpha, \beta_0 | \mathbf{X})) \\ &= \alpha[-n \ln(\beta_0) + T_1] + n \ln(\alpha) + g \end{aligned}$$

where the term g does not depend on α .

Note that we are interested in a specific $\beta_0 = \hat{\beta}_{MLE} = X_{(n)}$, and this yields that

$$[-n \ln(X_{(n)}) + T_1] = [n(-\ln(X_{(n)})) + n^{-1} \sum_{l=1}^n \ln(X_l)] \leq 0$$

with the equality iff $\alpha = 0$. Thus we have the following graphic:

We have a global maximum (you can also get this by checking the second derivative). Let us find the extreme point via taking the derivative,

$$dl(\alpha, \beta_0 | \mathbf{X})/d\alpha = -n \ln(\beta_0) + T_1 + n/\alpha.$$

Thus, the extreme point is the solution of the equation $-n \ln(\beta_0) + T_1 + n/\alpha = 0$ which yields

$$\hat{\alpha}_{MLE} = [\ln(\beta_0) - T_1/n]^{-1}.$$

Answer:

$$(\hat{\alpha}_{MLE}, \hat{\beta}_{MLE}) = (\frac{1}{\overline{Z_{(n)}} - \bar{Z}}, X_{(n)}),$$

where $Z_l := \ln(X_l)$.

Remark: Look at how nicely the problem would look if we had taken the logarithmic transformation of the random variable X . Do you know why? Let us consider U with the pdf $f(u|\alpha) = \beta^{-\alpha} \alpha u^{\alpha-1} I(0 \leq u \leq \beta)$. Then $V := \ln(U)$ has the cdf

$$F_V(v) = F_U(e^v) = \int_0^{e^v} \beta^{-\alpha} \alpha u^{\alpha-1} du = \beta^{-\alpha} [e^{\alpha v} - 1]$$

which is the exponential cdf with the pmf

$$f_V(v) = \alpha \beta^{-1} e^{\alpha v} I(-\infty < v < \ln(\beta)).$$

(c) Here $\hat{\beta}_{MLE} = \ln(25)$, and you need a calculator to find $\hat{\alpha}_{MLE}$. I skip it.

8. Exerc. 7.11. Here

$$f(x|\theta) = \theta x^{\theta-1} I(0 \leq x \leq 1), \quad \theta \in \Omega := (0, \infty).$$

(a). Write

$$l(\theta|\mathbf{x}) = n \ln(\theta) + (\theta - 1) \sum_{l=1}^n \ln(X_l).$$

Taking derivative gives us $l'(\theta|\mathbf{x}) = n/\theta + \sum_{l=1}^n \ln(X_l)$, so the extrema is $\theta^* = -[n^{-1} \sum_{l=1}^n \ln(X_l)]^{-1}$.

Is this the point of maximum? Note that $-n^{-1} \sum_{l=1}^n \ln(X_l) > 0$ because $X_l \in [0, 1]$. Thus the graphic of the log-likelihood looks like this:

Or, note that $l''(\theta|\mathbf{X}) = -n/\theta^2 < 0$ so we do have the maximum.

We conclude that $\hat{\theta}_{MLE} = -[n^{-1} \sum_{l=1}^n \ln(X_l)]^{-1}$.

(b) Method of moments:

$$\mu_1 = E(X) = \theta \int_0^1 x x^{\theta-1} dx = \frac{\theta}{\theta+1} x^{\theta+1} \Big|_0^1 = \frac{1}{1+1/\theta}.$$

This yields that

$$\frac{1}{1 + \hat{\theta}_{MME}} = \bar{X}$$

and solving the equation yields $\hat{\theta}_{MME} = \bar{X}/[1 - \bar{X}]$.

Look at how different the MLE and the MME are!

9. Exerc. 7.12. Given the pmf

$$p(x|\theta) = \theta^x(1 - \theta)^{1-x}I(x \in \{0,1\}), \quad \theta \in \Omega := [0, 1/2].$$

Note that this is Bernoulli pmf with the information that the probability of success is at most 1/2. We know that, in a general case of $\theta \in [0, 1]$, the sample mean \bar{X} is the MLE and MME. Let us see what a difference, if any, that prior makes.

(a). (i) We are searching after MME. Because $\mu_1 = E(X) = \theta$, we get $\hat{\theta}_{MME} = \bar{X}$ (no change). Note that this is always the case with a MME because Ω does not affect the estimator.

(ii) Consider MLE:

$$P(\mathbf{X} = \mathbf{x}^n | \theta) = e^{\ln(\theta) \sum_{l=1}^n x_l + \ln(1-\theta) \sum_{l=1}^n (1-x_l)} = e^{n[\ln(\theta)/(1-\theta))\bar{x} + \ln(1-\theta)]}.$$

Further, by taking logarithm we simplify a bit,

$$l(\theta|\mathbf{x}) = n\bar{x} \ln(\theta) + n \ln(1 - \theta)(1 - \bar{x}).$$

To find extreme points, let us take the derivative,

$$l'(\theta|\mathbf{x}) = n\bar{x}\theta^{-1} - n(1 - \bar{x})(1 - \theta)^{-1}.$$

This yields a single extreme point $\theta^* = \bar{x}$. Is it the maximum?

The second derivative is

$$l''(\theta|\mathbf{x}) = -n[\bar{x}\theta^{-2} + (1 - \bar{x})(1 - \theta)^{-2}] < 0.$$

Now is time for graphics.

Note that $\hat{\theta}_{MLE} \in [0, 1/2]$ by its definition because

$$\hat{\theta}_{MLE} := \operatorname{argmax}_{\theta \in \Omega} l(\theta|\mathbf{X}).$$

Answer: $\hat{\theta}_{MLE} = \min(0.5, \bar{X})$. Note how Ω affects the MLE. Also, do you see that the obtained MLE is a projection of our general MLE estimator \bar{X} on Ω ?

Clearly the MLE is better than the MME whenever $\bar{X} > 1/2$ because it is closer to the underlying θ .

b) By definition, the mean squared error is

$$MSE = E_{\theta}(\hat{\theta} - \theta)^2. \quad (3)$$

(i) For the MME, because $n\bar{X}$ is a Binomial RV, namely $Binom(\theta, n)$, we get

$$E(\hat{\theta}_{MME} - \theta)^2 = E(\bar{X} - \theta)^2 = n^{-2}\text{Var}(n\bar{X}) = n^{-2}[n\theta(1 - \theta)] = n^{-1}\theta(1 - \theta).$$

(ii) For the MLE the situation is more complicated,

$$E(\hat{\theta}_{MLE} - \theta)^2 = E\{(\bar{X} - \theta)^2 I(\bar{X} \leq .5)\} + (.5 - \theta)^2 E\{I(\bar{X} > .5)\}.$$

Now I use the fact that $Y := n\bar{X}$ is $Binom(\theta, n)$. Write (I denote by $\lfloor n/2 \rfloor$ the rounded down $n/2$)

$$\begin{aligned} E_{\theta}(\hat{\theta}_{MLE} - \theta)^2 &= E_{\theta}\{(Y/n - \theta)^2 I(Y \leq n/2)\} + (.5 - \theta)^2 P_{\theta}(Y > n/2) \\ &= \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{n!}{k!(n-k)!} \theta^k (1 - \theta)^{n-k} (k/n - \theta)^2 + (.5 - \theta)^2 \sum_{k=\lfloor n/2 \rfloor + 1}^n \frac{n!}{k!(n-k)!} \theta^k (1 - \theta)^{n-k}. \end{aligned}$$

Here we can stop and say that a software should be used to find the number. As I explained earlier, the MME is better than the MMR whenever $\bar{X} > .5$.

10. Exerc. 7.16. (e)(i). Given a sample of size n from the pdf

$$f(x|\theta) = e^{\theta h(x) - H(\theta)} g(x),$$

where $h(x) = dH(x)/dx$ and $h(x)$ is increasing in x .

To find the MLE, note that

$$l(\theta|\mathbf{x}) = \ln(f(\mathbf{x}|\theta)) = \theta \sum_{l=1}^n h(x_l) - nH(\theta) + \sum_{l=1}^n \ln(g(x_l)).$$

Then

$$l'(\theta|\mathbf{x}) = \sum_{l=1}^n h(x_l) - nh(\theta).$$

Because $h(z)$ is increasing, only one extrema θ^* exists, and

$$h(\theta^*) = n^{-1} \sum_{l=1}^n h(x_l).$$

Further, the inverse function h^{-1} exists, and we can continue that

$$\theta^* = h^{-1}(n^{-1} \sum_{l=1}^n h(x_l)).$$

The only remaining issue is to verify that the extrema is the maximum. Note that you cannot take the second derivative because $h(\theta)$ may not be differentiable (this is not

assumed). On the other hand, if h is differentiable then $l''(\theta|\mathbf{x}) = -nh'(\theta) < 0$ because $h(\theta)$ is increasing.

In general, please look at the above-written expression for $l'(\theta|\mathbf{x})$ and note that $l'(\theta|\mathbf{x})$ is negative for $\theta > \theta^*$ and $l'(\theta|\mathbf{x})$ is positive for $\theta < \theta^*$ because $h(\theta)$ is increasing function. This yields that the extrema is the maximum. Two graphics below highlight this conclusion.